

Scilab Textbook Companion for
Engineering Physics
by G. Aruldas¹

Created by
Pankaj Biswas
Modern Physics
Physics
Shri Mata Vaishno Devi University
College Teacher
Dr. Vinay Kumar
Cross-Checked by

July 31, 2019

¹Funded by a grant from the National Mission on Education through ICT, <http://spoken-tutorial.org/NMEICT-Intro>. This Textbook Companion and Scilab codes written in it can be downloaded from the "Textbook Companion Project" section at the website <http://scilab.in>

Book Description

Title: Engineering Physics

Author: G. Aruldas

Publisher: Phi Learning Pvt. Ltd.

Edition: 1

Year: 2010

ISBN: 9788120339163

Scilab numbering policy used in this document and the relation to the above book.

Exa Example (Solved example)

Eqn Equation (Particular equation of the above book)

AP Appendix to Example(Scilab Code that is an Appednix to a particular Example of the above book)

For example, Exa 3.51 means solved example 3.51 of this book. Sec 2.3 means a scilab code whose theory is explained in Section 2.3 of the book.

Contents

List of Scilab Codes	4
1 OSCILLATION AND WAVES	5
2 ELECTROMAGNETIC THEORY	12
3 INTERFERENCE	15
4 DIFFRACTION	21
5 POLARIZATION	28
6 CRYSTALLOGRAPHY	33
7 SUPERCONDUCTIVITY	44
8 SPECIAL THEORY OF RELATIVITY	48
9 QUANTUM MECHANICS	58
10 STATISTICAL MECHANICS	62
11 LASERS	65
12 HOLOGRAPHY AND FIBRE OPTICS	70

13 DIELECTRIC PROPERTIES OF MATERIALS	74
14 MAGNETIC PROPERTIES OF MATERIALS	80
15 THERMAL PROPERTIES	86
17 ULTRASONICS	90
18 ACOUSTICS OF BUILDINGS	91

List of Scilab Codes

Exa 1.1	Time period of SHM	5
Exa 1.2	Accelertion and maximum velocity in SHM .	5
Exa 1.3	Damped Vibrating System	6
Exa 1.4	Amplitude and Time Period in SHM	7
Exa 1.5	Oscillation of a spring mass system	8
Exa 1.6	Frequency of visible region	8
Exa 1.7	Characteristics of sound wave	9
Exa 1.8	Equation of a wave moving along X axis . .	10
Exa 1.9	Wave in the wire	10
Exa 2.1	Peak value of displacement current	12
Exa 2.2	Displacement current density in a good con- ductor	12
Exa 2.4	Poynting vector	13
Exa 2.5	Plane electromagnetic wave in a medium . .	13
Exa 3.1	Wavelength of Light using Young Double Slit experiment	15
Exa 3.2	Fringe shift due to change in wavelength . .	15
Exa 3.3	Refractive index from double slit experiment	16
Exa 3.4	Interference by thin soap film	17
Exa 3.5	Interference due to thin air wedge	17
Exa 3.6	Separation between consecutive bright fringes formed by an air wedge	18
Exa 3.7	Newton Rings by reflected light	18
Exa 3.8	Refractive index from Newton Rings arrange- ment	19
Exa 3.9	Wavelength of light using Michelson Interfer- ometer	19

Exa 3.10	Shift in movable mirror of Michelson Interferometer	20
Exa 4.1	Diffraction at a single slit	21
Exa 4.2	Diffraction at a circular aperture	22
Exa 4.3	Second order maximum for diffraction grating	23
Exa 4.4	The highest spectral order with diffraction grating	23
Exa 4.5	Overlapping spectra with diffraction grating	24
Exa 4.6	Width of first order spectrum	25
Exa 4.7	Resolution of wavelengths for grating	25
Exa 4.8	Angular separation to satisfy Rayleigh criterion	26
Exa 4.9	Linear separation between two points	27
Exa 5.1	Polarization by reflection	28
Exa 5.2	Percentage transmission of polarized light .	29
Exa 5.3	Thickness of Quarter Wave Plate	29
Exa 5.4	Behaviour of half wave plate for increased wavelength	30
Exa 5.5	Phase retardation for quartz	31
Exa 5.6	Brewster angle at the boundary between two materials	31
Exa 6.1	Lattice parameter of NaCl crystal	33
Exa 7.1	Variation of critical magnetic field with temperature	34
Exa 6.2	Miller indices of the crystal plane	34
Exa 6.3	Indices of lattice plane	35
Exa 6.5	Interplanar spacing in cubic crystal	35
Exa 6.6	Interplanar spacing in cubic crystal	36
Exa 6.7	ngle of reflection by using wavelength of X ray	37
Exa 6.8	Actual volume occupied by the spheres in fcc structure	38
Exa 6.9	X ray Diffraction by crystal planes	38
Exa 6.10	X ray Diffraction by crystal planes	39
Exa 6.11	Maximum order of diffraction	40
Exa 6.12	Bragg reflection angle for the second order diffraction	41
Exa 6.13	Distance between nearest neighbours of NaCl	42
Exa 6.14	Effect of structural change on volume	42

Exa 7.2	Frequency of Josephson current	44
Exa 7.3	Superconducting energy gap at 0K	44
Exa 7.4	Wavelength of photon to break up a Cooper pair	45
Exa 7.5	Variation of London penetration depth with temperature	45
Exa 7.6	Isotope Effect in mercury	46
Exa 8.1	Relativistic length contraction	48
Exa 8.2	Time Dilation	49
Exa 8.4	Relativistic velocity addition	49
Exa 8.5	Relativistic effects as observed for spaceship	50
Exa 8.6	Time difference and distance between the events	51
Exa 8.7	Speed of unstable particle in the Laboratory frame	52
Exa 8.8	Relativistic effects applied to mu meson . .	52
Exa 8.9	Speed of moving mass	53
Exa 8.10	Rate of decreasing mass of sun	54
Exa 8.11	Relativistic mass energy relation	54
Exa 8.13	Mass from relativistic energy	55
Exa 8.14	Relativistic momentum of high speed probe	56
Exa 8.15	Moving electron subjected to the electric field	56
Exa 9.1	De broglie wavelength of an electron from accelerating potential	58
Exa 9.2	De broglie wavelength of an electron from kinetic energy	58
Exa 9.4	Uncertainty principle for position and momentum	59
Exa 9.5	Uncertainty principle for energy and time .	60
Exa 9.6	Width of spectral line from Uncertainty principle	60
Exa 9.14	Probability of electron moving in 1D box . .	61
Exa 10.1	Ratio of occupancy of two states	62
Exa 10.4	Number density and fermi energy of silver .	62
Exa 10.5	Electronic contribution to the molar heat capacity of silver	63
Exa 10.6	Fermi energy and mean energy of aluminium	64
Exa 11.1	Ratio of spontaneous and stimulated emission	65
Exa 11.2	Energy of excited state of laser system . . .	66

Exa 11.3	Condition of equivalence of stimulated and spontaneous emission	66
Exa 11.4	Area and intensity of image formed by laser	67
Exa 11.5	Rate of energy released in a pulsed laser . . .	68
Exa 11.6	Angular and linear spread of laser beam . . .	68
Exa 12.1	Parameters of step index fibre	70
Exa 12.2	Parameters of optical fibre	71
Exa 12.3	Numerical aperture and acceptance angle of step index fibre	72
Exa 12.5	Output power in fibre optic communication	72
Exa 13.1	Electronic Polarizability of atom	74
Exa 13.2	Parallel plate capacitor	75
Exa 13.3	Dielectric displacement of medium	75
Exa 13.4	Relative dielectric constant	76
Exa 13.5	Atomic polarizability of sulphur	77
Exa 13.6	Electronic polarizability from refractive index	77
Exa 13.7	Ratio of electronic polarizability to ionic polarizability	78
Exa 14.1	Spontaneous magnetisation of the substance	80
Exa 14.2	Relative permeability of ferromagnetic material	81
Exa 14.3	Relative permeability from magnetisation . . .	81
Exa 14.4	Magnetic flux density and magnetisation of diamagnetic material	82
Exa 14.5	Magnetisation Magnetic flux density relative permeability of diamagnetic material	83
Exa 14.6	Mean radius of body centered cubic structure	83
Exa 14.7	Susceptibility and magnetisation of paramagnetic salt	84
Exa 15.1	Debye temperature of aluminium	86
Exa 15.2	Lattice specific heat of carbon	86
Exa 15.3	Einstein frequency for Cu	87
Exa 15.4	Electronic and lattice heat capacities for Cu	87
Exa 15.5	Einstein lattice specific heat	88
Exa 15.6	Molar electronic heat capacity of zinc	89
Exa 17.1	Thickness of vibrating quartz at resonance . .	90
Exa 18.1	Output power of the sound source	91
Exa 18.2	Change in sound level for doubling intensity	92

Exa 18.3	Total absorption of sound in the hall	92
Exa 18.4	Average absorption coefficient of the surfaces of the hall	92
Exa 18.5	Reverbration time for the hall	93
Exa 18.6	Gain of resultant sound intensity	94

Chapter 1

OSCILLATION AND WAVES

Scilab code Exa 1.1 Time period of SHM

```
1 // Scilab Code Ex1.1 : Page-23 (2010)
2 A = 4/2; // Amplitude of SHM, cm
3 x = 0; // Mean position of oscillating particle ,
  cm
4 v = 12; // Velocity of the particle at the mean
  position , cm/s
5 // As  $v = \omega \sqrt{A^2 - x^2}$ , solving for  $\omega$ 
6  $\omega = v / \sqrt{A^2 - x^2}$ ;
7 printf("\\nThe time period of SHM = %5.2f s", (2*%pi)
  / $\omega$ );
8
9 // Result
10 // The time period of SHM = 1.05 s
```

Scilab code Exa 1.2 Acceleration and maximum velocity in SHM

```
1 // Scilab Code Ex1.2 : Page-23 (2010)
2 T = 0.1; // Time period of oscillation in SHM, s
```

```

3 x = 0.2;    // Position of the particle from its
    mean position , cm
4 A = 4;     // Amplitude of the particle executing SHM
    , cm
5 // As  $T = 2*\%pi/\omega$ , solving for omega
6 omega = 2*%pi/T;    // Angular speed of particle
    executing SHM, per sec
7 a = omega^2*x;    // Accelertion of particle
    executing SHM, cm per sec square
8 v_max = omega*A;    // Maximum velocity of the
    particle in SHM, cm per sec
9 printf("\nThe accelertion of particle executing SHM
    = %5.1f cm per sec square", a);
10 printf("\nThe maximum velocity of the particle in
    SHM = %5.1f cm per sec", v_max);
11
12 // Result
13 // The accelertion of particle executing SHM = 789.6
    cm per sec square
14 // The maximum velocity of the particle in SHM =
    251.3 cm per sec

```

Scilab code Exa 1.3 Damped Vibrating System

```

1 // Scilab Code Ex1.3 : Page-24 (2010)
2 A1 = 40;    // First amplitude of oscillation , cm
3 An_plus_1 = 4;    // Amplitude after 100
    oscillations , cm
4 n = 100;    // Number of oscillations
5 T = 2.5;    // Time period of oscillations , s
6 t = T/4;    // Time taken to reach the first
    amplitude from the mean position , s
7 // Now  $A1 = x0*\exp(-\lambda*t)$  and  $An\_plus\_1 = x0*\exp$ 
     $(-\lambda*(t+nT))$ 
8 //  $A1/An\_plus\_1 = \exp(n*\lambda*T)$ , solving for

```

```

lambda
9 lambda = log(A1/An_plus_1)/(n*T);    // Damping
    constant. per sec
10 printf("\nDamping constant = %3.2e per sec", lambda)
    ;
11
12 // Result
13 // Damping constant = 9.21e-003 per sec

```

Scilab code Exa 1.4 Amplitude and Time Period in SHM

```

1 // Scilab Code Ex1.4 : Page-24 (2010)
2 v1 = 16;    // Velocity of particle executing SHM at
    position 3 cm
3 v2 = 12;    // Velocity of particle executing SHM at
    position 4 cm
4 x1 = 3;    // First position of the particle , cm
5 x2 = 4;    // Second position of the particle , cm
6 // As  $v = \omega \sqrt{A^2 - x^2}$  so
7 //  $(v1/v2)^2 = (A^2 - x1^2)/(A^2 - x2^2)$ , solving
    for A
8 A = poly(0, 'A');    // Declare variable A
9 A = roots((A^2 - x1^2)*v2^2 - (A^2 - x2^2)*v1^2);
10 printf("\nThe amplitude of SHM = %1d cm", A(1));
11 //  $v = \omega \sqrt{A^2 - x^2}$ , solving for omega
12 omega = v1/sqrt(A(1)^2 - x1^2);    // Angular speed
    of the particle , rad per sec
13 T = 2*pi/omega;    // Time period of oscillation ,
    sec
14 printf("\nThe time period of oscillation = %5.3f sec
    ", T);
15
16 // Result
17 // The amplitude of SHM = 5 cm
18 // The time period of oscillation = 1.571 sec

```

Scilab code Exa 1.5 Oscillation of a spring mass system

```
1 // Scilab Code Ex1.5 : Page-25 (2010)
2 m = 0.3; // Mass attached to the string , kg
3 g = 9.8; // Acceleration due to gravity , metre
    per sec square
4 x = 0.15; // Stretchness produced in the spring ,
    m
5 F = m*g; // Restoring force acting on the mass, N
6 k = F/x; // Spring constant, newton per metre
7 A = 0.1; // Amplitude of the string , m
8 omega = sqrt(k/m); // Angular frequency of
    oscillation , rad per sec
9 v0 = omega*A; // Maximum velocity during the
    oscillations , m/s
10 printf("\nThe spring constant = %4.1f newton per
    metre", k);
11 printf("\nThe amplitude of oscillation = %2.1f m", A
    );
12 printf("\nThe maximum velocity during oscillations =
    %3.2f m/s", v0);
13
14 // Result
15 // The spring constant = 19.6 newton per metre
16 // The amplitude of oscillation = 0.1 m
17 // The maximum velocity during oscillations = 0.81 m
    /s
```

Scilab code Exa 1.6 Frequency of visible region

```
1 // Scilab Code Ex1.6 : Page-25 (2010)
```

```

2 lambda1 = 400e-09;    // Lower limit of wavelength
   of visible region , m
3 lambda2 = 700e-09;    // Upper limit of wavelength
   of visible region , m
4 c = 3e+08;          // Speed of light in vacuum, m/s
5 f1 = c/lambda1;      // Upper limit of frequency of
   visible region , m
6 f2 = c/lambda2;      // Lower limit of frequency of
   visible region , m
7 printf("\nThe frequency equivalent of %3g nm to %3g
   nm is %3.1e Hz to %3.1e Hz", lambda1/1e-09,
   lambda2/1e-09, f1, f2);
8
9 // Result
10 // The frequency equivalent of 400 nm to 700 nm is
   7.5e+014 Hz to 4.3e+014 Hz

```

Scilab code Exa 1.7 Characteristics of sound wave

```

1 // Scilab Code Ex1.7 : Page-26 (2010)
2 // Comparing the standard equation
3 //  $u(x,t) = A \sin(2 * \pi(x/\lambda - t/T))$ 
4 // with the given equation, we get
5 A = 1.5e-03;        // Amplitude of the sound wave, m
6 lambda = 8;         // Wavelength of the sound wave, m
7 T = 1/40;          // Time period of the sound wave, s
8 nu = 1/T;          // Frequency of the sound wave, Hz
9 v = nu*lambda;     // Velocity of the sound wave, m/s
10 printf("\nThe amplitude of the sound wave = %3.1e m",
   , A);
11 printf("\nThe wavelength of the sound wave = %1d m",
   lambda);
12 printf("\nThe time period of the sound wave = %3.2 f
   s", T);
13 printf("\nThe frequency of the sound wave = %2d Hz",

```

```

    nu);
14 printf("\nThe velocity of the sound wave = %3d m/s",
    v);
15
16
17 // Result
18 // The amplitude of the sound wave = 1.5e-003 m
19 // The wavelength of the sound wave = 8 m
20 // The time period of the sound wave = 0.03 s
21 // The frequency of the sound wave = 40 Hz
22 // The velocity of the sound wave = 320 m/s

```

Scilab code Exa 1.8 Equation of a wave moving along X axis

```

1 // Scilab Code Ex1.8 : Page-26 (2010)
2 A = 2; // Amplitude of the wave, cm
3 T = 0.5; // Time period of the wave, sec
4 v = 200; // Wave velocity, cm/s
5 f = 1/0.5; // Frequency of the wave, Hz
6 lambda = v/f; // Wavelength of the wave, cm
7 printf("\nThe Equation of the wave moving along X-
    axis :");
8 printf("u = %1d*sin*2*pi*(x/%3d-t/%2.1 f)", A, lambda
    , T);
9
10
11 // Result
12 // The Equation of the wave moving along X-axis :u =
    2*sin*2*pi*(x/100-t/0.5)

```

Scilab code Exa 1.9 Wave in the wire

```

1 // Scilab Code Ex1.9 : Page-27 (2010)

```



```

2 T = 1000;    // Tension in the wire , N
3 m = 15/300; // Mass per unit length of the wire ,
    kg per metre
4 lambda = 0.30; // Wavelength of wave along wire ,
    m
5 v = sqrt(T/m); // Velocity of wave through wire ,
    m/s
6 nu = v/lambda; // Frequency of wave through
    string , Hz
7 printf("\nThe velocity and frequency of the wave
    through wire are %5.1f m/s and %5.1f Hz
    respectively" , v, nu);
8
9
10
11 // Result
12 // The velocity and frequency of the wave through
    wire are 141.4 m/s and 471.4 Hz respectively

```

Chapter 2

ELECTROMAGNETIC THEORY

Scilab code Exa 2.1 Peak value of displacement current

```
1 // Scilab Code Ex2.1 : Page-46 (2010)
2 function V = f(t)
3     V = 0.2*sin(120*%pi*t);
4 endfunction
5 t = 0; // Time when peak value of current occurs
6 C = 10e-012; // Capacitance of the capacitor ,
   farad
7 I = C*derivative(f,t);
8 printf("\nThe peak value of displacement current =
   %6.4e A", I);
9
10 // Result
11 // The peak value of displacement current = 7.5398e
   -010 A
```

Scilab code Exa 2.2 Displacement current density in a good conductor

```

1 // Scilab Code Ex2.2 : Page-46 (2010)
2 function E = fn(t)
3     E = sin(120*%pi*t);
4 endfunction
5 epsilon_r = 1; // Relative electrical
    permittivity of free space
6 epsilon_0 = 8.854e-012; // Absolute electrical
    permittivity of free space, farad per metre
7 t = 0; // Time when peak value of current occurs
8 J2 = epsilon_0*epsilon_r*derivative(fn,t);
9 printf("\nThe peak value of displacement current =
    %4.2e ampere per metre square", J2);
10
11 // Result
12 // The peak value of displacement current = 3.34e
    -009 ampere per metre square

```

Scilab code Exa 2.4 Poynting vector

```

1 // Scilab Code Ex2.4 : Page-47 (2010)
2 p = 60; // Power rating of bulb, watt
3 d = 0.5; // Distance from the blb, m
4 P = p/(4*%pi*d^2); // Value of Poynting vector,
    watt per metre square
5 printf("\nThe value of Poynting vector = %4.1f watt
    per metre square", P);
6
7 // Result
8 // The value of Poynting vector = 19.1 watt per
    metre square

```

Scilab code Exa 2.5 Plane electromagnetic wave in a medium

```

1 // Scilab Code Ex2.5 : Page-47 (2010)
2 E_peak = 6; // Peak value of electric field
   intensity , V/m
3 c = 3e+08; // Speed of electromagnetic wave in
   free space , m/s
4 mu_0 = 4*%pi*1e-07; // Absolute permeability of
   free space , tesla metre per ampere
5 epsilon_0 = 8.854e-012; // Absolute permittivity
   of free space , farad/m
6 mu_r = 1; // Relative permeability of medium
7 epsilon_r = 3; // Relative permittivity of the
   medium
8 v = c/sqrt(mu_r*epsilon_r); // Wave velocity , m/s
9 eta = sqrt((mu_0/epsilon_0)*(mu_r/epsilon_r)); //
   Intrinsic impedance of the medium, ohm
10 H_P = E_peak*sqrt((epsilon_0*epsilon_r)/(mu_0*mu_r))
   ; // Peak value of the magnetic intensity ,
   ampere per metre
11 printf("\nThe wave velocity = %5.3e m/s", v);
12 printf("\nThe intrinsic impedance of the medium = %6
   .2f ohm", eta);
13 printf("\nThe peak value of the magnetic intensity =
   %4.2e A/m", H_P);
14
15 // Result
16 // The wave velocity = 1.732e+008 m/s
17 // The intrinsic impedance of the medium = 217.51
   ohm
18 // The peak value of the magnetic intensity = 2.76e
   -002 A/m

```

Chapter 3

INTERFERENCE

Scilab code Exa 3.1 Wavelength of Light using Young Double Slit experiment

```
1 // Scilab Code Ex3.1 : Page-71 (2010)
2 beta = 0.51e-02; // Fringe width, cm
3 d = 2.2e-02; // Distance between the slits, cm
4 D = 2e+02; // Distance between the slits and the
   screen, cm
5 // As beta = D*lambda/d, solving for lambda
6 lambda = beta*d/D; // Wavelength of light, m
7 printf("\nThe wavelength of light = %4d angstrom",
   lambda/1e-010);
8
9 // Result
10 // The wavelength of light = 5610 angstrom
```

Scilab code Exa 3.2 Fringe shift due to change in wavelength

```
1 // Scilab Code Ex3.2 : Page-71 (2010)
2 lambda1 = 4250e-010; // First wavelength emitted
   by source of light, m
```

```

3 lambda2 = 5050e-010;    // Second wavelength emitted
    by source of light , m
4 D = 1.5;    // Distance between the source and the
    screen , m
5 d = 0.025e-03;    // Distance between the slits ,
    m
6 n = 3;    // Number of fringe from the centre
7 x3 = n*lambda1*D/d;    // Position of third bright
    fringe due to lambda1, m
8 x3_prime = n*lambda2*D/d;    // Position of third
    bright fringe due to lambda2, m
9 printf("\nThe separation between the third bright
    fringe due to the two wavelengths = %4.2f cm", (
    x3_prime - x3)/1e-02);
10
11 // Result
12 // The separation between the third bright fringe
    due to the two wavelengths = 1.44 cm

```

Scilab code Exa 3.3 Refractive index from double slit experiment

```

1 // Scilab Code Ex3.3 : Page-71 (2010)
2 lambda = 5.5e-05;    // Wavelength emitted by source
    of light , cm
3 n = 4;    // Number of fringes shifted
4 t = 3.9e-04;    // Thickness of the thin glass sheet
    , cm
5 mu = n*lambda/t+1;    // Refractive index of the
    sheet of glass
6 printf("\nThe refractive index of the sheet of glass
    = %6.4f", mu);
7
8 // Result
9 // The refractive index of the sheet of glass =
    1.5641

```

Scilab code Exa 3.4 Interference by thin soap film

```
1 // Scilab Code Ex3.4 : Page-72 (2010)
2 lambda = 5893e-010; // Wavelength of
   monochromatic light used, m
3 n = 1; // Number of fringe for the least
   thickness of the film
4 r = 0; // Value of refraction angle for normal
   incidence, degrees
5 mu = 1.42; // refractive index of the soap film
6 // As for constructive interference,
7 //  $2\mu t \cos(r) = (2n-1)\lambda/2$ , solving for t
8 t = (2*n-1)*lambda/(4*mu*cos(r)); // Thickness of
   the film that appears bright, m
9 printf("\nThe thickness of the film that appears
   bright = %6.1f angstrom", t/1e-010);
10 // As for destructive interference,
11 //  $2\mu t \cos(r) = n\lambda$ , solving for t
12 t = n*lambda/(2*mu*cos(r)); // Thickness of the
   film that appears bright, m
13 printf("\nThe thickness of the film that appears
   dark = %4d angstrom", t/1e-010);
14
15 // Result
16 // The thickness of the film that appears bright =
   1037.5 angstrom
17 // The thickness of the film that appears dark =
   2075 angstrom
```

Scilab code Exa 3.5 Interference due to thin air wedge

```

1 // Scilab Code Ex3.5 : Page-72 (2010)
2 lambda = 5893e-008; // Wavelength of
   monochromatic lihgt used, m
3 n = 10; // Number of fringe that are found in the
   distnace of 1 cm
4 d = 1; // Distance of 10 fringes , cm
5 beta = d/n; // Fringe width, cm
6 theta = lambda/(2*beta); // Angle of the wedge,
   rad
7 printf("\nThe angle of the wedge = %5.3e rad", theta
   );
8
9 // Result
10 // The angle of the wedge = 2.946e-004 rad

```

Scilab code Exa 3.6 Separation between consecutive bright fringes formed by an air

```

1 // Scilab Code Ex3.6 : Page-72 (2010)
2 lambda = 5900e-008; // Wavelength of
   monochromatic lihgt used, m
3 t = 0.010e-01; // Spacer thickness , cm
4 l = 10; // Wedge length, cm
5 theta = t/l; // Angle of the wedge, rad
6 beta = lambda/(2*theta); // Fringe width, cm
7 printf("\nThe separation between consecutive bright
   fringes = %5.3e cm", beta);
8
9 // Result
10 // The separation between consecutive bright fringes
   = 2.950e-001 cm

```

Scilab code Exa 3.7 Newton Rings by reflected light


```

1 // Scilab Code Ex3.7 : Page-72 (2010)
2 D4 = 0.4; // Diameter of 4th dark ring , cm
3 D12 = 0.7; // Diameter of 12th dark ring , cm
4 // We have  $dn\_puls\_k^2 - Dn^2 = 4*k*R*\lambda$ , so
5 //  $D12^2 - D4^2 = 32*R*\lambda$  and  $D20^2 - D12^2 = 32*R*$ 
//  $\lambda$  for  $k = 8$ , solving for D20
6 D20 = sqrt(2*D12^2-D4^2); // Diameter of 20th
// dark ring , cm
7 printf("\nThe diameter of 20th dark ring = %6.4f cm"
, D20);
8
9 // Result
10 // The diameter of 20th dark ring = 0.9055 cm

```

Scilab code Exa 3.8 Refractive index from Newton Rings arrangement

```

1 // Scilab Code Ex3.8 : Page-73 (2010)
2 Dn = 0.30; // Diameter of nth dark ring with air
// film , cm
3 dn = 0.25; // Diameter of nth dark ring with
// liquid film , cm
4 mu = (Dn/dn)^2; // Refractive index of the liquid
5 printf("\nThe refractive index of the liquid = %4.2f
", mu);
6
7 // Result
8 // The refractive index of the liquid = 1.44

```

Scilab code Exa 3.9 Wavelength of light using Michelson Interferometer

```

1 // Scilab Code Ex3.9 : Page-73 (2010)
2 x = 0.002945; // Distance through which movable
// mirror is shifted , cm

```

```

3 N = 100;      // Number of fringes shifted
4 lambda = 2*x/N;    // Wavelength of light , m
5 printf("\nThe wavelength of light = %4d angstrom",
        lambda/1e-008);
6
7 // Result
8 // The wavelength of light = 5890 angstrom

```

Scilab code Exa 3.10 Shift in movable mirror of Michelson Interferometer

```

1 // Scilab Code Ex3.10 : Page-73 (2010)
2 lambda1 = 5896e-008;    // Wavelength of D1 line of
        sodium , m
3 lambda2 = 5890e-008;    // Wavelength of D2 line of
        sodium , m
4 lambda = (lambda1+lambda2)/2;
5 // As lambda1 - lambda2 = lambda^2/(2*x), solving
        for x
6 x = lambda^2/(2*(lambda1 - lambda2));    // Shift in
        movable mirror of Michelson Interferometer , cm
7 printf("\nThe shift in movable mirror = %5.3f mm", x
        /1e-001);
8
9 // Result
10 // The shift in movable mirror = 0.289 mm

```

Chapter 4

DIFFRACTION

Scilab code Exa 4.1 Diffraction at a single slit

```
1 // Scilab Code Ex4.1 : Page-91 (2010)
2 D = 50; // Distance between source and the screen
   , cm
3 lambda = 6563e-008; // Wavelength of light of
   parallel rays , m
4 d = 0.385e-01; // Width of the slit , cm
5 n = 1; // Order of diffraction for first minimum
6 // As  $\sin(\theta_1) = n*\lambda/d = x_1/D$ , solving for
   x1
7 x1 = n*lambda*D/d; // Distance from the centre of
   the principal maximum to the first minimum, cm
8 printf("\nThe Distance from the centre of the
   principal maximum to the first minimum = %4.2 f mm
   ", x1/1e-001);
9 n = 5; // Order of diffraction for fifth minimum
10 x2 = n*lambda*D/d; // Distance from the centre of
   the principal maximum to the fifth minimum, cm
11 printf("\nThe Distance from the centre of the
   principal maximum to the fifth minimum = %4.2 f mm
   ", x2/1e-001);
12
```

```

13 // Result
14 // The Distance from the centre of the principal
    maximum to the first minimum = 0.85 mm
15 // The Distance from the centre of the principal
    maximum to the fifth minimum = 4.26 mm

```

Scilab code Exa 4.2 Diffraction at a circular aperture

```

1 // Scilab Code Ex4.2 : Page-91 (2010)
2 D = 0.04; // Diameter of circular aperture , cm
3 f = 20; // Focal length of convex lens , cm
4 lambda = 6000e-008; // Wavelength of light used ,
    m
5 // We have  $\sin(\theta) = 1.22*\lambda/D = \theta$  , for
    small  $\theta$  , such that
6 // For first dark ring
7 theta = 1.22*lambda/D; // The half angular width
    at central maximum, rad
8 r1 = theta*f; // The half width of central
    maximum for first dark ring , cm
9 // We have  $\sin(\theta) = 5.136*\lambda/(\pi*D) = \theta$ 
    , for small  $\theta$  , such that
10 // For second dark ring
11 theta = 5.136*lambda/(%pi*D); // The half angular
    width at central maximum, rad
12 r2 = theta*f; // The half width of central
    maximum for second dark ring , cm
13 printf("\nThe radius of first dark ring = %4.2e cm",
    r1);
14 printf("\nThe radius of second dark ring = %4.1e cm"
    , r2);
15
16 // Result
17 // The radius of first dark ring = 3.66e-002 cm
18 // The radius of second dark ring = 4.90e-002 cm

```

Scilab code Exa 4.3 Second order maximum for diffraction grating

```
1 // Scilab Code Ex4.3 : Page-91 (2010)
2 n = 2; // Order of diffraction
3 lambda = 650e-009; // Wavelength of light used, m
4 d = 1.2e-05; // Distance between two consecutive
  slits of grating, m
5 // We have  $\sin(\theta) = n\lambda/d$ , solving for theta
6 theta = asind(n*lambda/d); // Angle at which the
  650 nm light produces a second order maximum,
  degrees
7 printf("\nThe angle at which the 650 nm light
  produces a second order maximum = %4.2f degrees",
  theta);
8
9 // Result
10 // The angle at which the 650 nm light produces a
  second order maximum = 6.22 degrees
```

Scilab code Exa 4.4 The highest spectral order with diffraction grating

```
1 // Scilab Code Ex4.4 : Page-92 (2010)
2 lambda = 650e-009; // Wavelength of light used, m
3 N = 6000e+02; // Number of lines per m on grating
  , per m
4 theta = 90; // Angle at which the highest
  spectral order is obtained, degrees
5 // We have  $\sin(\theta) = n\lambda/d$ , solving for n
6 n = sind(theta)/(N*lambda); // The highest order
  of spectra with diffraction grating
```

```

7 printf("\nThe highest order of spectra obtained with
    diffraction grating = %ld", n);
8
9 // Result
10 // The highest order of spectra obtained with
    diffraction grating = 2

```

Scilab code Exa 4.5 Overlapping spectra with diffraction grating

```

1 // Scilab Code Ex4.5 : Page-92 (2010)
2 N = 4000e+02; // Number of lines per m on grating
    , per m
3 // For Blue Line
4 lambda = 450e-009; // Wavelength of blue light , m
5 n = 3; // Order of diffraction spectrum
6 // We have sin(theta) = n*N*lambda, solving for sin(
    theta)
7 sin_theta_3 = n*N*lambda; // Sine of angle at
    third order diffraction
8 // For Red Line
9 lambda = 700e-009; // Wavelength of blue light , m
10 n = 2; // Order of diffraction spectrum
11 // We have sin(theta) = n*N*lambda, solving for sin(
    theta)
12 sin_theta_2 = n*N*lambda; // Sine of angle at
    second order diffraction
13 // Check for overlapping
14 if abs(sin_theta_3 - sin_theta_2) < 0.05 then
15     printf("\nThe two orders overlap.");
16 else
17     printf("\nThe two orders do not overlap.");
18 end
19
20 // Result
21 // The two orders overlap.

```

Scilab code Exa 4.6 Width of first order spectrum

```
1 // Scilab Code Ex4.6 : Page-93 (2010)
2 n = 1; // Order of diffraction spectrum
3 N = 6000e+02; // Number of lines per m on
  diffraction grating, per m
4 D = 2; // Distance of screen from the source, m
5 lambda1 = 400e-009; // Wavelength of blue light,
  m
6 // We have  $\sin(\theta_1) = n \cdot N \cdot \lambda$ , solving for
  theta1
7 theta1 = asind(n*N*lambda1); // Angle at first
  order diffraction for Blue light, degrees
8 lambda2 = 750e-009; // Wavelength of blue light,
  m
9 // We have  $\sin(\theta_2) = n \cdot N \cdot \lambda$ , solving for
  theta2
10 theta2 = asind(n*N*lambda2); // Angle at first
  order diffraction for Red light, degrees
11 x1 = D*tand(theta1); // Half width position at
  central maximum for blue color, m
12 x2 = D*tand(theta2); // Half width position at
  central maximum for red color, m
13
14 printf("\nThe width of first order spectrum on the
  screen = %4.1f cm", (x2 - x1)/1e-02);
15
16 // Result
17 // The width of first order spectrum on the screen =
  51.3 cm
```

Scilab code Exa 4.7 Resolution of wavelengths for grating

```

1 // Scilab Code Ex4.7 : Page-93 (2010)
2 w = 5; // Width of the grating , cm
3 N = 320; // Number of lines per cm on grating ,
    per cm
4 N0 = w*N; // Total number of lines on the grating
5 lambda = 640; // Wavelength of light , nm
6 n = 2; // Order of diffraction
7 d_lambda = lambda/(n*N0); // Separation between
    wavelengths which the gratign can just resolve ,
    nm
8 printf("\nThe separation between wavelengths which
    the grating can just resolve = %3.1f nm",
    d_lambda);
9
10 // Result
11 // The separation between wavelengths which the
    grating can just resolve = 0.2 nm

```

Scilab code Exa 4.8 Angular separation to satisfy Rayleigh criterion

```

1 // Scilab Code Ex4.8 : Page-93 (2010)
2 lambda = 550e-09; // Wavelength of light , m
3 D = 3.2e-02; // Diameter of circular lens , m
4 f = 24e-02; // Focal length of the lens , m
5 theta_min = 1.22*lambda/D; // Minimum angle of
    resolution provided by the lens , rad
6 // As delta_x/f = theta_min , solving for delta_x
7 delta_x = theta_min*f; // Separation of the
    centres of the images in the focal plane of lens ,
    m
8 printf("\nThe separation of the centres of the
    images in the focal plane of lens = %1d micro-
    metre", delta_x/1e-06);
9
10 // Result

```



```
11 // The separation of the centres of the images in
    the focal plane of lens = 5 micro-metre
```

Scilab code Exa 4.9 Linear separation between two points

```
1 // Scilab Code Ex4.9 : Page-94 (2010)
2 lambda = 550e-09; // Wavelength of light , m
3 D = 20e-02; // Diameter of objective of telescope
  , m
4 d = 6e+003; // Distance of two points from the
  objective of telescope , m
5 theta = 1.22*lambda/D; // Angular separation
  between two points , rad
6 x = theta*d; // Linear separation between two
  points , m
7 printf("\nThe linear separation between two points =
  %5.2f mm", x/1e-03);
8
9 // Result
10 // The linear separation between two points = 20.13
  mm
```

Chapter 5

POLARIZATION

Scilab code Exa 5.1 Polarization by reflection

```
1 // Scilab Code Ex5.1 : Polarization by reflection:
   Page-113 (2010)
2 mu_g = 1.72; // Refractive index of glass
3 mu_w = 4/3; // Refractive index of water
4 // For polarization to occur on flint glass,  $\tan(i)$ 
   = mu_g/mu_w
5 // Solving for i
6 i = atand(mu_g/mu_w);
7 printf("\nThe angle of incidence for complete
   polarization to occur on flint glass = %4.1f
   degrees", i);
8 // For polarization to occur on water,  $\tan(i) = \mu_w$ 
   /mu_g
9 // Solving for i
10 i = atand(mu_w/mu_g);
11 printf("\nThe angle of incidence for complete
   polarization to occur on water = %5.2f degrees",
   i);
12
13 // Result
14 // The angle of incidence for complete polarization
```

```
to occur on flint glass = 52.2 degrees
15 // The angle of incidence for complete polarization
to occur on water = 37.78 degrees
```

Scilab code Exa 5.2 Percentage transmission of polarized light

```
1 // Scilab Code Ex5.2 : Percentage transmission of
polarized light: Page-113 (2010)
2 I0 = 1; // For simplicity, we assume the
intensity of light falling on the second Nicol
prism to be unity, watt per metre square
3 theta = 30; // Angle through which the crossed
Nicol is rotated, degrees
4 I = I0*cosd(90-theta)^2; // Intensity of the
emerging light from second Nicol, watt per metre
square
5 T = I/(2*I0)*100; // Percentage transmission of
incident light
6 printf("\nThe percentage transmission of incident
light after emerging through the Nicol prism = %4
.f percent", T);
7
8 // Result
9 // The percentage transmission of incident light
after emerging through the Nicol prism = 12.5
percent
```

Scilab code Exa 5.3 Thickness of Quarter Wave Plate

```
1 // Scilab Code Ex5.3 : Thickness of Quarter Wave
Plate : Page-113 (2010)
2 lambda = 6000e-008; // Wavelength of incident
light, cm
```

```

3 mu_e = 1.55;    // Refractive index of extraordinary
    ray
4 mu_o = 1.54;    // Refractive index of ordinary ray
5 t = lambda/(4*(mu_e - mu_o));    // Thickness of
    Quarter Wave plate of positive crystal, cm
6 printf("\nThe thickness of Quarter Wave plate = %6.4
    f cm", t);
7
8 // Result
9 // The thickness of Quarter Wave plate = 0.0015 cm

```

Scilab code Exa 5.4 Behaviour of half wave plate for increased wavelength

```

1 // Scilab Code Ex5.4 : Behaviour of half wave plate
    for increased wavelength : Page-114 (2010)
2 lambda = 1;    // For simplicity, wavelength of
    incident light is assumed to be , cm
3 mu_e = 1.55;    // Refractive index of extraordinary
    ray
4 mu_o = 1.54;    // Refractive index of ordinary ray
5 t = lambda/(2*(mu_e - mu_o));    // Thickness of
    Half Wave plate for given lambda, cm
6 t_prime = 2*lambda/(2*(mu_e - mu_o));    // Thickness
    of Half Wave plate for twice lambda, cm
7 printf("\nThe thickness of half wave plate is %2.1f
    times that of the quarter wave plate.", t/t_prime
    );
8 printf("\nThe half wave plate behaves as a quarter
    wave plate for twice the wavelength of incident
    light.");
9
10 // Result
11 // The thickness of half wave plate is 0.5 times
    that of the quarter wave plate.
12 // The half wave plate behaves as a quarter wave

```

plate for twice the wavelength of incident light.

Scilab code Exa 5.5 Phase retardation for quartz

```
1 // Scilab Code Ex5.5 : Phase retardation for quartz
  : Page-114 (2010)
2 lambda = 500e-09; // Wavelength of incident light
  , m
3 mu_e = 1.5508; // Refractive index of
  extraordinary ray
4 mu_o = 1.5418; // Refractive index of ordinary
  ray
5 t = 0.032e-03; // Thickness of quartz plate , m
6 dx = (mu_e - mu_o)*t; // Path difference between
  E-ray and O-ray , m
7 dphi = (2*pi)/lambda*dx; // Phase retardation
  for quartz for given wavelength , rad
8 printf("\nThe phase retardation for quartz for given
  wavelength = %5.3f pi rad", dphi/pi);
9
10 // Result
11 // The phase retardation for quartz for given
  wavelength = 1.152 pi rad
```

Scilab code Exa 5.6 Brewster angle at the boundary between two materials

```
1 // Scilab Code Ex5.6 : Brewster angle at the
  boundary between two materials : Page-114 (2010)
2 C = 52; // Critical angle for total internal
  reflection at a boundary between two materials ,
  degrees
3 // From Brewster 's law , tand(i_B) = 1_mu_2
4 // Also sind(C) = 1_mu_2 , so that
```

```
5 // tand(i_B) = sind(C), solving for i_B
6 i_B = atand(sind(C)); // Brewster angle at the
  boundary, degrees
7 printf("\nThe Brewster angle at the boundary between
  two materials = %2d degrees", i_B);
8
9 // Result
10 // The Brewster angle at the boundary between two
  materials = 38 degrees
```

Chapter 6

CRYSTALLOGRAPHY

Scilab code Exa 6.1 Lattice parameter of NaCl crystal

```
1 // Scilab Code Ex6.1 : Lattice parameter of NaCl
  crystal : Page-134 (2010)
2 M = 23+35.5; // Molecular weight of NaCl, kg
  per k-mole
3 d = 2.18e+03; // Density of rock salt, kg per
  metre cube
4 n = 4; // No. of atoms per unit cell for an fcc
  lattice of NaCl crystal
5 N = 6.023D+26; // Avogadro's No., atoms/k-mol
6 // Volume of the unit cell is given by
7 //  $a^3 = M*n/(N*d)$ 
8 // Solving for a
9 a = (n*M/(d*N))^(1/3); // Lattice constant of
  unit cell of NaCl
10 printf("\nLattice parameter for the NaCl crystal =
  %4.2f angstrom", a/1e-010);
11
12 // Result
13 // Lattice parameter for the NaCl crystal = 5.63
  angstrom
```

Scilab code Exa 7.1 Variation of critical magnetic field with temperature

```
1 // Scilab Code Ex7.1 : Variation of critical
   magnetic field with temperature : Page-152 (2010)
2 T_c = 3.722; // Critical temperature of
   superconducting transition , kelvin
3 B_c0 = 0.0306; // Critical magnetic field to
   destroy superconductivity , tesla
4 T = 2; // Temperature at which critical magnetic
   field is to be found out , kelvin
5 B_cT = B_c0*(1-(T/T_c)^2);
6 printf("\nThe critical magnetic field at %d K = %6.4
   f T", T, B_cT);
7
8 // Result
9 // The critical magnetic field at 2 K = 0.0218 T s
```

Scilab code Exa 6.2 Miller indices of the crystal plane

```
1 // Scilab Code Ex6.2 : Miller indices of the crystal
   plane : Page-134 (2010)
2 m = 3; n = 2; p = 1; // Coefficients of intercepts
   along three axes
3 m_inv = 1/m; // Reciprocate the first
   coefficient
4 n_inv = 1/n; // Reciprocate the second
   coefficient
5 p_inv = 1/p; // Reciprocate the third
   coefficient
6 mul_fact = double(lcm(int32([m,n,p]))); // Find l.c.
   m. of m,n and p
7 m1 = m_inv*mul_fact; // Clear the first fraction
```



```

8 m2 = n_inv*mul_fact;    // Clear the second fraction
9 m3 = p_inv*mul_fact;    // Clear the third fraction
10 printf("\nThe required miller indices are : (%d %d
    %d) ", m1,m2,m3);
11
12 // Result
13 // The required miller indices are : (2 3 6)

```

Scilab code Exa 6.3 Indices of lattice plane

```

1 // Scilab Code Ex6.3 : Indices of lattice plane :
    Page-135 (2010)
2 m = 2; // Coefficient of intercept along x-axis
3 n = %inf; // Coefficient of intercept along y-
    axis
4 p = 3/2; // Coefficient of intercept along z-axis
5 m_inv = 1/m; // Reciprocate m
6 n_inv = 1/n; // Reciprocate n
7 p_inv = 1/p; // Reciprocate p
8 mul_fact = 6; // multiplicative factor , L.C.M. of
    2 and 3 i.e. 6
9 m1 = m_inv*mul_fact; // Clear the first fraction
10 m2 = n_inv*mul_fact; // Clear the second fraction
11 m3 = p_inv*mul_fact; // Clear the third fraction
12 printf("\nThe required miller indices are : %d, %d,
    %d ", m1,m2,m3);
13
14 // Result
15 // The required miller indices are : 3, 0, 4

```

Scilab code Exa 6.5 Interplanar spacing in cubic crystal

```

1 // Scilab Code Ex6.5 : Interplanar spacing in cubic
   crystal: Page-136 (2010)
2
3 // For (110) planes
4 h = 1; k = 1; l = 0; // Miller Indices for planes in
   a cubic crystal
5 a = 0.43e-009; // Interatomic spacing, m
6 d = a/(h^2+k^2+l^2)^(1/2); // The interplanar
   spacing for cubic crystals, m
7 printf("\nThe interplanar spacing between
   consecutive (110) planes = %4.2f angstrom", d/1e
   -010);
8
9 // For (212) planes
10 h = 2; k = 1; l = 2; // Miller Indices for planes in
   a cubic crystal
11 a = 4.21D-10; // Interatomic spacing, m
12 d = a/(h^2+k^2+l^2)^(1/2); // The interplanar
   spacing for cubic crystals, m
13 printf("\nThe interplanar spacing between
   consecutive (212) planes = %4.3f angstrom", d/1e
   -010);
14
15 // Result
16 // The interplanar spacing between consecutive (110)
   planes = 3.04 angstrom
17 // The interplanar spacing between consecutive (212)
   planes = 1.403 angstrom

```

Scilab code Exa 6.6 Interplanar spacing in cubic crystal

```

1 // Scilab Code Ex6.6 : Interplanar spacing in cubic
   crystal: Page-136 (2010)
2 h = 2; k = 3; l = 1; // Miller Indices for planes in
   a cubic crystal

```

```

3 r = 0.175e-009;    // Atomic radius of fcc lattice ,
  m
4 a = 2*sqrt(2)*r;    // Interatomic spacing of fcc
  lattice , m
5 d = a/(h^2+k^2+l^2)^(1/2); // The interplanar
  spacing for cubic crystals , m
6 printf("\nThe interplanar spacing between
  consecutive (231) planes = %4.2f ansgtrom", d/1e
  -010);
7
8 // Result
9 // The interplanar spacing between consecutive (231)
  planes = 1.32 ansgtrom

```

Scilab code Exa 6.7 ngle of reflection by using wavelength of X ray

```

1 // Scilab Code Ex6.7 : Angle of reflection by using
  wavelength of X-ray: Page-136 (2010)
2 lambda = 1.440e-010; // Wavelength of X-rays , m
3 d = 2.8e-010; // Interplanar spacing of rocksalt
  crystal , m
4 // 2*d*sin(theta) = n*lambda **Bragg's law, n is
  the order of diffraction
5 // Solving for theta, we have
6
7 // For Ist Order diffraction
8 n = 1;
9 theta = asind(n*lambda/(2*d)); // Angle of
  diffraction , degrees
10 printf("\nThe angle of reflection for first order
  diffraction = %4.1f degrees", theta);
11
12 // For IInd Order diffraction
13 n = 2;
14 theta = asind(n*lambda/(2*d)); // Angle of

```

```

    diffraction , degrees
15 printf("\nThe angle of reflection for first order
    diffraction = %4.1f degrees", theta);
16
17 // Result
18 // The angle of reflection for first order
    diffraction = 14.9 degrees
19 // The angle of reflection for first order
    diffraction = 30.9 degrees

```

Scilab code Exa 6.8 Actual volume occupied by the spheres in fcc structure

```

1 // Scilab Code Ex6.8 : Actual volume occupied by the
    spheres in fcc structure Page-136 (2010)
2 N = 8*1/8 + 6*1/2; // total number of spheres in
    a unit cell
3 a = 1; // For convenience, assume interatomic
    spacing to be unity, m
4 r = a/(2*sqrt(2)); // The atomic radius, m
5 V_atom = N*4/3*pi*r^3; // Volume of atoms, metre
    cube
6 V_uc = a^3; // Volume of unit cell, metre cube
7 printf("\nThe percentage of actual volume occupied
    by the spheres in fcc structure = %4.2f percent",
    V_atom/V_uc*100);
8
9 // Result
10 // The percentage of actual volume occupied by the
    spheres in fcc structure = 74.05 percent

```

Scilab code Exa 6.9 X ray Diffraction by crystal planes

```

1 // Scilab Code Ex6.9 : X-ray Diffraction by crystal
  planes: Page-137 (2010)
2 // For (221) planes
3 h = 2; k = 2; l = 1; // Miller Indices for planes in
  a cubic crystal
4 a = 2.68e-010; // Interatomic spacing, m
5 n = 1; // First Order of diffraction
6 theta = 8.5; // Glancing angle at which Bragg's
  reflection occurs, degrees
7 d = a/(h^2+k^2+l^2)^(1/2); // The interplanar
  spacing for cubic crystal, m
8 lambda = 2*d*sind(theta); // Bragg's Law for
  wavelength of X-rays, m
9 n = 2; // Second order of diffraction
10 theta = asind(n*lambda/(2*d)); // Angle at which
  second order Bragg reflection occurs, degrees
11 printf("\nThe interplanar spacing between
  consecutive (221) planes = %5.3e", d);
12 printf("\nThe wavelength of X-rays = %5.3f angstrom"
  , lambda/1e-010);
13 printf("\nThe angle at which second order Bragg
  reflection occurs = %4.1f degrees", theta);
14
15 // Result
16 // The interplanar spacing between consecutive (221)
  planes = 8.933e-011
17 // The wavelength of X-rays = 0.264 angstrom
18 // The angle at which second order Bragg reflection
  occurs = 17.2 degrees

```

Scilab code Exa 6.10 X ray Diffraction by crystal planes

```

1 // Scilab Code Ex6.10 : Lattice parameter for (110)
  planes of cubic crystal: Page-137 (2010)
2 h = 1; k = 1; l = 0; // Miller Indices for planes in

```

```

    a cubic crystal
3 n = 1;    // First Order of diffraction
4 theta = 25;    // Glancing angle at which Bragg's
    reflection occurs, degrees
5 lambda = 0.7e-010;    // Wavelength of X-rays, m
6 // From Bragg's Law, n*lambda = 2*d*sind(theta),
    solving for d
7 d = n*lambda/(2*sind(theta));    // Interplanar
    spacing of cubic crystal, m
8 a = d*(h^2+k^2+l^2)^(1/2);    // The lattice parameter
    for cubic crystal, m
9 printf("\nThe lattice parameter for cubic crystal =
    %4.2f angstrom", a/1e-010);
10
11 // Result
12 // The lattice parameter for cubic crystal = 1.17
    angstrom

```

Scilab code Exa 6.11 Maximum order of diffraction

```

1 // Scilab Code Ex6.11 : Maximum order of diffraction
    : Page-138 (2010)
2 d = 0.31e-009;    // Interplanar spacing, m
3 n = 1;    // First Order of diffraction
4 theta = 9.25;    // Glancing angle at which Bragg's
    reflection occurs, degrees
5 // From Bragg's Law, n*lambda = 2*d*sind(theta),
    solving for lambda
6 lambda = 2*d*sind(theta)/n;    // Wavelength of X-
    rays, m (Bragg's Law)
7 theta_max = 90;    // Maximum possible angle at
    which reflection can occur, degrees
8 n = 2*d*sind(theta_max)/lambda;    // Maximum
    possible order of diffraction
9 printf("\nThe Maximum possible order of diffraction

```

```

    = %1d",n);
10
11 // Result
12 // The Maximum possible order of diffraction = 6

```

Scilab code Exa 6.12 Bragg reflection angle for the second order diffraction

```

1 // Scilab Code Ex6.12 : Bragg reflection angle for
  the second order diffraction: Page-138 (2010)
2 // For (110) planes
3 h = 1, k = 1, l = 0; // Miller indices for (110)
  planes
4 d_110 = 0.195e-009; // Interplanar spacing
  between (110) planes, m
5 // As  $d_{110} = a / (h^2 + k^2 + l^2)^{1/2}$ , solving for
  a
6 a = d_110*(h^2 + k^2 + l^2)^(1/2); // Lattice
  parameter for bcc crystal, m
7 // For (210) planes
8 h = 2, k = 1, l = 0; // Miller indices for (110)
  planes
9 d_210 = a/(h^2 + k^2 + l^2)^(1/2); // Interplanar
  spacing between (210) planes, m
10 n = 2; // Seconds Order of diffraction
11 lambda = 0.072e-009; // Wavelength of X-rays, m
12 // From Bragg's Law,  $n*\lambda = 2*d_{210}*\sin(\theta)$ ,
  solving for theta
13 theta = asind(n*lambda/(2*d_210)); // Bragg
  reflection angle for the second order diffraction
  , degrees
14 printf("\nBragg reflection angle for the second
  order diffraction = %5.2f degrees", theta);
15
16 // Result
17 // Bragg reflection angle for the second order

```

diffraction = 35.72 degrees

Scilab code Exa 6.13 Distance between nearest neighbours of NaCl

```
1 // Scilab Code Ex6.13 : Distance between nearest
  neighbours of NaCl: Page-138 (2010)
2 M = 23+35.5; // Molecular weight of NaCl, kg
  per k-mole
3 d = 2.18e+03; // Density of rock salt, kg per
  metre cube
4 n = 4; // No. of atoms per unit cell for an fcc
  lattice of NaCl crystal
5 N = 6.023D+26; // Avogadro's No., atoms/k-mol
6 // Volume of the unit cell is given by
7 // a^3 = M*n/(N*d)
8 // Solving for a
9 a = (n*M/(d*N))^(1/3); // Lattice constant of
  unit cell of NaCl
10 printf("\nThe distance between nearest neighbours of
  NaCl structure = %5.3e", a/2);
11
12 // Result
13 // The distance between nearest neighbours of NaCl
  structure = 2.814e-010
```

Scilab code Exa 6.14 Effect of structural change on volume

```
1 // Scilab Code Ex6.14 : Effect of structural change
  on volume : Page-139 (2010)
2 // For bcc structure
3 r = 1.258e-010; // Atomic radius of bcc structure
  of iron, m
```



```

4 a = 4*r/sqrt(3);    // Lattice parameter of bcc
   structure of iron , m
5 V = a^3;          // Volume of bcc unit cell , metre cube
6 N = 2;           // Number of atoms per unit cell in bcc
   structure
7 V_atom_bcc = V/N;    // Volume occupied by one atom ,
   metre cube
8 // For fcc structure
9 r = 1.292e-010;    // Atomic radius of fcc structure
   of iron , m
10 a = 2*sqrt(2)*r;    // Lattice parameter of fcc
   structure of iron , m
11 V = a^3;          // Volume of fcc unit cell , metre cube
12 N = 4;           // Number of atoms per unit cell in fcc
   structure
13 V_atom_fcc = V/N;    // Volume occupied by one atom ,
   metre cube
14 delta_V = (V_atom_bcc-V_atom_fcc)/V_atom_bcc*100;
   // Percentage change in volume due to
   structural change of iron
15 printf("\nThe percentage change in volume of iron =
   %4.2f percent", delta_V);
16
17 // Result
18 // The percentage change in volume of iron = 0.49
   percent

```

Chapter 7

SUPERCONDUCTIVITY

Scilab code Exa 7.2 Frequency of Josephson current

```
1 // Scilab Code Ex7.2 : Frequency of Josephson
  current : Page-152 (2010)
2 V = 1e-06; // DC voltage applied across the
  Josephson junction , volt
3 e = 1.6e-019; // Charge on an electron , C
4 h = 6.626e-034; // Planck's constant , Js
5 f = 2*e*V/h; // Frequency of Josephson current ,
  Hz
6 printf("\nThe frequency of Josephson current = %5.1 f
  MHz", f/1e+06);
7
8 // Result
9 // The frequency of Josephson current = 482.9 MHz
```

Scilab code Exa 7.3 Superconducting energy gap at 0K

```
1 // Scilab Code Ex7.3 : Superconducting energy gap at
  0K : Page-152 (2010)
```

```

2 T_c = 0.517;    // Critical temperature for cadmium,
  K
3 k = 1.38e-023; // Boltzmann constant, J/K
4 e = 1.6e-019;  // Energy equivalent of 1 eV, J/eV
5 E_g = 3.5*k*T_c/e; // Superconducting energy gap
  at absolute zero, eV
6 printf("\nThe superconducting energy gap for Cd at
  absolute zero = %4.2e eV",E_g);
7
8 // Result
9 // The superconducting energy gap for Cd at absolute
  zero = 1.56e-004 eV

```

Scilab code Exa 7.4 Wavelength of photon to break up a Cooper pair

```

1 // Scilab Code Ex7.4 : Wavelength of photon to break
  up a Cooper-pair: Page-152 (2010)
2 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
3 c = 3e+08;    // Speed of light in free space, m/s
4 h = 6.626e-034; // Planck's constant, Js
5 E_g = 1.5e-004; // Superconducting energy gap for
  a material, eV
6 // As  $E_g = h*f = h*c/\lambda$ , solving for  $\lambda$ 
7  $\lambda = h*c/(E_g*e)$ ; // Wavelength of photon to
  break up a Cooper-pair, m
8 printf("\nThe wavelength of photon to break up a
  Cooper-pair = %4.2e m", lambda);
9
10 // Result
11 // The wavelength of photon to break up a Cooper-
  pair = 8.28e-003 m

```

Scilab code Exa 7.5 Variation of London penetration depth with temperature

```

1 // Scilab Code Ex7.5: Variation of London
  penetration depth with temperature: Page-153
  (2010)
2 lambda_0 = 37e-009; // Penetration depth of lead
  at 0 kelvin, m
3 T_c = 7.193; // Critical temperature of
  superconducting transition for lead, kelvin
4 T = 5.2; // Temperature at which penetration
  depth for lead becomes lambda_T, kelvin
5 lambda_T = lambda_0*(1-(T/T_c)^4)^(-1/2); //
  Penetration depth of lead at 5.2 kelvin, m
6 printf("\nThe penetration depth of lead at %3.1f K =
  %4.1f nm",T, lambda_T/1e-009);
7
8 // Result
9 // The penetration depth of lead at 5.2 K = 43.4 nm

```

Scilab code Exa 7.6 Isotope Effect in mercury

```

1 // Scilab Code Ex7.6: Isotope Effect in mercury:
  Page-153 (2010)
2 M1 = 199; // Mass of an isotope of mercury, amu
3 T_C1 = 4.185; // Transition temperature of the
  isotope of Hg, K
4 T_C2 = 4.153; // Transition temperature of
  another isotope of Hg, K
5 alpha = 0.5; // Isotope coefficient
6 M2 = M1*(T_C1/T_C2)^(1/alpha); // Mass of another
  isotope of mercury, amu
7 printf("\nThe mass of another isotope of mercury at
  %5.3f K = %6.2f amu",T_C2, M2);
8
9 // Result
10 // The mass of another isotope of mercury at 4.153 K
  = 202.08 amu

```


Chapter 8

SPECIAL THEORY OF RELATIVITY

Scilab code Exa 8.1 Relativistic length contraction

```
1 // Scilab Code Ex8.1: Page-171 (2010)
2 L_0 = 1; // For simplicity, we assume classical
  length to be unity, m
3 c = 1; // For simplicity assume speed of light to
  be unity, m/s
4 L = (1-1/100)*L_0; // Relativistic length, m
5 // Relativistic length contraction gives
6 //  $L = L_0 \cdot \sqrt{1 - v^2/c^2}$ , solving for v
7 v = sqrt(1-(L/L_0)^2)*c; // Speed at which
  relativistic length is 1 percent of the classical
  length, m/s
8 printf("\nThe speed at which relativistic length is
  1 percent of the classical length = %5.3fc", v);
9
10 // Result
11 // The speed at which relativistic length is 1
  percent of the classical length = 0.141c
```

Scilab code Exa 8.2 Time Dilation

```
1 // Scilab Code Ex8.2: Page-171 (2010)
2 c = 1; // For simplicity assume speed of light to
        be unity, m/s
3 v = 0.9*c; // Speed at which beam of particles
        travel, m/s
4 delta_t = 5e-006; // Mean lifetime of particles
        as observed in the Lab. frame, s
5 delta_tau = delta_t*sqrt(1-(v/c)^2); // Proper
        lifetime of particle as per Time Dilation rule, s
6 printf("\nThe proper lifetime of particle = %4.2e s"
        , delta_tau);
7
8 // Result
9 // The proper lifetime of particle = 2.18e-006 s
```

Scilab code Exa 8.4 Relativistic velocity addition

```
1 // Scilab Code Ex8.4: Page-172 (2010)
2 c = 1; // For simplicity assume speed of light to
        be unity, m/s
3 v = 0.6*c; // Speed with which the rocket leaves
        the earth, m/s
4 u_prime = 0.9*c; // Relative speed of second
        rocket w.r.t. the first rocket, m/s
5 u = (u_prime+v)/(1+(u_prime*v)/c^2); // Speed of
        second rocket for same direction of firing as per
        Velocity Addition Rule, m/s
6 printf("\nThe speed of second rocket for same
        direction of firing = %5.3fc", u);
```

```

7 u = (-u_prime+v)/(1-(u_prime*v)/c^2);    // Speed of
    second rocket for opposite direction of firing
    as per Velocity Addition Rule, m/s
8 printf("\nThe speed of second rocket for opposite
    direction of firing = %5.3fc", u);
9
10 // Result
11 // The speed of second rocket for same direction of
    firing = 0.974c
12 // The speed of second rocket for opposite direction
    of firing = -0.652c

```

Scilab code Exa 8.5 Relativistic effects as observed for spaceship

```

1 // Scilab Code Ex8.5: Page-172 (2010)
2 c = 1;    // For simplicity assume speed of light to
    be unity, m/s
3 L0 = 1;    // For simplicity assume length in
    spaceship's frame to be unity, m
4 L = 1/2*L0;    // Length as observed on earth, m
5 // Relativistic length contraction gives
6 // L = L_0*sqrt(1-v^2/c^2), solving for v
7 v = sqrt(1-(L/L0)^2)*c;    // Speed at which length
    of spaceship is observed as half from the earth
    frame, m/s
8 tau = 1;    // Unit time in the spaceship's frame, s
9 t = tau/sqrt(1-(v/c)^2);    // Time dilation of the
    spaceship's unit time, s
10 printf("\nThe speed at which length of spaceship is
    observed as half from the earth frame = %5.3fc",
    v);
11 printf("\nThe time dilation of the spaceship unit
    time = %lg*tau", t);
12
13 // Result

```



```

14 // The speed at which length of spaceship is
    observed as half from the earth frame = 0.866c
15 // The time dilation of the spaceship unit time = 2*
    tau

```

Scilab code Exa 8.6 Time difference and distance between the events

```

1 // Scilab Code Ex8.6: Page-172 (2010)
2 c = 3e+008; // Speed of light in vacuum, m/s
3 v = 0.6*c; // Velocity with which S2 frame moves
    relative to S1 frame, m/s
4 L_factor = 1/sqrt(1-(v/c)^2); // Lorentz factor
5 t1 = 2e-007; // Time for which first event occurs
    , s
6 t2 = 3e-007; // Time for which second event
    occurs, s
7 x1 = 10; // Position at which first event occurs,
    m
8 x2 = 40; // Position at which second event occurs
    , m
9 delta_t = L_factor*(t2 - t1)+L_factor*v/c^2*(x1 - x2
    ); // Time difference between the events, s
10 delta_x = L_factor*(x2 - x1)-L_factor*v*(t2 - t1);
    // Distance between the events, m
11 printf("\nThe time difference between the events =
    %3.1e s", delta_t);
12 printf("\nThe distance between the events = %2d m",
    delta_x);
13
14 // Result
15 // The time difference between the events = 5.0e-008
    s
16 // The distance between the events = 15 m

```

Scilab code Exa 8.7 Speed of unstable particle in the Laboratory frame

```
1 // Scilab Code Ex8.7: Page-173 (2010)
2 c = 3e+008; // Speed of light in vacuum, m/s
3 tau = 2.6e-008; // Mean lifetime the particle in
  its own frame, s
4 d = 20; // Distance which the unstable particle
  travels before decaying, m
5 // As  $t = d/v$  and also  $t = \tau/\sqrt{1-(v/c)^2}$ , so
  that
6 //  $d/v = \tau/\sqrt{1-(v/c)^2}$ , solving for v
7 v = sqrt(d^2/(tau^2+(d/c)^2)); // Speed of the
  unstable particle in Lab. frame, m/s
8 printf("\nThe speed of the unstable particle in Lab.
  frame = %3.1e m/s", v)
9
10 // Result
11 // The speed of the unstable particle in Lab. frame
  = 2.8e+008 m/s
```

Scilab code Exa 8.8 Relativistic effects applied to mu meson

```
1 // Scilab Code Ex8.8: Page-174 (2010)
2 c = 1; // For simplicity assume speed of light to
  be unity, m/s
3 me = 1; // For simplicity assume mass of electron
  to be unity, kg
4 tau = 2.3e-006; // Average lifetime of mu-meson
  in rest frame, s
5 t = 6.9e-006; // Average lifetime of mu-meson in
  laboratory frame, s
```

```

6 // Fromm Time Dilation Rule, tau = t*sqrt(1-(v/c)^2)
  , solving for v
7 v = sqrt(1-(tau/t)^2)*c; // Speed of mu-meson in
  the laboratory frame, m/s
8 c
9 m0 = 207*me; // Rest mass of mu-meson, kg
10 m = m0/sqrt(1-(v/c)^2); // Relativistic variation
  of mass with velocity, kg
11 me = 9.1e-031; // Mass of an electron, kg
12 c = 3e+008; // Speed of light in vacuum, m/s
13 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
14 T = (m*me*c^2 - m0*me*c^2)/e; // Kinetic energy
  of mu-meson, J
15 printf("\nThe speed of mu-meson in the laboratory
  frame = %6.4fc", v);
16 printf("\nThe effective mass of mu-meson = %3d me",
  m);
17 printf("\nThe kinetic energy of mu-meson = %5.1f MeV
  ", T/1e+006);
18
19 // Result
20 // The speed of mu-meson in the laboratory frame =
  0.9428c
21 // The effective mass of mu-meson = 620 me
22 // The kinetic energy of mu-meson = 211.9 MeV

```

Scilab code Exa 8.9 Speed of moving mass

```

1 // Scilab Code Ex8.9: Page-174 (2010)
2 c = 1; // For simplicity assume speed of light to
  be unity, m/s
3 m0 = 1; // For simplicity assume rest mass to be
  unity, kg
4 m = (20/100+1)*m0; // Mass in motion, kg
5 // As m = m0/sqrt(1-(u/c)^2), solving for u

```

```

6 u = sqrt(1-(m0/m)^2)*c;    // Speed of moving mass,
    m/s
7 printf("\nThe speed of moving body, u = %5.3fc", u);
8
9 // Result
10 // The speed of moving body, u = 0.553c

```

Scilab code Exa 8.10 Rate of decreasing mass of sun

```

1 // Scilab Code Ex8.10: Page-175 (2010)
2 c = 3e+008;    // Speed of light in vacuum, m/s
3 dE = 4e+026;    // Energy radiated per second by the
    sun, J/s
4 dm = dE/c^2;    // Rate of decrease of mass of sun,
    kg/s
5 printf("\nThe rate of decrease of mass of sun = %4.2
    e kg/s", dm);
6
7 // Result
8 // The rate of decrease of mass of sun = 4.44e+009
    kg/s

```

Scilab code Exa 8.11 Relativistic mass energy relation

```

1 // Scilab Code Ex8.11: Page-175 (2010)
2 c = 1;    // For simplicity assume speed of light to
    be unity, m/s
3 m0 = 9.1e-031;    // Mass of the electron, kg
4 E0 = 0.512;    // Rest energy of electron, MeV
5 T = 10;    // Kinetic energy of electron, MeV
6 E = T + E0;    // Total energy of electron, MeV
7 // From Relativistic mass-energy relation
8 //  $E^2 = c^2 * p^2 + m0^2 * c^4$ , solving for p

```

```

9 p = sqrt(E^2-m0^2*c^4)/c;    // Momentum of the
    electron , MeV
10 // As E = E0/sqrt(1-(u/c)^2), solving for u
11 u = sqrt(1-(E0/E)^2)*c;    // Velocity of the
    electron , m/s
12 printf("\nThe momentum of the electron = %4.1f/c MeV
    ", p);
13 printf("\nThe velocity of the electron = %6.4fc", u)
    ;
14
15 // Result
16 // The momentum of the electron = 10.5/c MeV
17 // The velocity of the electron = 0.9988c

```

Scilab code Exa 8.13 Mass from relativistic energy

```

1 // Scilab Code Ex8.13: Page-176 (2010)
2 c = 3e+008;    // Speed of light in vacuum, m/s
3 E = 4.5e+017;    // Total energy of object, J
4 px = 3.8e+008;    // X-component of momentum, kg-m/s
5 py = 3e+008;    // Y-component of momentum, kg-m/s
6 pz = 3e+008;    // Z-component of momentum, kg-m/s
7 p = sqrt(px^2+py^2+pz^2);    // Total momentum of
    the object, kg-m/s
8 // From Relativistic mass-energy relation
9 // E^2 = c^2*p^2 + m0^2*c^4, solving for m0
10 m0 = sqrt(E^2/c^4 - p^2/c^2);    // Rest mass of the
    body, kg
11 printf("\nThe rest mass of the body = %4.2f kg", m0)
    ;
12
13 // Result
14 // The rest mass of the body = 4.56 kg

```

Scilab code Exa 8.14 Relativistic momentum of high speed probe

```
1 // Scilab Code Ex8.14: Page-176 (2010)
2 c = 3e+008; // Speed of light in vacuum, m/s
3 m = 50000; // Mass of high speed probe, kg
4 u = 0.8*c; // Speed of the probe, m/s
5 p = m*u/sqrt(1-(u/c)^2); // Momentum of the probe
   , kg-m/s
6 printf("\nThe momentum of the high speed probe = %lg
   kg-m/s", p);
7
8 // Result
9 // The momentum of the high speed probe = 2e+013 kg-
   m/s
```

Scilab code Exa 8.15 Moving electron subjected to the electric field

```
1 // Scilab Code Ex8.15: Page-177 (2010)
2 e = 1.6e-019; // Electronic charge, C = Energy
   equivalent of 1 eV, J/eV
3 m0 = 9.11e-031; // Rest mass of electron, kg
4 c = 3e+008; // Speed of light in vacuum, m/s
5 u1 = 0.98*c; // Initial speed of electron, m/s
6 u2 = 0.99*c; // Final speed of electron, m/s
7 m1 = m0/sqrt(1-(u1/c)^2); // Initial relativistic
   mass of electron, kg
8 m2 = m0/sqrt(1-(u2/c)^2); // Final relativistic
   mass of electron, kg
9 dm = m2 - m1; // Change in relativistic mass of
   the electron, kg
10 W = dm*c^2; // Work done on the electron to
   change its velocity, J
```

```

11 // As  $W = eV$ ,  $V =$  accelerating potential, solving
    for  $V$ 
12  $V = W/e$ ; // Accelerating potential, volt
13 printf("\nThe change in relativistic mass of the
    electron = %4.1e kg", dm);
14 printf("\nThe work done on the electron to change
    its velocity = %4.2f MeV",  $W/(e*1e+006)$ );
15 printf("\nThe accelerating potential = %4.2e volt",
    V);
16
17 // Result
18 // The change in relativistic mass of the electron =
    1.9e-030 kg
19 // The work done on the electron to change its
    velocity = 1.06 MeV
20 // The accelerating potential = 1.06e+006 volt

```

Chapter 9

QUANTUM MECHANICS

Scilab code Exa 9.1 De broglie wavelength of an electron from accelerating potenti

```
1 // Scilab Code Ex9.1: De-broglie wavelength of an
  electron from accelerating potential : Page-202
  (2010)
2 V = 100; // Accelerating potential for electron ,
  volt
3 lambda = sqrt(150/V)*1e-010; // de-Broglie
  wavelength of electron , m
4 printf("\nThe De-Broglie wavelength of electron = %4
  .2e m", lambda);
5
6 // Result
7 // The De-Broglie wavelength of electron = 1.22e-010
  m
```

Scilab code Exa 9.2 De broglie wavelength of an electron from kinetic energy

```
1 // Scilab Code Ex9.2: De-broglie wavelength of an
  electron from kinetic energy : Page-203 (2010)
```



```

2 e = 1.6e-019;    // Energy equivalent of 1 eV, J/eV
3 h = 6.626e-034; // Planck's constant, Js
4 m = 9.1e-031;   // Mass of the electron, kg
5 Ek = 10;        // Kinetic energy of electron, eV
6 // Ek = p^2/(2*m), solving for p
7 p = sqrt(2*m*Ek*e); // Momentum of the electron,
    kg-m/s
8 lambda = h/p ; // de-Broglie wavelength of
    electron from De-Broglie relation, m
9 printf("\nThe de-Broglie wavelength of electron = %4
    .2e nm", lambda/1e-009);
10
11 // Result
12 // The de-Broglie wavelength of electron = 3.88e-001
    nm

```

Scilab code Exa 9.4 Uncertainty principle for position and momentum

```

1 // Scilab Code Ex9.4: Uncertainty principle for
    position and momentum: Page-203 (2010)
2 h = 6.626e-034; // Planck's constant, Js
3 m = 9.1e-031;   // Mass of the electron, kg
4 v = 1.1e+006;   // Speed of the electron, m/s
5 p = m*v;        // Momentum of the electron, kg-m/s
6 dp = 0.1/100*p; // Uncertainty in momentum, kg-m/
    s
7 h_bar = h/(2*%pi); // Reduced Planck's constant,
    Js
8 // From Heisenberg uncertainty principle,
9 // dx*dp = h_bar/2, solving for dx
10 dx = h_bar/(2*dp); // Uncertainty in position, m
11 printf("\nThe uncertainty in position of electron =
    %4.2e m", dx);
12
13 // Result

```

```
14 // The uncertainty in position of electron = 5.27e
    -008 m
```

Scilab code Exa 9.5 Uncertainty principle for energy and time

```
1 // Scilab Code Ex9.5: Uncertainty principle for
    energy and time: Page-203 (2010)
2 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
3 h = 6.626e-034; // Planck's constant, Js
4 dt = 1e-008; // Uncertainty in time, s
5 h_bar = h/(2*%pi); // Reduced Planck's constant,
    Js
6 // From Heisenberg uncertainty principle,
7 //  $dE*dt = \hbar/2$ , solving for dE
8 dE = h_bar/(2*dt*e); // Uncertainty in energy of
    the excited state, m
9 printf("\nThe uncertainty in energy of the excited
    state = %4.2e eV", dE);
10
11 // Result
12 // The uncertainty in energy of the excited state =
    3.30e-008 eV
```

Scilab code Exa 9.6 Width of spectral line from Uncertainty principle

```
1 // Scilab Code Ex9.6: Width of spectral line from
    Uncertainty principle: Page-204 (2010)
2 c = 3e+008; // Speed of light, m/s
3 dt = 1e-008; // Average lifetime, s
4 lambda = 400e-009; // Wavelength of spectral line
    , m
5 // From Heisenberg uncertainty principle,
```

```

6 // dE = h_bar/(2*dt) and also dE = h*c/lambda^2*
  d_lambda, which give
7 // h_bar/(2*dt) = h*c/lambda^2*d_lambda, solving for
  d_lambda
8 d_lambda = lambda^2/(4*pi*c*dt); // Width of
  spectral line, m
9 printf("\nThe width of spectral line = %4.2e m",
  d_lambda);
10
11 // Result
12 // The width of spectral line = 4.24e-015 m

```

Scilab code Exa 9.14 Probability of electron moving in 1D box

```

1 // Scilab Code Ex9.14: Probability of electron
  moving in 1D box : Page-207 (2010)
2 a = 2e-010; // Width of 1D box, m
3 x1 = 0; // Position of first extreme of the box,
  m
4 x2 = 1e-010; // Position of second extreme of the
  box, m
5 P = integrate('2/a*(sin(2*pi*x/a))^2', 'x', x1, x2)
  ; // The probability of finding the electron
  between x = 0 and x = 1e-010
6 printf("\nThe probability of finding the electron
  between x = 0 and x = 1e-010 = %3.1f", P);
7
8 // Result
9 // The probability of finding the electron between x
  = 0 and x = 1e-010 = 0.5

```

Chapter 10

STATISTICAL MECHANICS

Scilab code Exa 10.1 Ratio of occupancy of two states

```
1 // Scilab Code Ex10.1: Page-222 (2010)
2 k = 1.38e-023; // Boltzmann constant, J/K
3 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
4 g1 = 2; // The degeneracy of ground state
5 g2 = 8; // The degeneracy of excited state
6 delta_E = 10.2; // Energy of excited state above
   the ground state, eV
7 T = 6000; // Temperature of the state, K
8 D_ratio = g2/g1; // Ratio of degeneracy of states
9 N_ratio = D_ratio*exp(-delta_E/(k*T/e)); // Ratio
   of occupancy of the excited to the ground state
10 printf("\nThe ratio of occupancy of the excited to
   the ground state at %d K = %4.2e", T, N_ratio);
11
12 // Result
13 // The ratio of occupancy of the excited to the
   ground state at 6000 K = 1.10e-008
```

Scilab code Exa 10.4 Number density and fermi energy of silver

```

1 // Scilab Code Ex10.4: Page-223 (2010)
2 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
3 N_A = 6.023e+023; // Avogadro's number
4 h = 6.626e-034; // Planck's constant, Js
5 me = 9.1e-031; // Mass of electron, kg
6 rho = 10.5; // Density of silver, g per cm
7 m = 108; // Molecular mass of silver, g/mol
8 N_D = rho*N_A/(m*1e-006); // Number density of
   conduction electrons, per metre cube
9 E_F = h^2/(8*me)*(3/%pi*N_D)^(2/3);
10 printf("\nThe number density of conduction electrons
   = %4.2e per metre cube", N_D);
11 printf("\nThe Fermi energy of silver = %4.2f eV",
   E_F/e);
12
13 // Result
14 // The number density of conduction electrons = 5.86
   e+028 per metre cube
15 // The Fermi energy of silver = 5.51 eV

```

Scilab code Exa 10.5 Electronic contribution to the molar heat capacity of silver

```

1 // Scilab Code Ex10.5: Page-224 (2010)
2 N_A = 6.023e+023; // Avogadro's number
3 k = 1.38e-023; // Boltzmann constant, J/K
4 T = 293; // Temperature of sodium, K
5 E_F = 3.24; // Fermi energy of sodium, eV
6 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
7 C_v = %pi^2*N_A*k^2*T/(2*E_F*e); // Molar
   specific heat of sodium, J/mole/K
8 printf("\nThe molar specific heat of sodium = %4.2f
   J/mole/K", C_v);
9
10 // Result
11 // The molar specific heat of sodium = 0.32 J/mole/K

```

Scilab code Exa 10.6 Fermi energy and mean energy of aluminium

```
1 // Scilab Code Ex10.6: Page-224 (2010)
2 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
3 h = 6.626e-034; // Planck's constant, Js
4 m = 9.1e-031; // Mass of the electron, kg
5 N_D = 18.1e+028; // Number density of conduction
    electrons in Al, per metre cube
6 E_F = h^2/(8*m)*(3/%pi*N_D)^(2/3); // Fermi
    energy of aluminium, J
7 Em_0 = 3/5*E_F; // Mean energy of the electron
    at 0K, J
8 printf("\nThe Fermi energy of aluminium = %5.2f eV",
    E_F/e);
9 printf("\nThe mean energy of the electron at 0K =
    %4.2f eV", Em_0/e);
10
11 // Result
12 // The Fermi energy of aluminium = 11.70 eV
13 // The mean energy of the electron at 0K = 7.02 eV
```

Chapter 11

LASERS

Scilab code Exa 11.1 Ratio of spontaneous and stimulated emission

```
1 // Scilab Code Ex11.1: Page-249 (2010)
2 h = 6.626e-034; // Planck's constant, Js
3 c = 3e+08; // Speed of light in free space, m/s
4 k = 1.38e-023; // Boltzmann constant, J/K
5 T = 300; // Temperature at absolute scale, K
6 lambda = 5500e-010; // Wavelength of visible
   light, m
7 rate_ratio = exp(h*c/(lambda*k*T))-1; // Ratio of
   spontaneous emission to stimulated emission
8 printf("\nThe ratio of spontaneous emission to
   stimulated emission for visible region = %1.0e",
   rate_ratio);
9 lambda = 1e-02; // Wavelength of microwave, m
10 rate_ratio = exp(h*c/(lambda*k*T))-1; // Ratio of
   spontaneous emission to stimulated emission
11 printf("\nThe ratio of spontaneous emission to
   stimulated emission for microwave region = %6.4f"
   , rate_ratio);
12
13 // Result
14 // The ratio of spontaneous emission to stimulated
```

```

    emission for visible region = 8e+037
15 // The ratio of spontaneous emission to stimulated
    emission for microwave region = 0.0048

```

Scilab code Exa 11.2 Energy of excited state of laser system

```

1 // Scilab Code Ex11.2: Page-250 (2010)
2 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
3 h = 6.626e-034; // Planck's constant, Js
4 c = 3e+08; // Speed of light in free space, m/s
5 lambda = 690e-009; // Wavelength of laser light,
    m
6 E_lower = 30.5; // Energy of lower state, eV
7 E = h*c/(lambda*e); // Energy of the laser light,
    eV
8 E_ex = E_lower + E; // Energy of excited state of
    laser system, eV
9 printf("\nThe energy of excited state of laser
    system = %4.1f eV", E_ex);
10
11 // Result
12 // The energy of excited state of laser system =
    32.3 eV

```

Scilab code Exa 11.3 Condition of equivalence of stimulated and spontaneous emission

```

1 // Scilab Code Ex11.3: Page-250 (2010)
2 h = 6.626e-034; // Planck's constant, Js
3 k = 1.38e-023; // Boltzmann constant, J/K
4 // Stimulated Emission = Spontaneous Emission <=>
    exp(h*f/(k*T))-1 = 1 i.e.
5 // f/T = log(2)*k/h = A

```



```

6 A = log(2)*k/h; // Frequency per unit
   temperature , Hz/K
7 printf("\nThe stimulated emission equals spontaneous
   emission iff f/T = %4.2e Hz/K", A);
8
9 // Result
10 // The stimulated emission equals spontaneous
   emission iff f/T = 1.44e+010 Hz/K

```

Scilab code Exa 11.4 Area and intensity of image formed by laser

```

1 // Scilab Code Ex11.4: Page-250 (2010)
2 lambda = 500e-009; // Wavelength of laser light ,
   m
3 f = 15e-02; // Focal length of the lens , m
4 d = 2e-02; // Diameter of the aperture of source ,
   m
5 a = d/2; // Radius of the aperture of source , m
6 P = 5e-003; // Power of the laser , W
7 A = %pi*lambda^2*f^2/a^2; // Area of the spot at
   the focal plane , metre square
8 I = P/A; // Intensity at the focus , W per metre
   square
9 printf("\nThe area of the spot at the focal plane =
   %4.2e metre square", A);
10 printf("\nThe intensity at the focus = %4.2e watt
   per metre square", I);
11
12 // Result
13 // The area of the spot at the focal plane = 1.77e
   -010 metre square
14 // The intensity at the focus = 2.83e+007 watt per
   metre square

```

Scilab code Exa 11.5 Rate of energy released in a pulsed laser

```
1 // Scilab Code Ex11.5: Page-251 (2010)
2 h = 6.626e-034; // Planck's constant, Js
3 c = 3e+08; // Speed of light in free space, m/s
4 lambda = 1064e-009; // Wavelength of laser light,
   m
5 P = 0.8; // Average power output per laser pulse,
   W
6 dt = 25e-003; // Pulse width of laser, s
7 E = P*dt; // Energy released per pulse, J
8 N = E/(h*c/lambda); // Number of photons in a
   pulse
9 printf("\nThe energy released per pulse = %2.0e J",
   E);
10 printf("\nThe number of photons in a pulse = %4.2e",
   N);
11
12 // Result
13 // The energy released per pulse = 2e-002 J
14 // The number of photons in a pulse = 1.07e+017
```

Scilab code Exa 11.6 Angular and linear spread of laser beam

```
1 // Scilab Code Ex11.6:Page-251 (2010)
2 lambda = 693e-009; // Wavelength of laser beam, m
3 D = 3e-003; // Diameter of laser beam, m
4 d_theta = 1.22*lambda/D; // Angular spread of
   laser beam, rad
5 d = 300e+003; // Height of a satellite above the
   surface of earth, m
```

```
6 a = d_theta*d;    // Diameter of the beam on the
   satellite , m
7 printf("\nThe height of a satellite above the
   surface of earth = %4.2e rad", d_theta);
8 printf("\nThe diameter of the beam on the satellite
   = %4.1f m", a);
9
10 // Result
11 // The height of a satellite above the surface of
   earth = 2.82e-004 rad
12 // The diameter of the beam on the satellite = 84.5
   m
```

Chapter 12

HOLOGRAPHY AND FIBRE OPTICS

Scilab code Exa 12.1 Parameters of step index fibre

```
1 // Scilab Code Ex12.1: Parameters of step index
   fibre : Page-271 (2010)
2 n1 = 1.43; // Refractive index of fibre core
3 n2 = 1.4; // Refractive index of fibre cladding
4 // As  $\sin(\alpha_c) = n_2/n_1$ , solving for  $\alpha_c$ 
5 alpha_c = asind(n2/n1); // Critical angle for
   optical fibre, degrees
6 // AS  $\cos(\theta_c) = n_2/n_1$ , solving for  $\theta_c$ 
7 theta_c = acosd(n2/n1); // Critical propagation
   angle for optical fibre, degrees
8 NA = sqrt(n1^2 - n2^2); // Numerical aperture for
   optical fibre
9 printf("\nThe critical angle for optical fibre = %5
   .2f degrees", alpha_c);
10 printf("\nThe critical propagation angle for optical
   fibre = %5.2f degrees", theta_c);
11 printf("\Numerical aperture for optical fibre = %4
   .2f", NA);
12
```

```

13 // Result
14 // The critical angle for optical fibre = 78.24
    degrees
15 // The critical propagation angle for optical fibre
    = 11.76 degrees
16 // Numerical aperture for optical fibre = 0.29

```

Scilab code Exa 12.2 Parameters of optical fibre

```

1 // Scilab Code Ex12.2: Parameters of optical fibre :
    Page-271 (2010)
2 n1 = 1.45; // Refractive index of fibre core
3 n2 = 1.4; // Refractive index of fibre cladding
4 NA = sqrt(n1^2 - n2^2); // Numerical aperture for
    optical fibre
5 // As  $\sin(\theta_a) = \sqrt{n1^2 - n2^2}$ , solving for
    theta_a
6 theta_a = asind(sqrt(n1^2 - n2^2)); // Half of
    acceptance angle of optical fibre, degrees
7 theta_accp = 2*theta_a; // Acceptance angle of
    optical fibre
8 Delta = (n1 - n2)/n1; // Relative refractive
    index difference
9 printf("\nNumerical aperture for optical fibre = %5
    .3f", NA);
10 printf("\nThe acceptance angle of optical fibre = %4
    .1f degrees", theta_accp);
11 printf("\nRelative refractive index difference = %5
    .3f", Delta);
12
13 // Result
14 // Numerical aperture for optical fibre = 0.377
15 // The acceptance angle of optical fibre = 44.4
    degrees
16 // Relative refractive index difference = 0.034

```

Scilab code Exa 12.3 Numerical aperture and acceptance angle of step index fibre

```
1 // Scilab Code Ex12.3: Numerical aperture and
  acceptance angle of step index fibre : Page-271
  (2010)
2 n1 = 1.55; // Refractive index of fibre core
3 n2 = 1.53; // Refractive index of fibre cladding
4 n0 = 1.3; // Refractive index of medium
5 NA = sqrt(n1^2 - n2^2); // Numerical aperture for
  optical fibre
6 // n0*sin(theta_a) = sqrt(n1^2 - n2^2) = NA, solving
  for theta_a
7 theta_a = asind(sqrt(n1^2 - n2^2)/n0); // Half of
  acceptance angle of optical fibre, degrees
8 theta_accp = 2*theta_a; // Acceptance angle of
  optical fibre
9 printf("\nNumerical aperture for step index fibre =
  %5.3f", NA);
10 printf("\nThe acceptance angle of step index fibre =
  %2d degrees", theta_accp);
11
12 // Result
13 // Numerical aperture for step index fibre = 0.248
14 // The acceptance angle of step index fibre = 22
  degrees
```

Scilab code Exa 12.5 Output power in fibre optic communication

```
1 // Scilab Code Ex12.5: Output power in fibre optic
  communication : Page-272 (2010)
2 alpha = 2; // Power loss through optical fibre ,
  dB/km
```

```
3 P_in = 500;    // Power input of optical fibre , micro
    -watt
4 z = 10;      // Length of the optical fibre , km
5 // As  $\alpha = 10/z * \log_{10}(P_{in}/P_{out})$ , solving for
    P_out
6 P_out = P_in/10^(alpha*z/10);    // Output power in
    fibre optic communication , W
7 printf("\nThe output power in fibre optic
    communication = %ld micro-watt", P_out);
8
9 // Result
10 // The output power in fibre optic communication = 5
    micro-watt
```

Chapter 13

DIELECTRIC PROPERTIES OF MATERIALS

Scilab code Exa 13.1 Electronic Polarizability of atom

```
1 // Scilab Code Ex13.1: Electronic Polarizability of
  atom : Page-287 (2010)
2 epsilon_0 = 8.854e-012; // Absolute electrical
  permittivity of free space, farad per metre
3 R = 0.52e-010; // Radius of hydrogen atom,
  angstrom
4 n = 9.7e+026; // Number density of hydrogen, per
  metre cube
5 alpha_e = 4*pi*epsilon_0*R^3; // Electronic
  polarizability of hydrogen atom, farad-metre
  square
6 printf("\nThe electronic polarizability of hydrogen
  atom = %4.2e farad-metre square", alpha_e);
7
8 // Result
9 // The electronic polarizability of hydrogen atom =
  1.56e-041 farad-metre square
```

Scilab code Exa 13.2 Parallel plate capacitor

```
1 // Scilab Code Ex13.2: Parallel plate capacitor:
   Page-287 (2010)
2 epsilon_0 = 8.854e-012; // Absolute electrical
   permittivity of free space, farad per metre
3 A = 100e-004; // Area of a plate of parallel
   plate capacitor, metre square
4 d = 1e-002; // Distance between the plates of the
   capacitor, m
5 V = 100; // Potential applied to the plates of
   the capacitor, volt
6 C = epsilon_0*A/d; // Capacitance of parallel
   plate capacitor, farad
7 Q = C/V; // Charge on the plates of the capacitor
   , coulomb
8 printf("\nThe capacitance of parallel plate
   capacitor = %5.3e F", C);
9 printf("\nThe charge on the plates of the capacitor
   = %5.3e C", Q);
10
11 // Result
12 // The capacitance of parallel plate capacitor =
   8.854e-012 F
13 // The charge on the plates of the capacitor = 8.854
   e-014 C
```

Scilab code Exa 13.3 Dielectric displacement of medium

```
1 // Scilab Code Ex13.3: Dielectric displacement of
   medium: Page-288 (2010)
```

```

2  epsilon_0 = 8.854e-012;    // Absolute electrical
    permittivity of free space, farad per metre
3  epsilon_r = 5.0;    // Dielectric constant of the
    material between the plates of capacitor
4  V = 15;    // Potential difference applied between
    the plates of the capacitor, volt
5  d = 1.5e-003;    // Separation between the plates of
    the capacitor, m
6  // Electric displacement,  $D = \epsilon_0 \epsilon_r E$ ,
    as  $E = V/d$ , so
7  D = epsilon_0*epsilon_r*V/d;    // Dielectric
    displacement, coulomb per metre square
8  printf("\nThe dielectric displacement = %5.3e
    coulomb per metre square", D);
9
10 // Result
11 // The dielectric displacement = 4.427e-007 coulomb
    per metre square

```

Scilab code Exa 13.4 Relative dielectric constant

```

1  // Scilab Code Ex13.4: Relative dielectric constant
    : Page-288 (2010)
2  epsilon_0 = 8.854e-012;    // Absolute electrical
    permittivity of free space, farad per metre
3  N = 3.0e+028;    // Number density of solid
    elemental dielectric, atoms per metre cube
4  alpha_e = 1e-040;    // Electronic polarizability,
    farad metre square
5  epsilon_r = 1 + N*alpha_e/epsilon_0;    // Relative
    dielectric constant of the material
6  printf("\nThe Relative dielectric constant of the
    material = %5.3f", epsilon_r);
7
8  // Result

```

```
9 // The Relative dielectric constant of the material
   = 1.339
```

Scilab code Exa 13.5 Atomic polarizability of sulphur

```
1 // Scilab Code Ex13.5: Atomic polarizability of
   sulphur : Page-288 (2010)
2 N_A = 6.023e+023; // Avogadro's number, per mole
3 epsilon_0 = 8.854e-012; // Absolute electrical
   permittivity of free space, farad per metre
4 epsilon_r = 3.75; // Relative dielectric constant
5 d = 2050; // Density of sulphur, kg per metre
   cube
6 y = 1/3; // Internal field constant
7 M = 32; // Atomic weight of sulphur, g/mol
8 N = N_A*1e+03*d/M; // Number density of atoms of
   sulphur, per metre cube
9 // Lorentz relation for local fields give
10 //  $E_{\text{local}} = E + P/(3*\epsilon_0)$  which gives
11 //  $(\epsilon_r - 1)/(\epsilon_r + 2) = N*\alpha_e/(3*$ 
    $\epsilon_0)$ , solving for  $\alpha_e$ 
12  $\alpha_e = (\epsilon_r - 1)/(\epsilon_r + 2)*3*$ 
    $\epsilon_0/N$ ; // Electronic polarizability of
   sulphur, farad metre square
13 printf("\\nThe electronic polarizability of sulphur =
   %5.3e farad metre square",  $\alpha_e$ );
14
15 // Result
16 // The electronic polarizability of sulphur = 3.292e
   -040 farad metre square
```

Scilab code Exa 13.6 Electronic polarizability from refractive index

```

1 // Scilab Code Ex13.6: Electronic polarizability
  from refractive index : Page-289 (2010)
2 N = 3e+028; // Number density of atoms of
  dielectric material, per metre cube
3 epsilon_0 = 8.854e-012; // Absolute electrical
  permittivity of free space, farad per metre
4 n = 1.6; // Refractive index of dielectric
  material
5 // As  $(n^2 - 1)/(n^2 + 2) = N \cdot \alpha_e / (3 \cdot \epsilon_0)$ ,
  solving for  $\alpha_e$ 
6 alpha_e = (n^2 - 1)/(n^2 + 2)*3*epsilon_0/N; //
  Electronic polarizability of dielectric material,
  farad metre square
7 printf("\nThe electronic polarizability of
  dielectric material = %4.2e farad metre square",
  alpha_e);
8
9 // Result
10 // The electronic polarizability of dielectric
  material = 3.03e-040 farad metre square

```

Scilab code Exa 13.7 Ratio of electronic polarizability to ionic polarizability

```

1 // Scilab Code Ex13.7: Ratio of electronic
  polarizability to ionic polarizability: Page-289
  (2010)
2 epsilon_r = 4.9; // Absolute relative dielectric
  constant of material, farad per metre
3 n = 1.6; // Refractive index of dielectric
  material
4 // As  $(n^2 - 1)/(n^2 + 2) \cdot (\alpha_e + \alpha_i) / \alpha_e = N \cdot (\alpha_e + \alpha_i) / (3 \cdot \epsilon_0) = (\epsilon_r - 1) / (\epsilon_r + 2)$ , solving for
   $\alpha_i / \alpha_e$ 
5 alpha_ratio = ((epsilon_r - 1)/(epsilon_r + 2))*(n^2

```

```
    + 2)/(n^2 - 1) - 1)^(-1);    // Ratio of
    electronic polarizability to ionic polarizability
6 printf("\nThe ratio of electronic polarizability to
    ionic polarizability = %4.2f", alpha_ratio);
7
8 // Result
9 // The ratio of electronic polarizability to ionic
    polarizability = 1.53
```

Chapter 14

MAGNETIC PROPERTIES OF MATERIALS

Scilab code Exa 14.1 Spontaneous magnetisation of the substance

```
1 // Scilab Code Ex14.1: Spontaneous magnetisation of
  the substance: Page-306 (2010)
2 N = 6.023e+023; // Avogadro's number. per mole
3 A = 56; // Atomic weight of the substance, g/mole
4 d = 7.9; // Density of the substance, gram per cm
  cube
5 m_B = 9.27e-024; // Bohr's Magnetron, joule per
  tesla
6 m = 2.2*m_B; // Magnetic moment of substance,
  joule per tesla
7 n = d*N/A*1e+006; // Number of atoms per unit
  volume of the substance, per metre cube
8 M = n*m; // Spontaneous magnetisation of the
  substance, ampere per metre
9 printf("\nThe spontaneous magnetisation of the
  substance = %4.2e ampere per metre", M);
10
11 // Result
12 // The spontaneous magnetisation of the substance =
```

1.73e+006 ampere per metre

Scilab code Exa 14.2 Relative permeability of ferromagnetic material

```
1 // Scilab Code Ex14.2: Relative permeability of
  ferromagnetic material : Page-307 (2010)
2 H = 200; // Field strength to which the
  ferromagnetic material is subjected, ampere per
  metre
3 M = 3100; // Magnetisation of the ferromagnetic
  material, ampere per metre
4 chi = M/H; // Magnetic susceptibility
5 mu_r = 1 + chi; // Relative permeability of
  ferromagnetic material
6 printf("\nThe relative permeability of ferromagnetic
  material = %4.1f", mu_r);
7
8 // Result
9 // The relative permeability of ferromagnetic
  material = 16.5
```

Scilab code Exa 14.3 Relative permeability from magnetisation

```
1 // Scilab Code Ex14.3: Relative permeability from
  magnetisation : Page-307 (2010)
2 H = 300; // Field strength to which the
  ferromagnetic material is subjected, ampere per
  metre
3 M = 4400; // Magnetisation of the ferromagnetic
  material, ampere per metre
4 chi = M/H; // Magnetic susceptibility
5 mu_r = 1 + chi; // Relative permeability of
  ferromagnetic material
```

```

6 printf("\nThe relative permeability of ferromagnetic
    material = %5.2f", mu_r);
7
8 // Result
9 // The relative permeability of ferromagnetic
    material = 15.67

```

Scilab code Exa 14.4 Magnetic flux density and magnetisation of diamagnetic materi

```

1 // Scilab Code Ex14.4: Magnetic flux density and
    magnetisation of diamagnetic material : Page-307
    (2010)
2 mu_0 = 4*pi*1e-07; // Magnetic permeability of
    free space, tesla metre per ampere
3 H = 10000; // Field strength to which the
    diamagnetic material is subjected, ampere per
    metre
4 chi = -0.4e-005; // Magnetic susceptibility
5 M = chi*H; // Magnetisation of the diamagnetic
    material, ampere per metre
6 B = mu_0*(H + M); // Magnetic flux density of
    diamagnetic material, T
7 printf("\nThe magnetisation of diamagnetic material
    = %4.2f ampere per metre", M);
8 printf("\nThe magnetic flux density of diamagnetic
    material = %6.4f T", B);
9
10 // Result
11 // The magnetisation of diamagnetic material = -0.04
    ampere per metre
12 // The Magnetic flux density of diamagnetic material
    = 0.0126 T

```

Scilab code Exa 14.5 Magnetisation Magnetic flux density relative permeability of

```
1 // Scilab Code Ex14.5: Magnetisation–Magnetic flux
  density–relative permeability of diamagnetic
  material : Page–307 (2010)
2 mu_0 = 4*%pi*1e-07; // Magnetic permeability of
  free space, tesla metre per ampere
3 H = 1.2e+005; // Field strength to which the
  diamagnetic material is subjected, ampere per
  metre
4 chi = -4.2e-006; // Magnetic susceptibility
5 M = chi*H; // Magnetisation of the diamagnetic
  material, ampere per metre
6 B = mu_0*(H + M); // Magnetic flux density of
  diamagnetic material, T
7 mu_r = M/H + 1; // The relative permeability of
  diamagnetic material
8 printf("\nThe magnetisation of diamagnetic material
  = %5.3f ampere per metre", M);
9 printf("\nThe magnetic flux density of diamagnetic
  material = %5.3f T", B);
10 printf("\nThe relative permeability of diamagnetic
  material = %f T", mu_r);
11 // Result
12 // The magnetisation of diamagnetic material =
  -0.504 ampere per metre
13 // The magnetic flux density of diamagnetic material
  = 0.151 T
14 // The relative permeability of diamagnetic material
  = 0.999996 T
```

Scilab code Exa 14.6 Mean radius of body centered cubic structure

```
1 // Scilab Code Ex14.6: Mean radius of body centered
  cubic structure: Page–308 (2010)
```

```

2 chi = 5.6e-006;    // Magnetic susceptibility of
    diamagnetic material
3 m = 9.1e-31;      // Mass of an electron , kg
4 mu_0 = 4*pi*1e-07; // Magnetic permeability of
    free space, tesla metre per ampere
5 Z = 1;           /// Atomic number
6 e = 1.6e-019;    // Electronic charge , C
7 a = 2.53e-010;    // Lattice parameter of bcc
    structure , m
8 N = 2/a^3;       // The number of electrons per unit
    volume, per metre cube
9 r = sqrt(chi*6*m/(mu_0*Z*e^2*N)); // Mean radius
    of body centered cubic structure as per Langevin
    relation for Diamagnetic susceptibility , m
10 printf("\nThe mean radius of body centered cubic
    structure = %5.3e angstrom", r/1e-010);
11
12 // Result
13 // The mean radius of body centered cubic structure
    = 8.773e-001 angstrom

```

Scilab code Exa 14.7 Susceptibility and magnetisation of paramagnetic salt

```

1 // Scilab Code Ex14.7: Susceptibility and
    magnetisation of paramagnetic salt: Page-308
    (2010)
2 mu_0 = 4*pi*1e-07; // Magnetic permeability of
    free space, tesla metre per ampere
3 N_A = 6.02e+026;   // Avogadro's number, per kmol
4 rho = 4370;        // Density of paramagnetic salt , kg
    per metre cube
5 M = 168.5;         // Molecular weight of paramagnetic
    salt , g/mol
6 T = 27+273;       // Temperature of paramagnetic salt ,
    K

```

```

7 H = 2e+005;    // Field strength to which the
    paramagnetic salt is subjected, ampere per metre
8 mu_B = 9.27e-024;    // Bohr's magneton, ampere
    metre square
9 p = 2;    // Number of Bohr magnetons per molecule
10 k = 1.38e-023;    // Boltzmann constant, J/K
11 N = rho*N_A/M;    // Total density of atoms in the
    paramagnetic salt, per metr cube
12 chi = mu_0*N*p^2*mu_B^2/(3*k*T);    // Magnetic
    susceptibility of paramagnetic salt
13 M = chi*H;    // Magnetisation of paramagnetic salt,
    ampere per metre
14 printf("\nThe magnetic susceptibility of
    paramagnetic salt = %4.2e per metre", chi);
15 printf("\nThe magnetisation of paramagnetic salt =
    %4.2e ampere per metre", M);
16
17 // Result
18 // The magnetic susceptibility of paramagnetic salt
    = 5.43e-004 per metre
19 // The magnetisation of paramagnetic salt = 1.09e
    +002 ampere per metre

```

Chapter 15

THERMAL PROPERTIES

Scilab code Exa 15.1 Debye temperature of aluminium

```
1 // Scilab Code Ex15.1: Page-323 (2010)
2 k = 1.38e-023; // Boltzmann constant, J/K
3 h = 6.626e-034; // Planck's constant, Js
4 f_D = 64e+011; // Debye frequency for Al, Hz
5 theta_D = h*f_D/k; // Debye temperature, K
6 printf("\nThe Debye temperature of aluminium = %5.1f
   K", theta_D);
7
8 // Result
9 // The Debye temperature of aluminium = 307.3 K
```

Scilab code Exa 15.2 Lattice specific heat of carbon

```
1 // Scilab Code Ex15.2: Page-323 (2010)
2 N = 6.02e+026; // Avogadro's number, per kmol
3 k = 1.38e-023; // Boltzmann constant, J/K
4 h = 6.626e-034; // Planck's constant, Js
5 f_D = 40.5e+012; // Debye frequency for Al, Hz
```

```

6 T = 30; // Temperature of carbon , Ks
7 theta_D = h*f_D/k; // Debye temperature , K
8 C_1 = 12/5*pi^4*N*k*(T/theta_D)^3; // Lattice
    specific heat of carbon , J/k-mol/K
9 printf("\nThe lattice specific heat of carbon = %4.2
    f J/k-mol/K", C_1);
10
11 // Result
12 // The lattice specific heat of carbon = 7.13 J/k-
    mol/K

```

Scilab code Exa 15.3 Einstein frequency for Cu

```

1 // Scilab Code Ex15.3: Page-323 (2010)
2 k = 1.38e-023; // Boltzmann constant , J/K
3 h = 6.626e-034; // Planck's constant , Js
4 theta_E = 1990; // Einstein temperature of Cu, K
5 f_E = k*theta_E/h; // Einstein frequency for Cu,
    K
6 printf("\nThe Einstein frequency for Cu = %4.2e Hz",
    f_E);
7 printf("\nThe frequency falls in the near infrared
    region");
8
9 // Result
10 // The Einstein frequency for Cu = 4.14e+013 Hz
11 // The frequency falls in the near infrared region

```

Scilab code Exa 15.4 Electronic and lattice heat capacities for Cu

```

1 // Scilab Code Ex15.4: Page-323 (2010)
2 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
3 N = 6.02e+023; // Avogadro's number , per mol

```

```

4 T = 0.05;      // Temperature of Cu, K
5 E_F = 7;      // Fermi energy of Cu, eV
6 k = 1.38e-023; // Boltzmann constant, J/K
7 h = 6.626e-034; // Planck's constant, Js
8 theta_D = 348; // Debye temperature of Cu, K
9 C_e = %pi^2*N*k^2*T/(2*E_F*e); // Electronic heat
    capacity of Cu, J/mol/K
10 C_V = 12/5*%pi^4*N*k*(T/theta_D)^3; // Lattice heat
    capacity of Cu, J/mol/K
11 printf("\nThe electronic heat capacity of Cu = %4.2e
    J/mol/K", C_e);
12 printf("\nThe lattice heat capacity of Cu = %4.2e J/
    mol/K", C_V);
13
14 // Result
15 // The electronic heat capacity of Cu = 2.53e-005 J/
    mol/K
16 // The lattice heat capacity of Cu = 5.76e-009 J/mol
    /K

```

Scilab code Exa 15.5 Einstein lattice specific heat

```

1 // Scilab Code Ex15.5: Page-324 (2010)
2 T = 1; // For simplicity assume temperature to be
    unity, K
3 R = 1; // For simplicity assume molar gas
    constant to be unity, J/mol/K
4 theta_E = T; // Einstein temperature, K
5 C_V = 3*R*(theta_E/T)^2*exp(theta_E/T)/(exp(theta_E/
    T)-1)^2; // Einstein lattice specific heat, J/
    mol/K
6 printf("\nThe Einstein lattice specific heat, C_v =
    %4.2f X 3R", C_V/3);
7
8 // Result

```

```
9 // The Einstein lattice specific heat,  $C_v = 0.92 \times$   
3R
```

Scilab code Exa 15.6 Molar electronic heat capacity of zinc

```
1 // Scilab Code Ex15.6: Page-324 (2010)  
2 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV  
3 v = 2; // Valency of Zn atom  
4 N = v*6.02e+023; // Avogadro's number, per mol  
5 T = 300; // Temperature of Zn, K  
6 E_F = 9.38; // Fermi energy of Zn, eV  
7 k = 1.38e-023; // Boltzmann constant, J/K  
8 h = 6.626e-034; // Planck's constant, Js  
9 C_e = %pi^2*N*k^2*T/(2*E_F*e); // Electronic heat  
capacity of Zn, J/mol/K  
10 printf("\nThe molar electronic heat capacity of zinc  
= %5.3f J/mol/K", C_e);  
11  
12 // Result  
13 // The molar electronic heat capacity of zinc =  
0.226 J/mol/K
```

Chapter 17

ULTRASONICS

Scilab code Exa 17.1 Thickness of vibrating quartz at resonance

```
1 // Scilab Code Ex17.1: Thickness of vibrating quartz
   at resonance : Page-352 (2010)
2 f = 3e+006; // Fundamental vibrational frequency
   of quartz crystal, MHz
3 Y = 7.9e+010; // Young's modulus of quartz,
   newton per metre
4 rho = 2650; // Density of quartz, kg per metre
   cube
5 // We have for resonant frequency
6 //  $f = 1/(2*l)*\sqrt{Y/\rho}$ , solving for l
7 l = 1/(2*f)*sqrt(Y/rho); // Thickness of
   vibrating quartz at resonance, m
8 printf("\nThe thickness of vibrating quartz at
   resonance = %3.1f mm", l/1e-003);
9
10 // Result
11 // The thickness of vibrating quartz at resonance =
   0.9 mm
```

Chapter 18

ACOUSTICS OF BUILDINGS

Scilab code Exa 18.1 Output power of the sound source

```
1 // Scilab Code Ex18.1: Output power of the sound
  source : Page-361 (2010)
2 r = 200; // Distance of the point of reduction
  from the source, m
3 I_0 = 1e-012; // Final intensity of sound, watt
  per metre square
4 I_f = 60; // Intensity gain of sound at the
  point of reduction, dB
5 // As  $A_I = 10 \cdot \log_{10}(I/I_0)$ , solving for I
6 I = I_0 * 10^(I_f/10); // Initial Intensity of
  sound, watt per metre square
7 P = 4 * %pi * r^2 * I; // Output power of the sound
  source, watt
8 printf("\n\nThe output power of the sound source = %3
  .1 f W", P);
9
10 // Result
11 // The output power of the sound source = 0.5 W
```

Scilab code Exa 18.2 Change in sound level for doubling intensity

```
1 // Scilab Code Ex18.2: Change in sound level for
   doubling intensity: Page-361 (2010)
2 I1 = 1; // For simplicity assume first intensity
   level to be unity, W per metre square
3 I2 = 2*I1; // Intensity level after doubling,
   watt per metre square
4 dA_I = 10*log10(I2/I1); // Difference in gain
   level, dB
5 printf("\nThe sound intensity level is increased by
   = %1d dB", dA_I);
6
7 // Result
8 // The sound intensity level is increased by = 3 dB
```

Scilab code Exa 18.3 Total absorption of sound in the hall

```
1 // Scilab Code Ex18.3: Total absorption of sound in
   the hall: Page-361 (2010)
2 V = 8000; // Volume of the hall, metre cube
3 T = 1.5; // Reverbration time of the hall, s
4 alpha_s = 0.167*V/T; // Sabine Formula giving
   total absorption of sound in the hall, OWU
5 printf("\nThe total absorption of sound in the hall
   = %5.1f OWU", alpha_s);
6
7 // Result
8 // The total absorption in the hall = 890.7 OWU
```

Scilab code Exa 18.4 Average absorption coefficient of the surfaces of the hall

```

1 // Scilab Code Ex18.4: Average absorption
  coefficient of the surfaces of the hall: Page-362
  (2010)
2 V = 25*20*8;          // Volume of the hall, metre cube
3 S = 2*(25*20+25*8+20*8); // Total surface area of
  the hall, metre square
4 T = 4;              // Reverbration time of the hall, s
5 alpha = 0.167*V/(T*S); // Sabine Formule giving
  total absorption in the hall, OWU
6 printf("\nThe total absorption in the hall = %5.3f
  OWU per metre square", alpha);
7
8 // Result
9 // The total absorption in the hall = 0.097 OWU per
  metre square

```

Scilab code Exa 18.5 Reverbration time for the hall

```

1 // Scilab Code Ex18.5: Reverbration time for the
  hall : Page-362 (2010)
2 V = 475;          // Volume of the hall, metre cube
3 s = [200, 100, 100]; // Area of wall, floor and
  ceiling of the hall resp., metre square
4 T = 4;          // Reverbration time of the hall, s
5 alpha = [0.025, 0.02, 0.55]; // Absorption
  coefficients of the wall, ceiling and floor resp
  ., OWU per metre square
6 alpha_s = 0;
7 for i=1:1:3
8     alpha_s = alpha_s + alpha(i)*s(i);
9 end
10 T = 0.167*V/alpha_s; // Sabine Formula for
  reverbration time, s
11 printf("\nThe reverbration time for the hall = %4.2f
  s", T);

```

```
12
13 // Result
14 // The reverbration time for the hall = 1.28 s
```

Scilab code Exa 18.6 Gain of resultant sound intensity

```
1 // Scilab Code Ex18.6: Gain of resultant sound
  intensity: Page-362 (2010)
2 I0 = 1; // For simplicity assume initial sound
  intensity to be unity, watt per metre square
3 A_I1 = 80; // First intensity gain of sound, dB
4 A_I2 = 70; // Second intensity gain of sound, dB
5 // As  $A_I = 10 \cdot \log_{10}(I/I_0)$ , solving for I1 and I2
6 I1 = 10^(A_I1/10)*I0; // First intensity of sound
  , watt per metre square
7 I2 = 10^(A_I2/10)*I0; // Second intensity of
  sound, watt per metre square
8 I = I1 + I2; // Resultant intensity level of
  sound, watt per metre square
9 A_I = 10*log10(I/I0); // Intensity gain of
  resultant sound, dB
10 printf("\nThe intensity gain of resultant sound = %6
  .3f dB", A_I);
11
12 // Result
13 // The intensity gain of resultant sound = 80.414 dB
```
