

Scilab Textbook Companion for
Physics- For Students Of Science And
Engineering(Part 2)
by D. Halliday and R. Resnick¹

Created by
Byragoni Spandana
Physics
Physics
IIT-B
College Teacher
Na
Cross-Checked by
Lavitha Pereira

July 30, 2019

¹Funded by a grant from the National Mission on Education through ICT, <http://spoken-tutorial.org/NMEICT-Intro>. This Textbook Companion and Scilab codes written in it can be downloaded from the "Textbook Companion Project" section at the website <http://scilab.in>

Book Description

Title: Physics- For Students Of Science And Engineering(Part 2)

Author: D. Halliday and R. Resnick

Publisher: Toppan, Tokyo, Japan

Edition: 2

Year: 1962

ISBN: 62-15336

Scilab numbering policy used in this document and the relation to the above book.

Exa Example (Solved example)

Eqn Equation (Particular equation of the above book)

AP Appendix to Example(Scilab Code that is an Appednix to a particular Example of the above book)

For example, Exa 3.51 means solved example 3.51 of this book. Sec 2.3 means a scilab code whose theory is explained in Section 2.3 of the book.

Contents

| | |
|----------------------------------|----|
| List of Scilab Codes | 4 |
| 26 CHARGE AND MATTER | 5 |
| 27 THE ELECTRIC FIELD | 8 |
| 28 GAUSS S LAW | 11 |
| 29 ELECTRIC POTENTIAL | 13 |
| 30 CAPACITORS AND DIELECTRICS | 16 |
| 31 CURRENT AND RESISTANCE | 19 |
| 33 THE MAGNETIC FIELD | 23 |
| 34 AMPERES LAW | 27 |
| 35 FARADAYS LAW | 30 |
| 36 INDUCTANCE | 33 |
| 37 MAGNETIC PROPERTIES OF MATTER | 37 |
| 38 ELECTROMAGNETIC OSCILLATIONS | 40 |

| | |
|---|----|
| 39 ELECTROMAGNETIC WAVES | 43 |
| 40 NATURE AND PROPOGATION OF LIGHT | 44 |
| 41 REFLECTION AND REFRACTION PLANE WAVES AND PLANE SURFACESREFLECTION AND REFRACTION PLANE WAVES AND PLANE SURFACES | 47 |
| 42 REFLECTION AND REFRACTION SPHERICAL WAVES AND SPHERICAL SURFACES | 49 |
| 43 INTERFERENCE | 52 |
| 44 DIFFRACTION | 54 |
| 45 GRATING AND SPECTRA | 56 |
| 46 POLARIZATION | 60 |
| 47 LIGHT AND QUANTUM PHYSICS | 62 |
| 48 WAVES AND PROPOGATION | 65 |

List of Scilab Codes

| | | |
|-----------|---|----|
| Exa 26.1 | Magnitude of total charges in a copper penny | 5 |
| Exa 26.2 | Seperation between total positive and negative charges | 5 |
| Exa 26.3 | Force acting on charge q1 | 6 |
| Exa 26.4 | Electrical and Gravitational force between two particles | 6 |
| Exa 26.5 | Repulsive force betwwen two protons in a nucleus of iron | 7 |
| Exa 27.1 | Electric field strength | 8 |
| Exa 27.4 | The point on the line joining two charges for the electric field strength to be zero | 8 |
| Exa 27.9 | Deflection of electron | 9 |
| Exa 27.11 | Torque and work done by external agent on electric dipole | 9 |
| Exa 28.3 | Electric field strength | 11 |
| Exa 28.4 | Electric field strength at the nuclear surface | 11 |
| Exa 29.3 | Magnitude of an isolated positive point charge | 13 |
| Exa 29.4 | Electric potential at the surface of a gold nucleus | 13 |
| Exa 29.5 | Potential at the center of the square | 14 |
| Exa 29.8 | Mutual potential energy | 14 |
| Exa 29.9 | Mutual potential energy | 15 |
| Exa 30.1 | Plate area | 16 |
| Exa 30.5 | To calculate Capacitance Free charge Electric field strength Potential diffrence between plates | 16 |
| Exa 30.6 | To calculate Electric displacement and Electric polarisation in dielectric and air gap | 17 |

| | | |
|----------|---|----|
| Exa 31.1 | Current density | 19 |
| Exa 31.2 | Drift speed | 20 |
| Exa 31.3 | Resistance and resistivity | 20 |
| Exa 31.4 | Mean time and Mean free path | 21 |
| Exa 31.5 | Power | 21 |
| Exa 33.1 | Force acting on a proton | 23 |
| Exa 33.3 | Torsional constant of the spring | 23 |
| Exa 33.4 | Work done | 24 |
| Exa 33.5 | Hall potential difference | 24 |
| Exa 33.6 | Orbital radius Cyclotron frequency and Pe- riod of revolution | 25 |
| Exa 33.7 | Magnetic induction and Deuteron energy . . | 26 |
| Exa 34.3 | Distance | 27 |
| Exa 34.5 | Magnetic field and Magnetic flux | 27 |
| Exa 34.9 | Magnetic field and Magnetic dipole moment | 28 |
| Exa 35.1 | Induced EMF | 30 |
| Exa 35.7 | Induced electric field and EMF | 31 |
| Exa 36.1 | Inductance of a toroid | 33 |
| Exa 36.2 | Time | 33 |
| Exa 36.3 | MAximum Current and Energy stored . . . | 34 |
| Exa 36.4 | Rate at which energy is stored and delivered and appeared | 34 |
| Exa 36.6 | Energy | 35 |
| Exa 37.2 | Orbital dipole moment | 37 |
| Exa 37.4 | Change in magnetic moment | 37 |
| Exa 37.5 | Precession frequency | 38 |
| Exa 37.6 | Magnetic field strength Magnetisation Effec- tive magnetising current and Permeability . | 38 |
| Exa 38.1 | Current | 40 |
| Exa 38.2 | Angular frequency | 40 |
| Exa 38.3 | Angular frequency and time | 41 |
| Exa 38.5 | Magnetic field | 41 |
| Exa 38.6 | Calculation of current | 42 |
| Exa 39.6 | Magnitude of electric and magnetic field . . | 43 |
| Exa 40.1 | Force and energy reflected | 44 |
| Exa 40.2 | Angular speed | 44 |
| Exa 40.3 | Calculation of c | 45 |
| Exa 40.4 | Percentage error | 45 |

| | | |
|----------|--|----|
| Exa 41.1 | angel between two refracted beam | 47 |
| Exa 41.4 | Index of glass | 48 |
| Exa 41.5 | Angel | 48 |
| Exa 42.4 | Location of image | 49 |
| Exa 42.5 | Location of image | 50 |
| Exa 42.7 | Location of image | 50 |
| Exa 42.8 | Location of image | 50 |
| Exa 43.1 | Angular position of first minimum | 52 |
| Exa 43.2 | Linear distance | 52 |
| Exa 43.4 | Refraction | 53 |
| Exa 43.5 | Refraction | 53 |
| Exa 44.1 | Calculation of wavelength | 54 |
| Exa 44.2 | Calculation of wavelength | 54 |
| Exa 44.5 | Current | 55 |
| Exa 44.7 | delta y | 55 |
| Exa 45.1 | calculation of angel | 56 |
| Exa 45.2 | Calculation of angel theta | 56 |
| Exa 45.3 | Calculation of sodium doublet | 57 |
| Exa 45.4 | Calculation of dispersion | 57 |
| Exa 45.5 | Calculation of angels | 58 |
| Exa 46.1 | Calculation of theta | 60 |
| Exa 46.2 | Angle of refraction | 60 |
| Exa 46.3 | Thickness of slab | 61 |
| Exa 47.1 | velocity | 62 |
| Exa 47.2 | Time calculation | 62 |
| Exa 47.3 | Work function for sodium | 63 |
| Exa 47.4 | Kinetic energy to be imparten on recoiling electron | 63 |
| Exa 47.5 | Binding energy of hydrogen atom | 64 |
| Exa 1.1 | Wavelength of particle | 65 |
| Exa 48.1 | velocity and angular wavelength | 65 |
| Exa 48.2 | Quantized energy | 66 |
| Exa 48.3 | Quantum number | 66 |
| Exa 48.5 | Position of electron | 67 |
| Exa 48.6 | Position of electron | 67 |

Chapter 26

CHARGE AND MATTER

Scilab code Exa 26.1 Magnitude of total charges in a copper penny

```
1 //chapter26
2 //Example 1.1
3 clc
4 m=3.1 //mass of copper penny in grams
5 e=4.6*10^-18 //charge in coulombs
6 N0=6*10^23 //avogadro's number atoms/mole
7 M=64 //molecular weight of copper in gm/mole
8 //Calculation
9 N=(N0*m)/M; //No.of copper atoms in penny
10 q=N*e; //magnitude of the charges in coulombs
11 disp(q,"magnitude of the charges in coul is")
```

Scilab code Exa 26.2 Separation between total positive and negative charges

```
1 //Chapter26
2 //Example 2
3 clc
4 F=4.5 //Force of attraction in nt
```

```

5 q=1.3*10^5 //total charge in coul
6 r=q*sqrt((9*10^9)/F);
7 disp(r,"Seperation between total positive and
    negative charges in meters is")

```

Scilab code Exa 26.3 Force acting on charge q1

```

1 //Chapter26
2 //example 3
3 clc
4 //given three charges q1,q2,q3
5 q1=-1.0*10^-6 //charge in coul
6 q2=+3.0*10^-6
7 q3=-2.0*10^-6
8 r12=15*10^-2 //seperation between q1 and q2 in m
9 r13=10*10^-2 //seperation between q1 and q3 in m
10 angle=%pi/6 //in degrees
11 F12=(9.0*10^9)*q1*q2/(r12^2) //in nt
12 F13=(9.0*10^9)*q1*q3/(r13^2) //in nt
13 F12x=-F12 ; //ignoring signs of charges
14 F13x=F13*sin(angle);
15 F1x=F12x+F13x
16 F12y=0 //from fig.263
17 F13y=-F13*cos(angle);
18 F1y=F12y+F13y //in nt
19 disp(F1x,"X component of resultant force acting on
    q1 in nt is")
20 disp(F1y,"Y component of resultant force acting on
    q1 in nt is")

```

Scilab code Exa 26.4 Electrical and Gravitational force between two particles

```

1 //chapter26

```

```

2 //Example 4
3 clc
4 r=5.3*10^-11 //distance b/w electron and proton in
   the hydrogen atom in meter
5 e=1.6*10^-19 //charge in coul
6 G=6.7*10^-11 //gravitatal constant in nt-m2/kg2
7 m1=9.1*10^-31 //mass of electron in kg
8 m2=1.7*10^-27 //mass of proton in kg
9 F1=(9*10^9)*e*e/(r^2) //coulombs law
10 F2=G*m1*m2/(r^2) //gravitational force
11 disp(F1,"Coulomb force in nt is")
12 disp(F2,"Gravitational force in nt is")

```

Scilab code Exa 26.5 Repulsive force betwten two protons in a nucleus of iron

```

1 //chapter 26
2 //Example 5
3 clc
4 r=4*10^-15 //separation b/w proton annd nucleus in
   iron in meters
5 q=1.6*10^-19 //charge in coul
6 F=(9*10^9)*(q^2)/(r^2); //coulombs law
7 printf("Repulsive coulomb force F=%d nt",F)

```

Chapter 27

THE ELECTRIC FIELD

Scilab code Exa 27.1 Electric field strength

```
1 //chapter27
2 //Example 1
3 clc
4 m=9.1*10^-31 //mass of electron in kg
5 g=9.8 //acceleration due to gravity in m/s
6 q=1.6*10^-19 //charge of electron in coul
7 disp("Electric field strength E=F/q where F=mg")
8 E=m*g/q
9 disp(E,"electric field strength in nt/coul is")
```

Scilab code Exa 27.4 The point on the line joining two charges for the electric fi

```
1 //chapter27
2 //example 4
3 clc
4 //given
5 q1=1.0*10^-6 //in coul
6 q2=2.0*10^-6 //in coul
```

```

7 l=10 //separation b/w q1 and q2 in cm
8 disp("for the electric field strength to be zero the
      point should lie between the charges where E1=E2
      ")
9 // "Refer to the fig 27.9"
10 //E1=electric field strength due to q1
11 //E2=electric field strength due to q2
12 disp("E1=E2 which implies  $q_1/4x^2 = q_2/4(1-x)^2$ ")
13 x=1/(1+sqrt(q2/q1))
14 printf("Electric field strength is zero at x=%f cm",
      x)

```

Scilab code Exa 27.9 Deflection of electron

```

1 //chapter27
2 //example 9
3 clc
4 //given
5 e=1.6*10^-19 //charge in coul
6 E=1.2*10^4 //electric field in nt/coul
7 x=1.5*10^-2 //length of deflecting assembly in m
8 K0=3.2*10^-16 //kinetic energy of electron in joule
9 //calculation
10 y=e*E*x^2/(4*K0)
11 disp(y,"Corresponding deflection in meters is")

```

Scilab code Exa 27.11 Torque and work done by external agent on electric dipole

```

1 //chapter 27
2 //Example 11
3 clc
4 //Given

```

```

5 q=1.0*10^-6 //magnitude of two opposite charges of a
   electric dipole in coul
6 d=2.0*10^-2 // seperation b/w charges in m
7 E=1.0*10^5 //external field in nt/coul
8 //calculations
9 //(a)Max torque if found when theta=90 degrees
10 //Torque =pEsin(theta)
11 p=q*d //electric dipole moment
12 T=p*E*sin(%pi/2)
13 disp("(a)Maximum torque exerted by the fied in nt-m
   is")
14 disp(T)
15 //(b)work done by the external agent is the
   potential energy b/w the positions theta=180 and
   0 degree
16 W=(-p*E*cos(%pi))-(-p*E*cos(0))
17 disp("(b) work done by the external agent to turn
   dipole end for end in joule is ")
18 disp(W)

```

Chapter 28

GAUSS S LAW

Scilab code Exa 28.3 Electric field strength

```
1 //chapter 28
2 //Example 3
3 clc
4 //Given
5 r=1*10^-10 //radius of the atom in meter
6 Z=79 //gold atomic number
7 e=1.6*10^-19 //charge in coul
8 q=Z*e //total positive charge in coul
9 E=(9.0*10^9)*q/r^2
10 disp(E,"Electric field strength at the surface of
    the gold atom in nt/coul is")
```

Scilab code Exa 28.4 Electric field strength at the nuclear surface

```
1 //chapter 28
2 //Example 4
3 clc
4 //given
```

```
5 r=6.9*10^-15 //radius of the gold nucleus in meter
6 Z=79 //gold atomic number
7 e=1.6*10^-19 //charge in coul
8 q=Z*e //total positive charge in coul
9 E=(9.0*10^9)*q/r^2
10 disp("Electric field strength at the surface of the
      gold atom in nt/coul is")
11 disp(E)
```

Chapter 29

ELECTRIC POTENTIAL

Scilab code Exa 29.3 Magnitude of an isolated positive point charge

```
1 //chapter 29
2 //example 3
3 clc
4 //given
5 V=100 //electric potential in volts
6 r=10*10^-2 //in meters
7 epsilon0=8.85*10^-12 //coul2/nt-m2
8 disp("Potential due to a point charge is  $V=q/4*\pi*$ 
      epislon0*r")
9 q=V*4*%pi*epsilon0*r
10 disp(q,"Magnitude of positive point charge in coul
      is ")
```

Scilab code Exa 29.4 Electric potential at the surface of a gold nucleus

```
1 //chapter 29
2 //example 4
3 clc
```

```

4 //given
5 r=6.6*10^-15 //radius of the gold nucleus in meter
6 Z=79 //gold atomic number
7 e=1.6*10^-19 //charge in coul
8 q=Z*e //total positive charge in coul
9 epsilon0=8.85*10^-12 //coul2/nt-m2
10 V=q/(4*pi*epsilon0*r)
11 disp(V,"Electric potential at the surface of the
    nucleus in volts is")

```

Scilab code Exa 29.5 Potential at the center of the square

```

1 //chapter 29
2 //example 5
3 clc
4 //given
5 q1=1.0*10^-8 //in coul
6 q2=-2.0*10^-8 //in coul
7 q3=3.0*10^-8 //in coul
8 q4=2.0*10^-8 //in coul
9 a=1 //side of square in meter
10 epsilon0=8.85*10^-12 //coul2/nt-m2
11 //refer to the fig 29.7
12 r=a/sqrt(2) //distance of charges from centre in
    meter
13 V=(q1+q2+q3+q4)/(4*pi*epsilon0*r)
14 disp(V,"Potential at the center of the square in
    volts is")

```

Scilab code Exa 29.8 Mutual potential energy

```

1 //chapter 29
2 //example 8

```

```

3  clc
4  //given
5  q1=1.6*10^-19 //charge in coul
6  q2=1.6*10^-19 //charge in coul
7  r=6.0*10^-15 //seperation b/w two protons in meter
8  epsilon0=8.85*10^-12 //coul2/nt-m2
9  U=(q1*q2)/(4*pi*epsilon0*r)
10 disp("Mutual electric potential energy of two proton
      in joules is")
11 disp(U)
12 V=U/q1
13 disp(V,"Mutual electric potential energy of two
      proton in ev is")

```

Scilab code Exa 29.9 Mutual potential energy

```

1  //chapter 29
2  //example 9
3  clc
4  //given
5  q=1.0*10^-7 //charge in coul
6  a=10*10^-2 //side of triangle in meter
7  q1=q
8  q2=-4*q
9  q3=2*q
10 epsilon0=8.85*10^-12 //coul2/nt-m2
11 disp("Total energy is the sum of each pair of
      particles ")
12 U=(1/(4*pi*epsilon0))*(((q1*q2)/a)+((q1*q3)/a)+((q2
      *q3)/a))
13 disp(U,"Mutual potential energy of the particles in
      joules is")

```

Chapter 30

CAPACITORS AND DIELECTRICS

Scilab code Exa 30.1 Plate area

```
1 //chapter 30
2 //example 1
3 clc
4 //given
5 C=1.0 //capacitance in farad
6 d=1.0*10^-3 //separation b/w plates in meter
7 epsilon0=8.85*10^-12 //coul2/nt-m2
8 A=d*C/epsilon0
9 disp(A,"Plate area in square meter is")
```

Scilab code Exa 30.5 To calculate Capacitance Free charge Electric field strength

```
1 //chapter 30
2 //example 5
3 clc
4 //given
```

```

5  epsilon0=8.85*10^-12 //coul2/nt-m2
6  A=100*10^-4//area of the plate in square meter
7  d=1*10^-2 //separation b/w plates in meter
8  b=5*10^-3 //thickness of dielectric lab in meter
9  V0=100//in volts
10 k=7
11 //(a)
12 C0=epsilon0*A/d
13 disp(C0,"(a) Capacitance before the slab is inserted
    in farad is")
14 //(b)
15 q=C0*V0
16 disp(q,"(b) Free charge in coul is")
17 //(c)
18 E0=q/(epsilon0*A)
19 disp(E0,"(c) Electric field strength in the gap in
    volts/meter is")
20 //(d)
21 E=q/(k*epsilon0*A)
22 disp(E,"(d) Electric field strength in the dielectric
    in volts/meter is")
23 //(e)
24 //Refer to fig30-12
25 V=E0*(d-b)+E*b
26 disp(V,"(e) Potential difference between the plates
    in volts is")
27 //(f)
28 C=q/V
29 disp(C,"Capacitance with the slab in place in farads
    is")

```

Scilab code Exa 30.6 To calculate Electric displacement and Electric polarisation

```

1 //chapter 30
2 //example 6

```

```

3  clc
4  //given
5  epsilon0=8.85*10^-12 //coul2/nt-m2
6  A=100*10^-4//area of the plate in square meter
7  d=1*10^-2 //separation b/w plates in meter
8  V0=100//in volts
9  E0=1*10^4 //Electric field in the air gap in volts/
    meter
10 k=7
11 k0=1
12 E=1.43*10^3 //in volts/metre
13 D=k*E*epsilon0
14 P=epsilon0*(k-1)*E
15 //(a)
16 disp(D,"(a) Electric displacement in dielectric in
    coul/square metre is ")
17 disp(P," Electric polarisation in dielectric in coul/
    square meter is")
18 //(b)
19 D0=k0*epsilon0*E0
20 disp(D0,"(b) Electric displacement in air gap in coul
    /square metre is ")
21 P0=epsilon0*(k0-1)*E0
22 disp(P0," Electric polarisation in air gap in coul/
    square meter is")

```

Chapter 31

CURRENT AND RESISTANCE

Scilab code Exa 31.1 Current density

```
1 //chapter 31
2 //example1
3 clc
4 //given
5 d1=0.10 //diameter of aluminium wire in inches
6 d2=0.064 //diameter of copper wire in inches
7 i=10 //current carried by composite wire in amperes
8 A1=%pi*(d1/2)^2 //crosssectional area of aluminium
   wire in square inches
9 A2=%pi*(d2/2)^2 //crosssectional area of copper wire
   in square inches
10 j1=i/A1
11 j2=i/A2
12 disp(j1,"Current density in Aluminium wire in amp/
   square inches")
13 disp(j2,"Current density in copper wire in amp/
   square inches")
```

Scilab code Exa 31.2 Drift speed

```
1 //chapter 31
2 //example2
3 clc
4 //given
5 j=480//current density for copper wire in amp/cm2
6 N0=6*10^23 //avagadro number in atoms/mole
7 M=64//molecular wt in gm/mole
8 d=9.0 //density in gm/cm3
9 e=1.6*10^-19//elecron charge in coul
10 n=d*N0/M
11 disp(n,"No.of free electrons per unit volume in
        atoms/mole")
12 Vd=j/(n*e)
13 disp(Vd,"Drift speed of electron in cm/sec is")
```

Scilab code Exa 31.3 Resistance and resistivity

```
1 //chapter 31
2 //example3
3 clc
4 //given
5 disp("Dimensions of rectangular carbon block are 1.0
        cm*1.0cm*50cm")
6 l=1.0*10^-2 //in meter
7 b=1.0*10^-2//in meter
8 h=50*10^-2 //in meter
9 p=3.5*10^-5 //resisivity of carbon in ohm-m
10 //(a)Resistance b/w two square ends
11 l1=h
12 A1=b*l
```



```

13 R1=p*l1/A1
14 disp(R1,"(a) Resistance measured b/w the two square
    ends in ohm is")
15 l2=1
16 A2=b*h
17 R2=p*l2/A2
18 disp(R2,"(a) Resistance measured b/w the two
    opposite rectangular faces in ohm is")

```

Scilab code Exa 31.4 Mean time and Mean free path

```

1 //chapter 31
2 //example4
3 clc
4 //given
5 m=9.1*10^-31 //in kg
6 n=8.4*10^28 //in m-1
7 e=1.6*10^-19 //in coul
8 p=1.7*10^-8 //in ohm-m
9 v=1.6*10^8 //in cm/sec
10 T=2*m/(n*p*e^2)
11 disp(T,"(a) Mean time b/w collisions in sec is")
12 Lambda=T*v
13 disp(Lambda,"(b) Mean free path in cm is")

```

Scilab code Exa 31.5 Power

```

1 //chapter 31
2 //example5
3 clc
4 //given
5 V=110 //in volt
6 R=24 //ohms

```

```
7 P1=V^2/R
8 disp(P1,"(a)Power for the single coil in watts is")
9 P2=V^2/(R/2)
10 disp(P2,"(b)Power for a coil of half the length in
    watts is")
```

Chapter 33

THE MAGNETIC FIELD

Scilab code Exa 33.1 Force acting on a proton

```
1 //chapter 33
2 //example1
3 clc
4 //given
5 K=5*10^6 //ev
6 e=1.6*10^-19 //in coul
7 K1=K*e //in joules
8 m=1.7*10^-27 //in kg
9 B=1.5 //wb/m
10 theta=%pi/2
11 v=sqrt(2*K1/m)
12 disp(v,"Speed of the proton in meters/sec is")
13 F=e*v*B*sin(theta)
14 disp(F,"Force acting on proton in nt is")
```

Scilab code Exa 33.3 Torsional constant of the spring

```
1 //chapter 33
```

```

2 //example3
3 clc
4 //given
5 N=250 //turns in coil
6 i=1.0*10^-4 //in amp
7 B=0.2 //wb/m2
8 h=2*10^-2 // coil height in m
9 w=1.0*10^-2 //width of coil in m
10 Q=30 //angular deflection in degrees
11 theta=%pi/2
12 A=h*w
13 k=N*i*A*B*sin(theta)/Q
14 disp(k,"Torssional constant in nt-m/deg is")

```

Scilab code Exa 33.4 Work done

```

1 //chapter 33
2 //example4
3 clc
4 //given
5 N=100 // turns in circular coil
6 i=0.10 //in amp
7 B=1.5 // in wb/m2
8 a=5*10^-2 //radius of coil in meter
9 u=N*i*%pi*(a^2) //u is dipole moment
10 U1=(-u*B*cos(0))
11 U2=-u*B*cos(%pi)
12 W=U2-U1
13 disp(W," Work required to turn current in an
    external magnetic field from theta=0 to theta=180
    degree in joule is ")

```

Scilab code Exa 33.5 Hall potential difference

```

1 //chapter 33
2 //example5
3 clc
4 //given
5 i=200 //current in the strip in amp
6 B=1.5 //magnetic field in wb/m2
7 n=8.4*10^28 //in m-3
8 e=1.6*10^-19 //in coul
9 h=1.0*10^-3 //thickness of copper strip in metre
10 w=2*10^-2 //width of copper strip in meter
11 //calculation
12 Vxy=i*B/(n*e*h)
13 disp(Vxy,"Hall potential difference across strip in
    volt is")

```

Scilab code Exa 33.6 Orbital radius Cyclotron frequency and Period of revolution

```

1 //chapter 33
2 //example6
3 clc
4 //given
5 m=9.1*10^-31 // in kg
6 v=1.9*10^6 //in m/sec
7 q=1.6*10^-19 //charge in coul
8 B=1.0*10^-4 //in wb/m2
9 //calculations
10 //(A)
11 r=m*v/(q*B)
12 disp(r,"(A) Orbit radius in meter is")
13 //(B)
14 f=q*B/(2*%pi*m)
15 disp(f,"(B) Cyclotron frequency in rev/sec is")
16 //(C)
17 T=1/f
18 disp(T,"(C) Period of revolution in sec is")

```

Scilab code Exa 33.7 Magnetic induction and Deuteron energy

```
1 //chapter 33
2 //example7
3 clc
4 //given
5 f0=12*10^6 //cyclotron frequency in cycles/sec
6 r=21//dee radius in inches
7 R=r*0.0254 //dee radius in meter
8 q=1.6*10^-19 //charge in coul
9 m=3.3*10^-27 //in kg
10 //(A)
11 B=2*%pi*f0*m/q
12 disp(B,"(A) Magnetic induction needed to accelerate
        deuterons in wb/m2 is")
13 //(B)
14 K=((q^2*B^2*R^2)/(2*m))
15 disp(K,"(B) Deuteron energy in joule is")
16 K1=K*(1/(1.6*10^-19))
17 disp(K1," Deuteron energy in ev is")
```

Chapter 34

AMPERES LAW

Scilab code Exa 34.3 Distance

```
1 //chapter 34
2 //example3
3 clc
4 //given
5 i1=100 //in amp
6 i2=20 //in amp
7 W=0.073 //weight of second wire  $W=F/l$  in nt/m
8 u0=4*%pi*10^-7 //in weber/amp-m
9 //calculations
10 d=u0*i1*i2/(2*%pi*W)
11 disp(d,"seperation between two wires in metres")
```

Scilab code Exa 34.5 Magnetic field and Magnetic flux

```
1 //chapter 34
2 //example5
3 clc
4 //given
```

```

5 l=1.0 //length of solenoid in meter
6 d=3*10^-2 //diameter of solenoid in meter
7 n=5*850 //number of layers and turns of wire
8 u0=4*%pi*10^-7 //in weber/amp-m
9 i0=5.0 //current in amp
10 //(A)
11 B=u0*i0*n
12 disp(B,"Magnetic field at center in wb/m2 is")
13 //(B)
14 A=%pi*(d/2)^2
15 Q=B*A
16 disp(Q,"Magnetic flux at the center of the solenoid
    in weber is")

```

Scilab code Exa 34.9 Magnetic field and Magnetic dipole moment

```

1 //chapter 34
2 //example9
3 clc
4 //given
5 e=1.6*10^-19 //in coul
6 R=5.1*10^-11 //radius of th enucleus in meter
7 f=6.8*10^15 //frequency with which electron
    circulates in rev/sec
8 u0=4*%pi*10^-7 //in weber/amp-m
9 x=0 //x is any point on the orbit , since at center x
    =0
10 //(A)
11 i=e*f
12 B=u0*i*R^2*0.5/((R^2+x^2)^(3/2))
13 disp(B,"(A) Magnetic field at the center of the
    orbit in wb/m2")
14 N=1 //no.of turns
15 A=%pi*R^2
16 U=N*i*A

```


17 `disp(U,"(B) Equivalent magnetic dipole moment in
amp-m2 is ")`

Chapter 35

FARADAYS LAW

Scilab code Exa 35.1 Induced EMF

```
1 //chapter 35
2 //example1
3 clc
4 //given
5 l=1.0 //length of solenoid in meter
6 r=3*10^-2 //radius of solenoid in meter
7 n=200*10^2 //number of turns in solenoid per meter
8 u0=4*%pi*10^-7 //in weber/amp-m
9 i=1.5 //current in amp
10 N=100 //no.of turns in a close packed coil placed at
    the center of solenoid
11 d=2*10^-2 //diameter of coil in meter
12 delta_T=0.050 //in sec
13 //(A)
14 B=u0*i*n
15 disp(B,"Magnetic field at center in wb/m2 is")
16 //(B)
17 A=%pi*(d/2)^2
18 Q=B*A
19 disp(Q,"Magnetic flux at the center of the solenoid
    in weber is")
```

```

20 delta_Q=Q-(-Q)
21 E=-(N*delta_Q/delta_T)
22 disp(E,"Induced EMF in volts is ")

```

Scilab code Exa 35.7 Induced electric field and EMF

```

1 //chapter 35
2 //example7
3 clc
4 //given
5 //refer to fig 35-16
6 B=2 //magnetic field in wb/m2
7 l=10*10^-2 //in m
8 v=1.0 //in m/sec
9 q=1.6*10^-19 //charge in coul
10 disp("Let S be the frame of reference fixed w.r.t
        the magnet and Z be the frame of reference w.r.t
        the loop")
11 //(A)
12 E=v*B
13 disp(E,"(A) Induced electric field in volt/m
        observed by Z")
14 //(B)
15 F=q*v*B
16 disp(F,"(B) Force acting on charge carrier in nt w.
        r.t S is")
17 F1=q*E
18 disp(F1," Force acting on charge carrier in nt w
        .r.t Z is")
19 //(C)
20 emf1=B*l*v
21 disp(emf1,"(C) Induced emf in volt observed by S is
        ")
22 emf2=E*l
23 disp(emf2,"Induced emf in volt observed by Z is")

```


Chapter 36

INDUCTANCE

Scilab code Exa 36.1 Inductance of a toroid

```
1 //chapter 36
2 //example1
3 clc
4 //given
5 u0=4*%pi*10^-7//in weber/amp-m   Mu-not=u0
6 N=10^3//no.of turns
7 a=5*10^-2//in meter
8 b=10*10^-2 //in meter
9 h=1*10^-2// in metre
10 L=(u0*N^2*h)/(2*%pi)*log(b/a)
11 disp(L,"Inductance of a toroid of recyangular cross
    section in henry is")
```

Scilab code Exa 36.2 Time

```
1 //chapter 36
2 //example2
3 clc
```

```

4 //given
5 L=50 //inductance in henry
6 R=30 //resistance in ohms
7 t0=log(2)*(L/R)
8 disp(t0,"Time taken for the current to reach one-
      half of its final equilibrium in sec is")

```

Scilab code Exa 36.3 MAXimum Current and Energy stored

```

1 //chapter 36
2 //example3
3 clc
4 //given
5 L=5 //inductance in henry
6 V=100 //emf in volts
7 R=20 //resistance in ohms
8 i=V/R
9 disp(i,"Maximum current in amp is")
10 U=(L*i^2)/2
11 disp(U,"Energy stored in the magnetic field in
      joules is")

```

Scilab code Exa 36.4 Rate at which energy is stored and delivered and appeared

```

1 //chapter 36
2 //example4
3 clc
4 //given
5 L=3//inductance in henry
6 R=10 //resistance in ohm
7 V=3 //emf in volts
8 t=0.30//in sec
9 T=0.30 //inductive time constant in sec

```

```

10 //(a)
11 i=(V/R)*(1-exp(-t/T))
12 P1=V*i
13 disp(P1,"The rate at which energy is delivred by the
    battery in watt is")
14 //(b)
15 P2=i^2*R
16 disp(P2,"The rate at which energy appears as Joule
    heat in the resistor in watt is")
17 //(c)
18 disp(" Let D=di/dt")
19 D=(V/L)*exp(-t/T) // in amp/sec
20 P3=L*i*D
21 disp(P3,"The desired rate at which energy is being
    stored in the magnetic field in watt is")

```

Scilab code Exa 36.6 Energy

```

1 //chapter 36
2 //example6
3 clc
4 //given
5 epsilon0=8.9*10^-12//in coul2/nt-m2
6 E=10^5//elelectric field in volts/meter
7 B=1 //magnetic field in weber/meter2
8 u0=4*%pi*10^-7//in weber/amp-m Mu-not=u0
9 a=0.1 //side of the cube in meter
10 V0=a^3 //volume of the cube in meter3
11 //(a)
12 U1=epsilon0*E^2*V0/2 //in elelectric field
13 disp(U1,"(a)Energy required to set up in the given
    cube on edge in electric field in joules is")
14 //(b)
15 U2=(B^2/(2*u0))*V0
16 disp(U2,"(b)Energy required to set up in the given

```

cube on edge in magnetic field in joules is”)

Chapter 37

MAGNETIC PROPERTIES OF MATTER

Scilab code Exa 37.2 Orbital dipole moment

```
1 //chapter 37
2 //example2
3 clc
4 //given
5 e=1.6*10^-19 //in coul
6 r=5.1*10^-11 //radius of hydrogen atom in meter
7 m=9.1*10^-31 // mass of electron in kg
8 epsilon0=8.9*10^-12 //in coul2/nt-m2
9 p=((e^2)/4)*sqrt(r/(%pi*epsilon0*m))
10 disp(p,"Orbital dipole moment in amp-m2 is")
```

Scilab code Exa 37.4 Change in magnetic moment

```
1 //chapter 37
2 //example4
3 clc
```

```

4 //given
5 e=1.6*10^-19 //in coul
6 r=5.1*10^-11 //radius of hydrogen atom in meter
7 m=9.1*10^-31 // mass of electron in kg
8 epsilon0=8.9*10^-12 //in coul2/nt-m2
9 B=2 //in wb/m2
10 delta_p=(e^2*B*r^2)/(4*m)
11 disp(delta_p,"Change in Orbital dipole moment in amp
    -m2 is + 0r -")

```

Scilab code Exa 37.5 Precession frequency

```

1 //chapter 37
2 //example5
3 clc
4 //given
5 u=1.4*10^-26 //in amp-m2
6 B=0.50 //wb/m2
7 Lp=0.53*10^-34 //in joule-sec
8 fp=u*B/(2*%pi*Lp)
9 disp(fp,"Precession frequency of phoyon in given
    magnetic field in cps is")

```

Scilab code Exa 37.6 Magnetic field strength Magnetisation Effective magnetising c

```

1 //chapter 37
2 //example6
3 clc
4 //given
5 n=10*10^2 //turns/m
6 i=2 //in amp
7 B=1.0 //in wb/m
8 u0=4*%pi*10^-7 //in wb/amp-m

```

```

9 // (A)
10 H=n*i
11 disp(H,"(A) Magnetic field strength in amp/m is")
12 // (B)
13 M=(B-u0*H)/u0
14 disp(" (B) Magnetisation is Zero when core is
        removed")
15 disp(M," Magnetisation when th ecore is replaced in
        amp/m")
16 // (C)
17 disp(" (C) Effective magnetizing current i=i(M,0)=M
        *(2*pi*r0/N0)=M/n")
18 i=M/n
19 disp(i," Effective magnetizing current in amp is")
20 // (D)
21 Km=B/(u0*H)
22 disp(Km," (D) Permeability ")

```

Chapter 38

ELECTROMAGNETIC OSCILLATIONS

Scilab code Exa 38.1 Current

```
1 //Example 1
2 //Chapter 38
3 //clc()
4 V_o=50// in volts
5 C=1*10^-6 //in farad
6 L=10*10^-3
7 i_m=V_o*(sqrt(C/L))
8 disp(i_m,"Max current in amps")
```

Scilab code Exa 38.2 Angular frequency

```
1 //chapter 38
2 //Example 2
3 clc
4 //given
5 L=10*(10^-3)// in henry
```

```
6 C=(10)^-6 //in farad
7 w=sqrt(1/(L*C))
8 disp(" Angular frequency in radians/sec=")
9 disp(w)
```

Scilab code Exa 38.3 Angular frequency and time

```
1 //chapter 38
2 //Example 3
3 clc
4
5
6 //given
7 L=10*(10^-3) // in henry
8 C=(10)^-6 //in farad
9 R=0.1 //in ohm
10 w=sqrt(1/(L*C))
11 disp(" Angular frequency in radians/sec=")
12 disp(w)
13 t=(2*L*log(2))/R
14 disp(" time in sec=")
15 disp(t)
```

Scilab code Exa 38.5 Magnetic field

```
1 //chapter 38
2 //Example 5
3 clc
4 //given
5 m_0=(4*%pi*10^-7) //in weber
6 e_0=(8.9*10^-12)
7 R=5*10^-2 //meters
8 dEbydT=10^12
```

```
9 B=(0.5*m_0*e_0*R*dEbydT)
10 disp(" magnetic field in weber/m^2=")
11 disp(B)
```

Scilab code Exa 38.6 Calculation of current

```
1 //chapter 38
2 //Example 6
3 clc
4 //given
5 m_0=(4*%pi*10^-7)//in weber
6 e_0=(8.9*10^-12)
7 R=5*10^-2//meters
8 dEbydT=10^12
9 i_d=(e_0*%pi*R*R*dEbydT)
10 disp(" current in amp=")
11 disp(i_d)
```

Chapter 39

ELECTROMAGNETIC WAVES

Scilab code Exa 39.6 Magnitude of electric and magnetic field

```
1 //Example 6
2 //chapter 39
3 //clc ()
4 r=1//in m
5 p=10^3//
6 m=4*%pi*10^-7//weber/amp-m
7 c=3*10^8
8 x=2*%pi
9 E_m=(1/r)*(sqrt((p*m*c)/x))
10 disp(E_m,"The value of E in volts/meter=")
11 B=E_m/c
12 disp(B,"B in weber/meter^2")
```

Chapter 40

NATURE AND PROPOGATION OF LIGHT

Scilab code Exa 40.1 Force and energy reflected

```
1 //chapter 40
2 //Example 1
3 //clc()
4 u=(10)*(1.0)*3600// in Joules
5 c=3*10^8// in m/sec
6 t=3600//in sec
7 disp("solution (a)")
8 disp(u,"energy reflected from mirror in joule=")
9 p=(2*u)/c
10 disp(p,"momentum after i hr illumination in kg-m/sec
    =")
11 disp("solution (b)")
12 f=p/t
13 disp(f,"force in newton=")
```

Scilab code Exa 40.2 Angular speed


```

1 //chapter 40
2 // example 2
3 //clc ()
4 theta=1/1440
5 c=3*10^8// in m/sec
6 l=8630 //in m
7 w=(c*theta)/(2*l)
8 disp(w,"Angular speed in rev/sec= ")

```

Scilab code Exa 40.3 Calculation of c

```

1 //chapter 40
2 //example 3
3 //clc ()
4 l=15.6//in cm
5 n=8
6 lambda_g=(2*l)/n
7 disp(lambda_g,"lambda_g in cm=")
8 lambda=3.15//in cm
9 f=9.5*10^9//cycles/sec
10 c=lambda*f
11 disp(c,"value of c in m/sec=")

```

Scilab code Exa 40.4 Percentage error

```

1 //Example 4
2 //chapter 4
3 //clc ()
4 v_1=25000//miles/hr
5 u=25000//miles/hr
6 c=6.7*10^8// miles/hr
7 x=1+((v_1*u)/(c)^2)
8

```

```
9 v=(v_1+u)/x
10 disp(v,"Speed of light in miles/hour=")
```

Chapter 41

REFLECTION AND REFRACTION PLANE WAVES AND PLANE SURFACES REFLECTION AND REFRACTION PLANE WAVES AND PLANE SURFACES

Scilab code Exa 41.1 angel between two refracted beam

```
1 //example_2
2 //chapter 41
3 theta_1=30
4 n_qa=1.4702
5 theta2=asind(sind(theta_1)/n_qa)
6 disp("For 4000 A beam, theta_2 in degree=")
7 disp(theta2)
8
```

```
9 theta_1=30
10 n_qa=1.4624
11 theta2=asind(sind(theta_1)/n_qa)
12 disp("For 5000 A beam, theta_2 in degree=")
13 disp(theta2)
```

Scilab code Exa 41.4 Index of glass

```
1 //example 4
2 // chapter 41
3 //clc()
4 disp("Index reflection=")
5 n=1/sind(45)
6 disp(n)
```

Scilab code Exa 41.5 Angel

```
1 //Example 4
2 //Chapter 41
3 //clc()
4 n2=1.33
5 n1=1.50
6 theta_c=asind(n2/n1)
7 disp(theta_c,"Angle theta_c in degree=")
8 disp("Actual angle of indices =45 is less than
      theta_c, so there \n is no internal angle
      reflection")
9 disp("angle of refraction=")
10 x=n1/n2
11 theta_2=asind(x*sind(45))
12 disp(theta_2,"theta_2 in degree=")
```

Chapter 42

REFLECTION AND REFRACTION SPHERICAL WAVES AND SPHERICAL SURFACES

Scilab code Exa 42.4 Location of image

```
1 //example_4
2 //chapter 42
3 //clc ()
4 n1=1
5 n2=2
6 o=20//in cm
7 r=10//in cm
8 disp("x=n2/i")
9 x=((n2-n1)/r)-(n1/o)
10 disp(x)
11 i=n2/x
12 disp(i,"The value of i in cm=")
```

Scilab code Exa 42.5 Location of image

```
1 //example_5
2 //chapter 42
3 //clc()
4 n1=2
5 n2=1
6 o=15//in cm
7 r=-10//in cm
8 disp("x=n2/i")
9 x=((n2-n1)/r)-(n1/o)
10 disp(x)
11 i=n2/x
12 disp(i,"The value of i in cm=")
```

Scilab code Exa 42.7 Location of image

```
1 //Example 7
2 //chapter 42
3 //clc()
4 n=1.65
5 r_1=40//in cm
6 r_2=-40//in cm
7 disp("x=1/f in cm=")
8 x=(n-1)*((1/r_1)-(1/r_2))
9 disp(x)
10 disp("f=1/x")
11 f=1/x
12 disp(f,"f in cm=")
```

Scilab code Exa 42.8 Location of image

```
1 //Example 8
```

```
2 //chapter 42
3 //clc()
4 o=9//in c
5 f=24//in cm
6 x=(1/f)-(1/o)
7 disp("x=1/i in cm=")
8 disp(x)
9 i=1/x
10 disp("i in cm=")
11 disp(i)
12 disp("lateral magnification =")
13 m=-(i/o)
14 disp(m)
```

Chapter 43

INTERFERENCE

Scilab code Exa 43.1 Angular position of first minimum

```
1 //Example
2 //chapter 43
3 //given
4 m=1
5 lambda=546*10^-9
6 d=0.10*10^-3//in m
7 sin_theta=((m-0.5)*lambda)/(d)
8 disp(sin_theta,"sin theta =")
9 theta=asind(sin_theta)
10 disp(theta,"angle in degree=")
```

Scilab code Exa 43.2 Linear distance

```
1 //Example 2
2 //chapter 43
3 delta=546*10^-9//in meter
4 D=20*10^-2//in meter
5 d=0.10*10^-3//in meter
```



```
6 delta_y=(delta*D)/d
7 disp(delta_y,"Linear distance in meter=")
```

Scilab code Exa 43.4 Refraction

```
1 //Example 4
2 //chapter 43
3 d=3200//in A
4 n=1.33
5 for m=1:2
6     lambda_max=(2*d*n)/(m+0.5)
7     lambda_min=(8500/m)
8     disp(m,"when m=")
9     disp(lambda_max,"lambda_max=")
10    disp(lambda_min,"lambda_min=")
11 end
```

Scilab code Exa 43.5 Refraction

```
1 //Example 5
2 //chapter 43
3 //clc()
4 lambda=5000//in A
5 n=1.38
6 for m=0:3
7     disp(m,"when m=")
8     d=((m+0.5)*lambda)/(2*n)
9     disp(d,"d in A=")
10 end
```

Chapter 44

DIFFRACTION

Scilab code Exa 44.1 Calculation of wavelength

```
1 //example 1
2 //chapter 44
3 //clc ()
4 m=1
5 lambda=6500//in A
6
7 a=(m*lambda)/sind(30)
8 disp(a,"a in A=")
```

Scilab code Exa 44.2 Calculation of wavelength

```
1 //Example 2
2 //Chapter 44
3 lambda=6500
4 lambda_1=lambda/1.5
5 disp(lambda_1,"wavelength in A=")
```

Scilab code Exa 44.5 Current

```
1 //chapter 44
2 //Example 5
3 clc
4 //given
5 m_0=(4*%pi*10^-7)//in weber
6 e_0=(8.9*10^-12)
7 R=5*10^-2//meters
8 dEbydT=10^12
9 i_d=(e_0*%pi*R*R*dEbydT)
10 disp(" current in amp=")
11 disp(i_d)
```

Scilab code Exa 44.7 delta y

```
1 //chapter 44
2 //example 7
3 //given
4 //clc()
5
6 lambda=480*10^-9//in m
7 d=0.10*10^-3//in m
8 D=50*10^-2// in m
9 a=0.02*10^-3
10 delta_y=(lambda*D)/d
11 disp(" solution (a)")
12 disp(delta_y,"D in m=")
```

Chapter 45

GRATING AND SPECTRA

Scilab code Exa 45.1 calculation of angel

```
1 //example 1
2 //chapter 45
3 //clc ()
4 m=1
5 lambda=4000//in A
6 d=31700//in A
7 theta=asind((m*lambda)/d)
8 disp(theta,"The first order diffraction pattern in
   degree=")
```

Scilab code Exa 45.2 Calculation of angel theta

```
1 //chapter 45
2 //example 2
3 //given
4 //clc ()
5 m=1
6 lambda=5890//in A
```

```

7 d=25400//in A
8 theta=asind((m*lambda)/d)
9 disp("solution (a)")
10 disp(theta,"The first order diffraction pattern in
    degree=")
11 disp("solution (b)")
12 //given
13 del_lambda=5.9//in A
14 delta_theta=(m*(del_lambda))/(d*cosd(theta))
15 disp(delta_theta,"Angle of seperation in degree=")

```

Scilab code Exa 45.3 Calculation of sodium doublet

```

1 //Example 3
2 //chapter 45
3 //Given
4 lambda=5890//A
5 m=3
6 delta_lambda=(5895.9-5890.0)//in A
7 R=lambda/(delta_lambda)
8 disp(R,"Resolving power=")
9 N=(R/m)
10 disp(N,"Number of rulings needed is=")

```

Scilab code Exa 45.4 Calculation of dispersion

```

1 //Example 4
2 //Chapter 45
3 //clc ()
4 m=3
5 m1=5
6 lambda=5460//in A
7 d=31700//in A

```

```

8 theta=asind((m*lambda)/d)
9 disp(theta,"The first order diffraction pattern in
    degree=")
10 D=m/(d* cosd(theta))
11 disp("Solution (a)")
12 disp(D,"The dispersion in radian/A=")
13 disp("Solution (b)")
14 N=8000
15 lambda=5460
16 R=N*m1
17 delta_lambda=lambda/R
18 disp(delta_lambda,"Wave length difference in A=")

```

Scilab code Exa 45.5 Calculation of angels

```

1 //Example 5
2 //chapter 45
3 //clc()
4 a_o=5.63//A
5 d=a_o/sqrt(5)
6 lambda=1.10//in A
7 disp(d,"Interplanar spacing d in A=")
8
9 disp("diffracted beam occurs when m=1,m=2 and m=3")
10 disp("when m1=1, theta in degree=")
11 m1=1
12 x=(m1*lambda)/(2*d)
13 theta_1=asind(x)
14 disp(theta_1)
15 disp("when m1=2, theta in degree=")
16 m2=2
17 x=(m2*lambda)/(2*d)
18 theta_2=asind(x)
19 disp(theta_2)
20 disp("when m1=3, theta in degree=")

```

```
21 m3=3
22 x=(m3*lambda)/(2*d)
23 theta_3=asind(x)
24 disp(theta_3)
```

Chapter 46

POLARIZATION

Scilab code Exa 46.1 Calculation of theta

```
1 //example 1
2 //chapter 46
3 theta=acosd(1/sqrt(2))
4 disp(180-theta,"Polarization angle theta=")
```

Scilab code Exa 46.2 Angle of refraction

```
1 //example 2
2 // Chapter 46
3 theta_p= atand(1.5)
4 disp(theta_p,"theta_p in degrees")
5 sin_theta_r= sind(theta_p)/1.5
6 theta_r=asind(sin_theta_r)
7 disp(theta_r,"angle of refraction from snells law in
degrees=")
```

Scilab code Exa 46.3 Thickness of slab

```
1 //Example 3
2 //chapter 46
3 // from the equation
4 lambda=5890//A
5 n_e=1.553
6 n_o=1.544
7 s=(n_e)-(n_o)
8 x=(lambda)/(4*s)
9 //textbook answer is wrong
10 disp(x,"The value of x in m=")
```

Chapter 47

LIGHT AND QUANTUM PHYSICS

Scilab code Exa 47.1 velocity

```
1 //Chapter 47
2 //Example1
3 k=20//in nt/m
4 m=1//in kg
5 //solution
6 v=(sqrt((k)/(m)))*(1/(2*%pi))
7
8 disp(" velocity in cycles/s" )
9 disp(v)
```

Scilab code Exa 47.2 Time calculation

```
1 //Example 2
2 //chapter 47
3 P=(10-3)*(3*10-18)/(300)
4 disp(P,"Power in j-sec")
```

```

5 s=1.6*(10^(-19))
6 t=(5*s)/P
7 disp(t,"time required in sec =")
8 one_sec=0.000277778// hr
9 in_hour=one_sec*t
10 disp(in_hour,"time required in hour")

```

Scilab code Exa 47.3 Work function for sodium

```

1 //Example_3
2 //Chapter 47
3 h=6.63*10^(-34)// in joule/sec
4 v=4.39*10^(14)// cycles/sec
5 E_o=h*(v)
6 disp(E_o,"Energy in joule=")

```

Scilab code Exa 47.4 Kinetic energy to be imparted on recoiling electron

```

1 //Example 4
2 disp("solution a")
3 h=(6.63)*10^-34
4 m=9.11*10^-31
5 c=3*10^8
6 delta_h=(h/(m*c))*(1-cos(90))
7 disp(delta_h,"compton shift in meter")
8 disp("solution b")
9 delta=1*10^-10
10 k=(h*c*delta_h)/(delta*(delta+delta_h))
11
12 disp(k,"Kinetic energy in joules")

```

Scilab code Exa 47.5 Binding energy of hydrogen atom

```
1 //example5
2 m=9.11*10^-31//in kg
3 e=8.85*10^-12// in coul^2/nt-m^2
4 h=6.63*10^-34//in j-sec
5 E=(-m*(e^4))/(8*(e^2)*(h^2))
6 disp(E,"Binding energy of hydrogen in joule")
```

Chapter 48

WAVES AND PROPOGATION

Scilab code Exa 1.1 Wavelength of particle

```
1 //Chapter 48
2 //Example1
3 k=100*(1.6*(10^-19))
4 m=9.1*(10^-31)
5 //solution
6 v=sqrt(((2*k)/(m)))
7
8 disp(" velocity in m/s" )
9 disp(v)
10 h=6.6*(10^-34)
11 p=5.4*(10^-34)
12 lambda=h/p
13 disp(" wavelength in A")
14 disp(lambda)
```

Scilab code Exa 48.1 velocity and angular wavelength

```

1 //Chapter 48
2 //Example1
3 k=100*(1.6*(10^-19))
4 m=9.1*(10^-31)
5 //solution
6 v=sqrt(((2*k)/(m)))
7
8 disp(" velocity in m/s" )
9 disp(v)
10 h=6.6*(10^-34)
11 p=5.4*(10^-34)
12 lambda=h/p
13 disp(" wavelength in A")
14 disp(lambda)

```

Scilab code Exa 48.2 Quantized energy

```

1 //Example 2
2 //given data
3 n=1
4 h=(6.6)*10^-34 //j/sec
5 m=9.1*(10^-31)//in kg
6 l=1*(10^-9)//in m
7 //Solution
8 E=(n^2)*((h^2)/(8*m*(l^2)))
9 disp(" Energy in Joule=")
10 disp(E)

```

Scilab code Exa 48.3 Quantum number

```

1 //Example3
2 //given
3 m=10^-9//in kg

```

```

4 v=10^-6//in m/s
5 l=10^-4//in m
6 h=(6.6)*(10^-34)//j/s
7 E=(0.5)*m*(v^2)
8
9 disp("Energy in joule=")
10 disp(E)
11
12 n=(1/h)*(sqrt(8*m*E))
13 disp("Quantum number=")
14 disp(n)

```

Scilab code Exa 48.5 Position of electron

```

1 //Example 5
2 //given
3 m=9.1*(10^-31)//in kg
4 v=300//in m/s
5 h=6.6*(10^-34)// in j-s
6 p=m*v
7 disp("The electrom momentum in kg-m/s=")
8 disp(p)
9 delta_p=(0.0001)*p
10
11 disp("delta_p in kg-m/s=")
12 disp(delta_p)
13
14 delta_x=(h/delta_p)
15 disp("Minimum uncertainty in m=")
16 disp(delta_x)

```

Scilab code Exa 48.6 Position of electron

```
1 //Example_6
2 m=0.05// in kg
3 v=300 //m/s
4 delta_p=m*v
5 disp("Momentum in kg-m/s=")
6 disp(delta_p)
7 delta_x=(6.6*10^-34)/delta_p
8 disp("delta_x in meter=")
9 disp(delta_x)
```
