

Scilab Textbook Companion for  
Modern Physics  
by R. A. Serway<sup>1</sup>

Created by  
Nusrat Ali  
Modern Physics  
Electronics Engineering  
Model Institute of Engineering and Technology  
College Teacher  
Ms. Bhavna Sharma  
Cross-Checked by  
Lavitha Pereira

July 31, 2019

<sup>1</sup>Funded by a grant from the National Mission on Education through ICT, <http://spoken-tutorial.org/NMEICT-Intro>. This Textbook Companion and Scilab codes written in it can be downloaded from the "Textbook Companion Project" section at the website <http://scilab.in>

# Book Description

**Title:** Modern Physics

**Author:** R. A. Serway

**Publisher:** Thomson Learning, USA

**Edition:** 3

**Year:** 2005

**ISBN:** 0-534-49339-4

Scilab numbering policy used in this document and the relation to the above book.

**Exa** Example (Solved example)

**Eqn** Equation (Particular equation of the above book)

**AP** Appendix to Example(Scilab Code that is an Appednix to a particular Example of the above book)

For example, Exa 3.51 means solved example 3.51 of this book. Sec 2.3 means a scilab code whose theory is explained in Section 2.3 of the book.

# Contents

|   |    |
|---|----|
| List of Scilab Codes                    | 4  |
| 1 Relativity I                          | 5  |
| 2 Relativity II                         | 11 |
| 3 The Quantum theory of light           | 18 |
| 4 The particle nature of matter         | 26 |
| 5 Matter waves                          | 35 |
| 6 Quantum mechanics in one dimension    | 40 |
| 7 Tunnelling phenomena                  | 49 |
| 8 Quantum Mechanics in Three Dimensions | 53 |
| 9 Atomic Structure                      | 57 |
| 10 Statistical Physics                  | 62 |
| 11 Molecular Structure                  | 68 |
| 12 The Solid State                      | 71 |

|  |           |
|--|-----------|
| <b>13 Nuclear Structure</b>            | <b>74</b> |
| <b>14 Nuclear Physics Applications</b> | <b>81</b> |
| <b>15 Elementary Particle</b>          | <b>86</b> |
| <b>16 Cosmology</b>                    | <b>93</b> |

# List of Scilab Codes

|          |  |    |
|----------|--|----|
| Exa 1.2  | Period of the pendulum wrt different frames of references . . . . .          | 5  |
| Exa 1.3  | Contraction of spaceship . . . . .   | 5  |
| Exa 1.4  | Altitude of spaceship wrt different frames of references . . . . .           | 6  |
| Exa 1.5  | Shape of spaceship seen from different frames of references . . . . .        | 7  |
| Exa 1.6  | Speed of recession of the galaxy Hydra . . . . .                             | 7  |
| Exa 1.8  | Relative velocity of spaceships . . . . .                                    | 8  |
| Exa 1.9  | Velocity of ball wrt stationary observer . . . . .                           | 9  |
| Exa 1.10 | Relative velocity of recession of two gang leaders . . . . .                 | 9  |
| Exa 2.1  | Momentum of an electron . . . . .  | 11 |
| Exa 2.3  | Energy of a speedy electron . . . . .  | 12 |
| Exa 2.4  | Energy of a speedy proton . . . . .  | 12 |
| Exa 2.5  | Increase in mass of colliding balls . . . . .                                | 13 |
| Exa 2.7  | A Fission Reaction . . . . .   | 14 |
| Exa 2.8  | Energy conservation . . . . .  | 15 |
| Exa 2.9  | Mass of Pion . . . . .   | 16 |
| Exa 3.1  | Temperature of sun . . . . .   | 18 |
| Exa 3.2  | Quantum oscillator vs classical oscillator . . . . .                         | 18 |
| Exa 3.3  | Stefan law from Planck distribution . . . . .                                | 20 |
| Exa 3.4  | Time lag between start of illumination and photocurrent generation . . . . . | 20 |
| Exa 3.5  | Time lag between start of illumination and photocurrent generation . . . . . | 21 |
| Exa 3.6  | Photoelectric effect for iron . . . . .                                      | 21 |
| Exa 3.7  | Compton shift for carbon . . . . .   | 22 |

|          |   |    |
|----------|---|----|
| Exa 3.8  | Xray photons vs visible photons . . . . .                                 | 23 |
| Exa 3.9  | Gravitational redshift for a white dwarf . . . . .                        | 24 |
| Exa 4.1  | Electrolysis of barium chloride . . . . .                                 | 26 |
| Exa 4.2  | Deflection of electron beam by E and B Fields . . . . .                   | 27 |
| Exa 4.3  | Experimental determination of e . . . . .                                 | 27 |
| Exa 4.4  | Collision of alpha particle with proton . . . . .                         | 30 |
| Exa 4.5  | Radius of Aluminium Nucleus . . . . .                                     | 31 |
| Exa 4.7  | Collision of alpha particle with proton . . . . .                         | 31 |
| Exa 4.8  | series for Hydrogen . . . . .   | 32 |
| Exa 4.9  | Hydrogen in its first excited state . . . . .                             | 33 |
| Exa 5.1  | Wave properties of a baseball . . . . .                                   | 35 |
| Exa 5.2  | de Broglie wavelength of an electron . . . . .                            | 35 |
| Exa 5.3  | Diffraction of neutrons at the crystal lattice . . . . .                  | 36 |
| Exa 5.8  | Uncertainty principle for macroscopic objects . . . . .                   | 37 |
| Exa 5.9  | Kinetic energy of electron confined within the nucleus . . . . .          | 37 |
| Exa 5.10 | Width of spectral lines . . . . .   | 38 |
| Exa 6.2  | Probability from wave function . . . . .                                  | 40 |
| Exa 6.4  | Dispersion of matter waves . . . . .                                      | 40 |
| Exa 6.5  | Energy Quantization for Macroscopic Object . . . . .                      | 41 |
| Exa 6.6  | Model of an Atom . . . . .  | 42 |
| Exa 6.7  | Probabilities for a particle in a Box . . . . .                           | 43 |
| Exa 6.8  | Ground state energy of an electron confined to a potential well . . . . . | 44 |
| Exa 6.12 | The quantum oscillator in nonclassical region . . . . .                   | 45 |
| Exa 6.13 | Quantization of vibrational energy . . . . .                              | 45 |
| Exa 6.14 | Standard Deviations from Averages . . . . .                               | 46 |
| Exa 6.15 | Location of a particle in the box . . . . .                               | 47 |
| Exa 7.1  | Transmission coefficient for an oxide layer . . . . .                     | 49 |
| Exa 7.2  | Tunnelling current through an oxide layer . . . . .                       | 50 |
| Exa 7.5  | Tunnelling in a parallel plate capacitor . . . . .                        | 50 |
| Exa 7.6  | Estimating halflives of Thorium and Polonium . . . . .                    | 51 |
| Exa 8.4  | Orbital quantum number for a stone . . . . .                              | 53 |
| Exa 8.6  | Space quantisation for an atomic electron . . . . .                       | 54 |
| Exa 8.7  | Energy of Hydrogen atom at first excited state . . . . .                  | 55 |
| Exa 8.8  | Probabilities for the Electron in Hydrogen . . . . .                      | 55 |
| Exa 9.1  | Magnetic energy of electron in Hydrogen . . . . .                         | 57 |

|           |  |    |
|-----------|--|----|
| Exa 9.2   | Angles between z axis and the spin angular momentum vector . . . . .             | 57 |
| Exa 9.3   | Zeeman Spectrum of Hydrogen Including Spin                                       | 58 |
| Exa 9.4   | Spin orbit energy of Sodium doublet . . . . .                                    | 59 |
| Exa 9.5   | Ground state of Helium atom . . . . .  | 60 |
| Exa 9.6   | Effective atomic number for 3s electron in Na                                    | 60 |
| Exa 10.1  | Population of excited states with respect to ground states in Hydrogen . . . . . | 62 |
| Exa 10.2  | Validity of Maxwell Boltzmann Statistics . .                                     | 63 |
| Exa 10.3  | Photons in a box . . . . .   | 65 |
| Exa 10.4  | Specific Heat of Diamond . . . . .   | 65 |
| Exa 10.5  | Fermi Energy of Gold . . . . .   | 66 |
| Exa 11.1  | Rotation of CO molecule . . . . .  | 68 |
| Exa 11.2  | Variation of CO molecule . . . . .   | 69 |
| Exa 12.1  | Classical free electron model . . . . .  | 71 |
| Exa 12.2  | Conduction in diamond . . . . .  | 72 |
| Exa 12.3  | Forward and reverse currents in diode . . . .                                    | 73 |
| Exa 13.1  | The Atomic Mass Unit . . . . .   | 74 |
| Exa 13.2  | The Volume and Density of Nucleus . . . . .                                      | 74 |
| Exa 13.3  | Binding energy of the Deuteron . . . . .   | 75 |
| Exa 13.4  | Left out sample during radioactive decay . .                                     | 75 |
| Exa 13.5  | The Activity of Radium . . . . .   | 76 |
| Exa 13.6  | The Activity of Carbon . . . . .   | 77 |
| Exa 13.7  | The Radiactive Isotope of Iodine . . . . .                                       | 78 |
| Exa 13.8  | Energy Liberated during Decay of Radium .  | 79 |
| Exa 13.9  | Probability of Alpha Decay . . . . .   | 79 |
| Exa 13.11 | Radioactive Dating . . . . .   | 80 |
| Exa 14.1  | Energy released in Fission . . . . .   | 81 |
| Exa 14.2  | Neutron capture by Al . . . . .  | 82 |
| Exa 14.4  | Energy released in the Fission of U235 . . . .                                   | 82 |
| Exa 14.5  | A Rough Mechanism for Fission Process . . .                                      | 83 |
| Exa 14.6  | The Fusion of Two Deutrons . . . . .   | 84 |
| Exa 14.7  | Half value thickness . . . . .   | 85 |
| Exa 15.2  | Checking Baryon Numbers . . . . .  | 86 |
| Exa 15.3  | Checking Lepton Numbers . . . . .  | 88 |
| Exa 15.4  | Conservation of strangeness . . . . .  | 90 |
| Exa 15.5  | Making virtual particle real . . . . .   | 92 |
| Exa 16.1  | Hubbles law . . . . .  | 93 |



|          |  |    |
|----------|--|----|
| Exa 16.2 | Critical density of universe . . . . . | 94 |
|----------|--|----|

# Chapter 1

## Relativity I

Scilab code Exa 1.2 Period of the pendulum wrt different frames of references

```
1 // Scilab code Ex1.2: Pg.18 (2005)
2 clc; clear;
3 c = 3e+08; // Velocity of light , m/s
4 v = 0.95*c; // Velocity of observer , m/s
5 T_proper = 3; // Proper time period of pendulum
   in rest frame , s
6 gama = 1/(sqrt(1 - (v/c)^2)); // Multiplying
   factor
7 // From time-dilation formula , we have
8 T = gama*T_proper; // Time period
   of pendulum w.r.t to moving observer , s
9 printf("\\nTime period of pendulum w.r.t to moving
   observer = %3.1f s", T);
10
11 // Result
12 // Time period of pendulum w.r.t to moving observer
   = 9.6 s
```

---

Scilab code Exa 1.3 Contraction of spaceship

```

1 // Scilab code Ex1.3: Pg 20 (2005)
2 clc; clear;
3 c = 3e+08; // Velocity of light , m
  /s
4 L_p = 100; // Proper length of
  spaceship , m
5 v = 0.99*c; // Velocity of spaceship
  , m/s
6 // Using length contraction formula ,
7 L = L_p*sqrt(1 - (v/c)^2); //
  Observed length of spaceship , m
8 printf("Observed length of spaceship = %2d m", L);
9
10 // Result
11 // Observed length of spaceship = 14 m

```

---

**Scilab code Exa 1.4** Altitude of spaceship wrt different frames of references

```

1 // Scilab code Ex1.4: Pg 20 (2005)
2 clc; clear;
3 c = 3e+08; // Velocity of light , m
  /s
4 L_p = 435; // Proper altitude of
  spaceship , m
5 v = 0.970*c; // Velocity of
  spaceship , m/s
6 // Using length contraction formula ,
7 L = L_p*sqrt(1 - (v/c)^2); //
  Observed altitude of spaceship , m
8 printf("Observed altitude of spaceship = %2d m",
  ceil(L));
9
10 // Result
11 // Observed altitude of spaceship = 106 m

```

---

**Scilab code Exa 1.5** Shape of spaceship seen from different frames of references

```
1 // Scilab code Ex1.5: Pg 20 (2005)
2 clc; clear;
3 c = 3e+08; // Velocity of light , m
  /s
4 L_p = 50; // Proper distance
  between points x & y of spaceship , m
5 v = 0.950*c; // Velocity of
  spaceship , m/s
6 // Using length contraction formula ,
7 L = L_p*sqrt(1 - (v/c)^2); //
  Observed distance between points x & y of
  spaceship , m
8 printf("\\nObserved distance between points x and y
  of spaceship = %4.1f m", L);
9 printf("\\nThe spaceship will get contracted in the
  direction of motion");
10
11 // Result
12 // Observed distance between points x and y of
  spaceship = 15.6 m
13 // The spaceship will get contracted in the
  direction of motion
```

---

**Scilab code Exa 1.6** Speed of recession of the galaxy Hydra

```
1 // Scilab code Ex1.6: Pg 25 (2005)
2 clc; clear;
3 // For simplification assume velocity of light equal
  to unity
4 c = 1 // Velocity of light , m/s
```

```

5 lamda_obs = 474e-09;           // Wavelength measured
  by observer , m
6 lamda_source = 394e-09;       // Wavelength measured
  in the source's rest frame, m
7 v = ((lamda_obs^2 - lamda_source^2)/(lamda_obs^2 +
  lamda_source^2))*c;           //
  Receding velocity of Hydra, m/s
8 printf("\nReceding velocity of Hydra = %5.3fc m/s",
  v);
9
10 // Result
11 // Receding velocity of Hydra = 0.183c m/s

```

---

#### Scilab code Exa 1.8 Relative velocity of spaceships

```

1 // Scilab code Ex1.8: Pg 30 (2005)
2 clc; clear;
3 // For simplification assume velocity of light equal
  to unity
4 c = 1;           // Velocity of light , m/s
5 v = 0.750*c;    // Velocity of spaceship A
  relative to S frame, m/s
6 u_x = (-0.850)*c; // Velocity of spaceship B
  relative to S frame, m/s
7 // Using Lorentz velocity transformation
8 U_x = (u_x - v)/(1 - u_x*v/c^2); //
  Velocity of spaceship B with respect to spaceship
  A, m/s
9 printf("\nVelocity of spaceship B with respect to
  spaceship A = %6.4fc m/s", U_x);
10
11 // Result
12 // Velocity of spaceship B with respect to spaceship
  A = -0.9771c m/s

```

---

### Scilab code Exa 1.9 Velocity of ball wrt stationary observer

```
1 // Scilab code Ex1.9: Pg 30 (2005)
2 clc; clear;
3 // For simplification assume velocity of light equal
  to unity
4 c = 1; // Velocity of light , m/s
5 v = 0.800*c; // Velocity of motorcycle w
  .r.t stationary observer , m/s
6 U_x = 0.700*c; // Velocity of ball in the
  reference frame of motorcyclist , m/s
7 // Using inverse Lorentz velocity transformation
8 u_x = (U_x + v)/(1 + U_x*v/ c^2); //
  Velocity of ball relative to stationary observer
  , m/s
9 printf("\\nVelocity of ball relative to stationary
  observer = %6.4fc m/s", u_x);
10
11 // Result
12 // Velocity of ball relative to stationary observer
  = 0.9615c m/s
```

---

### Scilab code Exa 1.10 Relative velocity of recession of two gang leaders

```
1 // Scilab code Ex1.10: Pg 30–31 (2005)
2 clc; clear;
3 // For simplification assume velocity of light equal
  to unity
4 c = 1; // Velocity of light , m/s
5 ux = 0.75*c; // Velocity of pack
  leader alpha , m/s
6 gama = 1/(sqrt(1 - (ux/c)^2));
```

```

7  u_x = 0; // Velocity component of
    beta measured in S frame, m/s
8  U_x = (u_x - ux)/(1 - u_x*ux/c^2); //
    Velocity component of beta along X-axis measured
    in S' frame, (Velocity Addition Rule), m/s
9  u_y = -0.90*c; // Velocity
    component of beta long Y-axis measured in S frame
10 U_y = u_y/(gama*(1 - u_x*ux/c^2)); // Velocity
    component of beta along Y-axis measured in S'
    frame, m/s
11 U = sqrt(U_x^2+U_y^2); // Relative velocity of
    recession of two gang leaders, m/s
12 printf("\nThe relative velocity of recession of two
    gang leaders = %4.2 fc", U);
13
14 // Result
15 // The relative velocity of recession of two gang
    leaders = 0.96c

```

---

# Chapter 2

## Relativity II

Scilab code Exa 2.1 Momentum of an electron

```
1 // Scilab code Ex2.1: Pg.44 (2005)
2 clc; clear;
3 c = 3e08; // Velocity of light , m/s
4 u = 0.750*c; // Velocity of electron , m/s
5 m = 9.11e-31; // Rest mass of electron , kg
6 p_r = m*u/(sqrt(1 - (u/c)^2)); // Relativistic
    momentum of electron , kgm/s
7 p = m*u; // Classical momentum of electron , kg-m/s
8 printf("\\nThe relativistic momentum of electron = %4
    .2fe-22 kg-m/s", p_r*1e+22);
9 printf("\\nThe classical momentum of electron = %4.2
    fe-22 kg-m/s", p*1e+22);
10 printf("\\nThe relativistic result is 50 percent
    greater than the classical result.");
11
12 //Result
13 // The relativistic momentum of electron = 3.10e-22
    kg-m/s
14 // The classical momentum of electron = 2.05e-22 kg-
    m/s
15 // The relativistic result is 50 percent greater
```



than the classical result.

---

### Scilab code Exa 2.3 Energy of a speedy electron

```
1 // Scilab code Ex2.3: Pg.47 (2005)
2 clc; clear;
3 c = 3e+08; // Velocity of light, m/s
4 u = 0.85*c; // Velocity of electron, m/s
5 E_0 = 0.511; // Rest energy of electron, MeV
6 E = E_0/(sqrt(1-(u/c)^2)); // Total energy of
  electron, MeV
7 K = E - E_0; // Kinetic energy of electron, MeV
8 printf("\nThe total energy of electron = %5.3f MeV",
  E);
9 printf("\nThe kinetic energy of electron = %5.3f MeV
  ", K);
10
11 // Result
12 // The total energy of electron = 0.970 MeV
13 // The kinetic energy of electron = 0.459 MeV
```

---

### Scilab code Exa 2.4 Energy of a speedy proton

```
1 // Scilab code Ex2.4: Pg.47 (2005)
2 clc; clear;
3 c = 3e+08; // Velocity of light, m/s
4 u = 0.85*c; // Velocity of electron, m/s
5 m_p = 1.67e-27; // Rest mass of proton, kg
6
7 // Part (a)
8 E_o = m_p*c^2/1.602e-019; // Rest energy of proton
  , MeV
```

```

9 printf("\nRest energy of proton = %3d MeV", E_o/1e
    +06);
10
11 // Part (b)
12 // Since given that  $E = 3 * E_o = (3 * m_p * c^2) / \sqrt{1 - (u/c)^2}$ , solving for u
13 u = sqrt(8/9)*c; // Velocity of proton, m/s
14 printf("\nVelocity of proton = %4.2 fe+08 m/s", u*1e
    -08);
15
16 // Part (c)
17 // Since  $K = E - m_p * (c^2) = 3 * m_p * (c^2) - m_p * (c^2) = 2 * m_p * (c^2)$ 
18 K = 2 * E_o; // Kinetic energy of proton, MeV
19 printf("\nThe kinetic energy of proton = %4d MeV", K
    *1e-06);
20
21 // Part (d)
22 p = sqrt(8) * (E_o);
23 printf("\nThe momentum of proton = %4d MeV/c", p*1e
    -06);
24
25 // Result
26 // Rest energy of proton = 938 MeV
27 // Velocity of proton = 2.83e+08 m/s
28 // The kinetic energy of proton = 1876 MeV
29 // The momentum of proton = 2653 MeV/c

```

---

### Scilab code Exa 2.5 Increase in mass of colliding balls

```

1 // Scilab code Ex2.5: Pg.49 (2005)
2 clc; clear;
3 u = 450; // Velocity of each ball, m/s
4 m = 5; // Mass of each ball, kg
5 c = 3e+08; // Velocity of light, m/s

```

```

6 // Since  $(u/c)^2 \ll 1$ , therefore, substituting  $1/\sqrt{1 - (u/c)^2} = 1 + (1/2)*(u/c)^2$ 
7 delta_M = m*(u/c)^2; // Increase in the mass of
  balls, kg
8 printf("\nIncrease in the mass of the balls = %3.1fe
  -11 kg", delta_M*1e+11);
9
10 // Result
11 // Increase in the mass of the balls = 1.1e-11 kg

```

---

### Scilab code Exa 2.7 A Fission Reaction

```

1 // Scilab code Ex2.7: Pg.50 (2005)
2 clc; clear;
3 u = 1.660e-27; // Atomic mass unit
4 M_U = 236.045563; // Atomic mass of Uranium, u
5 M_Rb = 89.914811; // Atomic mass of Rubidium, u
6 M_Cs = 142.927220; // Atomic mass of Caesium, u
7 m_n = 1.008665; // Mass of neutrons, u
8
9 // Part (a)
10 printf("\nU(92,235) --> Rb(37,90) + Cs(55,143) + 3n
  (0,1)");
11 printf("\nSo three neutrons are produced per fission
  .\n");
12
13 // Part (b)
14 delta_M = (M_U - (M_Rb + M_Cs + 3*m_n))*u; //
  Combined mass of all products, kg
15 printf("\nCombined mass of all products = %6.4fe-28
  kg\n", delta_M*1e+28);
16
17 // Part (c)
18 // For simplification let velocity of light = 1 m/s
19 c = 1; // Velocity of light, m/s

```

```

20 // Since  $1u = 931.5 \text{ MeV}/(c^2)$ , therefore
21 Q = (delta_M/u)*931.5*(c^2); // Energy given out
    per fission event, MeV
22 printf("\nEnergy given out per fission event = %5.1f
    MeV\n", Q);
23
24 // Part (d)
25 N = ((6.02e23)*1000)/236; // Number of nuclei
    present
26 efficiency = 0.40;
27 E = efficiency*N*Q*(4.45e-20); // Total energy
    released, kWh
28 printf("\nTotal energy released = %4.2fe+06 kWh\n",
    E*1e-06);
29
30 printf("\nThis amount of energy will keep a 100-W
    lightbulb burning for %d years", E
    *1000/(100*24*365));
31
32 // Result
33 //  $U(92,235) \rightarrow Rb(37,90) + Cs(55,143) + 3n(0,1)$ 
34 // So three neutrons are produced per fission.
35 // Combined mass of all products =  $2.9471e-28 \text{ kg}$ 
36 // Energy given out per fission event =  $165.4 \text{ MeV}$ 
37 // Total energy released =  $7.51e+06 \text{ kWh}$ 
38 // This amount of energy will keep a 100-W lightbulb
    burning for 8571 years

```

---

### Scilab code Exa 2.8 Energy conservation

```

1 // Scilab code Ex2.8: Pg.52 (2005)
2 clc; clear;
3
4 // Part (a)
5 // Since  $1 \text{ eV} = 1.6e-19 \text{ J}$ , therefore  $3 \text{ eV} = 3*1.6e$ 
```

```

-19
6 BE = 3*1.6e-19; // Binding energy of water, J
7 c = 3e+08; // Velocity of light, m/s
8 delta_m = BE/(c^2); // Mass difference of water
  molecule & it constituents, kg
9 printf("\nMass difference of water molecule & it
  constituents = %3.1fe-36 kg", delta_m*1e+36);
10
11 // Part (b)
12 M = 3.0e-26; // Mass of water molecule, kg
13 M_f = delta_m/M; // Fractional loss of mass per
  molecule
14 printf("\nThe fractional loss of mass per molecule =
  %3.1fe-10", M_f*1e+10);
15
16 // Part (c)
17 E = M_f*(c^2); // Energy released when 1 g of
  water is formed, kJ
18 printf("\nEnergy released when 1 g of water is
  formed = %2.0f kJ", E*1e-06);
19
20 // Result
21 // Mass difference of water molecule & it
  constituents = 5.3e-36 kg
22 // The fractional loss of mass per molecule = 1.8e
  -10
23 // Energy released when 1 g of water is formed = 16
  kJ

```

---

### Scilab code Exa 2.9 Mass of Pion

```

1 // Scilab code Ex2.9: Pg.53 (2005)
2 clc; clear;
3 K_mu = 4.6; // Kinetic energy of muon, MeV
4 // For convinience let m_mew*(c^2) = E_mew

```

```
5 E_mu = 106;          // Energy of muon, MeV
6 E_pion = sqrt((E_mu^2) + (K_mu^2) + (2*K_mu*E_mu)) +
    sqrt((K_mu^2) + (2*K_mu*E_mu));
7 m_pion = E_pion;    // Mass of pion, MeV/(c^2)
8 printf("\nMass of Pion = %3.0f MeV/(c^2)", m_pion);
9
10 // Result
11 // Mass of Pion = 142 MeV/(c^2)
```

---

# Chapter 3

## The Quantum theory of light

Scilab code Exa 3.1 Temperature of sun

```
1 // Scilab code Ex3.1: Pg 69 (2005)
2 clc; clear;
3 e_total = 1400; // Total power per unit area, W/(m
   ^2)
4 sigma = 5.6e-08; // Stefan-Boltzmann constant
5 R = 1.5e+11; // Earth-sun distance, m
6 R_s = 7.0e+08; // Radius of sun, m
7 // Using Stefan's law & solving for T, we get
8 T = (e_total*R^2/(sigma*R_s^2))^0.25; //
   Temperature of sun, K
9 printf("\n\nThe temperature of sun = %4d K", T);
10
11 // Result
12 // The temperature of sun = 5820 K
```

---

Scilab code Exa 3.2 Quantum oscillator vs classical oscillator

```
1 // Scilab code Ex3.2: Pg 75 (2005)
```

```

2  clc; clear;
3
4  // Part (a)
5  h = 6.63e-34;    // Plank's constant, Js
6  c = 3e+08;     // Velocity of light, m/s
7  lamda_green = 540e-09;    // Wavelength of green
    light, nm
8  delta_E_green = h*c/lamda_green/1.602e-19;    //
    Minimum energy change in green light, eV
9  lamda_red = 700e-09;    // Wavelength of red light,
    nm
10 delta_E_red = h*c/lamda_red/1.602e-19;    // Minimum
    energy change in red light, eV
11
12 printf("\nMinimum energy change in green light = %4
    .2f eV", delta_E_green);
13 printf("\nMinimum energy change in red light = %4.2f
    eV", delta_E_red);
14
15 // Part (b)
16 f = 0.50;    // Frequency, Hz
17 m = 0.1;    // Mass of pendulum, kg
18 l = 1;    // Length of pendulum, m
19 theta = %pi/180*10;    // Angle, radians
20 g = 9.8;    // Acceleration due to gravity, m/s^2
21 E = m*g*l*(1-cos(theta));
22 delta_E = (h*f)/(1.6e-19);    // Minimum energy
    change in pendulum, eV
23 delta_E_f = (delta_E*1.6e-19)/E ;    // Fractional
    energy change
24 printf("\nFractional energy change = %3.1fe-32",
    delta_E_f*1e+32);
25
26 // Result
27 // Minimum energy change in green light = 2.30 eV
28 // Minimum energy change in red light = 1.77 eV
29 // Fractional energy change = 2.2e-32

```

---



### Scilab code Exa 3.3 Stefan law from Planck distribution

```
1 // Scilab code Ex3.3: Pg 80 (2005)
2 clc; clear;
3 k_B = 1.381e-23; // Boltzmann's constant, J/K
4 c = 3e+08; // Velocity of light, m/s
5 h = 6.626e-34; // Plank's constant, Js
6 // Since  $e_{total} = \sigma \cdot (T^4) = (2 \cdot (\%pi)^5 \cdot (k_B)^4) / (15 \cdot (c^2) \cdot (h^3)) \cdot T^4$ 
7  $\sigma = (2 \cdot (\%pi)^5 \cdot (k_B)^4) / (15 \cdot (c^2) \cdot (h^3));$ 
8 printf("\\nThe value of sigma = %3.2fe-08 W/Sq.m/K^4"
9     , sigma*1e+08);
10
11 // Result
12 // The value of sigma = 5.67e-08 W/Sq.m/K^4
```

---

### Scilab code Exa 3.4 Time lag between start of illumination and photocurrent genera

```
1 // Scilab code Ex3.4: Pg 83 (2005)
2 clc; clear;
3 phi = 2.38; // Work function for sodium, eV
4 I = 1e-07; // Absorbed light intensity, mJcm^2/s
5 A = %pi*1e-16; // Cross-sectional area, m^2
6 t = phi*1.6e-16/(I*A) // Time lag, days
7 printf("\\nTime lag between start of illumination and
8     photocurrent generation = %3.1fe+07 s", t*1e-07)
9     ;
10
11 // Result
12 // Time lag between start of illumination and
13     photocurrent generation = 1.2e+07 s
```

---

**Scilab code Exa 3.5** Time lag between start of illumination and photocurrent genera

```
1 // Scilab code Ex3.5: Pg 85 (2005)
2 clc; clear;
3 e = 1.6e-19; // Electric charge, C
4 V_s = 4.3; // Stopping potential, V
5 K_max = e*V_s; // Maximum kinetic energy attained
    by photoelectrons, J
6 m_e = 9.11e-31 // Mass of electron, kg
7 // Since K.E = eV_s = 0.5m_e(v_max^2), therefore
8 v_max = sqrt((2*K_max)/m_e); // Maximum velocity
    attained by photoelectron, m/s
9 printf("\nMaximum velocity attained by photoelectron
    = %3.1fe+06 m/s", v_max*1e-06);
10
11 // Result
12 // Maximum velocity attained by photoelectron = 1.2e
    +06 m/s
```

---

**Scilab code Exa 3.6** Photoelectric effect for iron

```
1 // Scilab code Ex3.6: Pg 85 (2005)
2 clc; clear;
3
4 // Part (a)
5 I_o = 1; // Total intensity of light, micro-W/cm
    ^2
6 I = (0.030)*(0.040)*I_o; // Intensity available to
    produce photoelectric effect, nW/cm^2
7 printf("\nIntensity available to produce
    photoelectric effect = %3.1f nW/cm^2", I*1e+03);
8
```

```

 9 // Part (b)
10 h = 6.6e-34; // Planck's constant, Js
11 c = 3e+08; // Velocity of light, m/s
12 lamda = 250e-09; // Wavelength, m
13 e_per_sec = (I*lamda*1e-06)/(h*c); // Number of
    electrons emitted per second
14 printf("\nNumber of electrons emitted per second =
    %3.1e", e_per_sec);
15
16 // Part (c)
17 e = 1.6e-019; // Energy equivalent of 1 eV, C
18 i = (e_per_sec)*e; // Electric current in
    phototube, A
19 printf("\nElectric current in phototube = %3.1e A",
    i);
20
21 // Part (d)
22 f_o = 1.1e+15; // Cut-off frequency, Hz
23 phi = (h*f_o)/e; // Work function for iron, eV
24 printf("\nWork function for iron = %3.1f eV", phi);
25
26 // Part (e)
27 V_s = (h*c/(e*lamda))-phi; // Stopping
    voltage, V
28 printf("\nStopping voltage = %4.2f V", V_s);
29
30 // Result
31 // Intensity available to produce photoelectric
    effect = 1.2 nW/cm^2
32 // Number of electrons emitted per second = 1.5e-09
33 // Electric current in phototube = 2.4e-10 A
34 // Work function for iron = 4.5 eV
35 // Stopping voltage = 0.41 V

```

---

Scilab code Exa 3.7 Compton shift for carbon

```

1 // Scilab code Ex3.7: Pg 93 (2005)
2 clc; clear;
3 h = 6.63e-34; // Plank's constant, Js
4 m_e = 9.11e-31; // Mass of electron, kg
5 c = 3e+08; // Velocity of light, m/s
6 theta = ((%pi)/180)*45; // Angle, radians
7 delta_lamda = (h/(m_e*c)*(1-cos(theta))); //
    Compton shift, nm
8 lamda_o = 0.200e-09; // Wavelength of X-ray, nm
9 lamda = delta_lamda+lamda_o // Increased
    wavelength of scattered X-ray, nm
10 printf("\\nIncreased wavelength of scattered X-ray =
    %8.6f nm", lamda*1e+09);
11
12 // Result
13 // Increased wavelength of scattered X-ray =
    0.200711 nm

```

---

### Scilab code Exa 3.8 Xray photons vs visible photons

```

1 // Scilab code Ex3.8: Pg 93 (2005)
2 clc; clear;
3
4 // Part (a)
5 h = 6.63e-34; // Plank's constant, Js
6 q = 1.6e-19; // Electric charge, C
7 m_e = 9.11e-31; // Mass of electron, kg
8 c = 3e+08; // Velocity of light, m/s
9 theta = ((%pi)/180)*90; // Angle, radians
10 delta_lamda = (h/(m_e*c)*(1-cos(theta)));
    // Compton shift, Angstrom
11 lamda_C = 0.0106; // Wavelength of gamma-rays from
    Cobalt,
12 f_d1_C = delta_lamda/ lamda_C; // Fractional
    change in wavelength of gamma rays from cobalt

```

```

13 printf("\nFractional change in wavelength of gamma
    rays from Cobalt = %4.2f", f_dl_C*1e+10);
14 lamda_Mo = 0.712;    // Wavelength of gamma-rays
    from Molybdenum, Angstrom
15 f_dl_Mo = delta_lamda/ lamda_Mo;    //
    Fractional change in wavelength of gamma rays
    from Molybdenum
16 printf("\nFractional change in wavelength of gamma
    rays from Molybdenum = %6.4f", f_dl_Mo*1e+10);
17 lamda_Hg = 5461;    // Wavelength of gamma-rays from
    Mercury, Angstrom
18 f_dl_Hg = delta_lamda/ lamda_Hg;    //
    Fractional change in wavelength of gamma rays
    from mercury
19 printf("\nFractional change in wavelength of gamma
    rays from Mercury = %4.2fe-06", f_dl_Hg*1e+16);
20
21 // Part (b)
22 lamda = 0.712e-10;    // Wavelength of X-rays,
    Angstrom
23 E = (h*c)/(q*lamda);    // Energy of X-rays' photon,
    eV
24 printf("\nEnergy of X-rays photon = %5.0f eV\n", E);
25
26 // Result
27 // Fractional change in wavelength of gamma rays
    from Cobalt = 2.29
28 // Fractional change in wavelength of gamma rays
    from Molybdenum = 0.0341
29 // Fractional change in wavelength of gamma rays
    from Mercury = 4.45fe-06
30 // Energy of X-rays photon = 17460 eV

```

---

Scilab code Exa 3.9 Gravitational redshift for a white dwarf

```
1 // Scilab code Ex3.9: Pg 96 (2005)
2 clc; clear;
3 M = 1.99e+30; // Mass of sun, kg
4 R_s = 6.37e+06; // Radius of earth, m
5 G = 6.67e-11; // Gravitational constant, Nm^2/kg^2
6 lamda = 300e-09; // Wavelength, m
7 c = 3e+08; // Velocity of light, m/s
8 delta_lamda = lamda*((G*M)/(R_s*c^2)); //
   Gravitational redshift, angstrom
9 printf("\\nGravitational redshift = %3.1f angstrom",
   delta_lamda*1e+10);
10
11 // Result
12 // Gravitational redshift = 0.7 angstrom
```

---

# Chapter 4

## The particle nature of matter

Scilab code Exa 4.1 Electrolysis of barium chloride

```
1 // Scilab code Ex4.1: Pg 109 (2005)
2 clc; clear;
3 I = 10; // Electric current, A
4 t = 3600; // Time, s
5 q = I*t; // Electric charge liberated, C
6 mm_Ba = 137; // Molar mass of Barium, g
7 mm_Cl = 35.5; // Molar mass of Chlorine, g
8 valence_Ba = 2; // Valence electrons of Barium
9 valence_Cl = 1; // Valence electrons of
    Chlorine
10 // Using Faradays law of electrolysis, we have
11 m_Ba = (q*mm_Ba)/(96500*valence_Ba); // Mass of
    Barium obtained, g
12 m_Cl = (q*mm_Cl)/(96500*valence_Cl); // Mass of
    Chlorine obtained, g
13 printf("\\nMass of Barium obtained = %4.1f g", m_Ba);
14 printf("\\nMass of Chlorine obtained = %4.1f g", m_Cl
    );
15
16 // Result
17 // Mass of Barium obtained = 25.6 g
```

18 // Mass of Chlorine obtained = 13.2 g

---

#### Scilab code Exa 4.2 Deflection of electron beam by E and B Fields

```
1 // Scilab code Ex4.2: Pg 113 (2005)
2 clc; clear;
3 V = 200; // Electric potential, V
4 theta = 0.20; // Angle, radians
5 l = 0.050; // Length of plates, m
6 d = 1.5e-02; // Distance between two plates, m
7 c_m_r = 1.76e+11; // Charge-to-mass ratio, C/kg
8 // Since  $e/m_e = (V*\theta)/(B^2*l*d)$ , solving for B
9 B = sqrt((V*theta)/(l*d*c_m_r)); // Magnetic
   field, T
10 printf("\nThe magnetic field required to produce the
   deflection of %4.2f rad = %3.1e T", theta, B);
11
12 // Result
13 // The magnetic field required to produce the
   deflection of 0.20 rad = 5.5e-04 T
```

---

#### Scilab code Exa 4.3 Experimental determination of e

```
1 // Scilab code Ex4.3: Pg 117 (2005)
2 clc; clear;
3
4 // Part (a)
5 delta_y = 0.600; // Distance of rise or fall of a
   droplet, cm
6 t_av = 21.0; // Average time of fall of droplet,
   s
7 delta_t = [46.0, 15.5, 28.1, 12.9, 45.3, 20.0]; //
   Rise time of the droplet in succession, s
```



```

8 v = delta_y/t_av; // Average speed of the falling
   droplet , cm/s
9 v_prime = zeros(6);
10 for i = 1:1:6
11     v_prime(i) = delta_y/delta_t(i); // Successive
   speeds of the rising drops , cm/s
12 end
13
14 // Calculate charge ratios
15 q1byq2 = (v+v_prime(1))/(v + v_prime(2));
16 q2byq3 = (v+v_prime(2))/(v + v_prime(3));
17 q3byq4 = (v+v_prime(3))/(v + v_prime(4));
18 q4byq5 = (v+v_prime(4))/(v + v_prime(5));
19 q5byq6 = (v+v_prime(5))/(v + v_prime(6));
20 printf("\nq1/q2 = %5.3 f", q1byq2);
21 printf("\nq2/q3 = %5.3 f", q2byq3);
22 printf("\nq3/q4 = %5.3 f", q3byq4);
23 printf("\nq4/q5 = %5.3 f", q4byq5);
24 printf("\nq5/q6 = %5.3 f", q5byq6);
25 printf("\nThe charge ratios are ratios of small
   whole numbers\n");
26
27 // Part (b)
28 eta = 1.83e-05; // Viscosity of
   air , kg/ms
29 rho = 858; // Oil density , kg
   /m^3
30 g = 9.81; // Acceleration due
   to gravity , m/s^2
31 a = sqrt((9*eta*v*1e-02)/(2*rho*g)); // Radius of
   oil droplet , m
32 V = 4/3*(%pi)*a^3; // Volume of oil
   droplet , m^3
33 m = rho*V; // Mass of oil
   droplet , kg
34 printf("\nRadius of oil droplet = %4.2 e m", a);
35 printf("\nVolume of oil droplet = %4.2 e m^3", V);
36 printf("\nMass of oil droplet = %4.2 e kg", m);

```

```

37
38 // Part (c)
39 V = 4550; // Potential difference across the
    plates of the capacitor, volt
40 d = 0.0160; // Distance between the plates
41 E = V/d; // Electric field between plates, V/m
42 q = zeros(6), e = zeros(6);
43 for i=1:1:6
44     q(i) = m*g/E*((v+v_prime(i))/v); // Charge on
        first drop, C
45     printf("\nq%d = %4.2e V/m", i, q(i));
46 end
47 e(1) = q(1)/5;
48 e(2) = q(2)/8;
49 e(3) = q(3)/6;
50 e(4) = q(4)/9;
51 e(5) = q(5)/5;
52 e(6) = q(6)/7;
53 e_tot = 0;
54 for i = 1:1:6
55     e_tot = e_tot + e(i);
56 end
57 e = e_tot/6;
58 printf("\nThe average charge on an electron = %5.3e
    C", e);
59
60 // Result
61 // q1/q2 = 1.105
62 // q2/q3 = 0.958
63 // q3/q4 = 1.053
64 // q4/q5 = 0.899
65 // q5/q6 = 1.086
66 // The charge ratios are ratios of small whole
    numbers
67
68 // Radius of oil droplet = 1.67e-06 m
69 // Volume of oil droplet = 1.96e-17 m^3
70 // Mass of oil droplet = 1.68e-14 kg

```

```

71
72 // q1 = 8.44e-019 V/m
73 // q2 = 1.36e-018 V/m
74 // q3 = 1.01e-018 V/m
75 // q4 = 1.52e-018 V/m
76 // q5 = 8.48e-019 V/m
77 // q6 = 1.19e-018 V/m
78 // The average charge on an electron = 1.694e-019 C

```

---

#### Scilab code Exa 4.4 Collision of alpha particle with proton

```

1 // Scilab code Ex4.4: Pg 121 (2005)
2 clc; clear;
3
4 // Part (b)
5 // For easy calculations , assume all variables to be
   unity
6 m_p = 1; // Mass of proton , a.m.u
7 m_a = 4*m_p; // Mass of alpha particle , a.m.u
8 Valpha = 1; // Velocity of alpha particle before
   collision , m/s
9 v_p = (2*m_a*Valpha)/(m_a + m_p); // Velocity of
   proton after collision , m/s
10 v_a = ((m_a - m_p)*(Valpha))/(m_a + m_p); //
   Velocity of alph particle after collision , m/s
11 p_change = ((v_a - Valpha)/(Valpha))*100; //
   Percentage change in velocity of alpha particle
12 printf("\\nVelocity of proton after collision = %4.2
   fVa m/s", v_p);
13 printf("\\nVelocity of alpha particle after collision
   = %4.2fVa m/s", v_a);
14 printf("\\nPercentage change in velocity of alpha
   particle = %2d percent", p_change);
15
16 // Result

```

```

17 // Velocity of proton after collision = 1.60 V_a m/s
18 // Velocity of alph particle after collision = 0.60
    V_a m/s
19 // Percentage change in velocity of alpha particle =
    -40 percent

```

---

#### Scilab code Exa 4.5 Radius of Aluminium Nucleus

```

1 // Scilab code Ex4.5: Pg 124 (2005)
2 clc; clear;
3 Z = 13; // Atomic number of Aluminium
4 e = 1.6e-19; // Charge on electron , C
5 k = 8.99e+09; // Coulomb constant , Nm^2/C^2
6 K_a = 7.7e+06*e; // Since K_a = (k*Z_e*2*e)/d_min
    , solving for d_min
7 d_min = (k*2*Z*e^2)/K_a; // Radius of Aluminum, m
8 printf("\\nRadius of Aluminum = %3.1e m", d_min);
9
10 // Result
11 // Radius of Aluminum = 4.9e-15 m

```

---

#### Scilab code Exa 4.7 Collision of alpha particle with proton

```

1 // Scilab code Ex4.7: Pg 135 (2005)
2 clc; clear;
3 // Part (a)
4 n_i = 2; // Initial level of electron
5 n_f = 1; // Final level of electron
6 R = 1.097e+07; // Rydberg constant , per metre
7 c = 3e+08; // Velocity of light , m/s
8 h = 4.136e-15; // Planck's constant , eV
9 lamda = n_i^2*n_f^2/((n_i^2-n_f^2)*R); //
    Wavelength of emitted photon, m

```

```

10 f = c/lamda;      // Frequency of emitted photon, Hz
11 E = h*f;         // Energy of emitted photon, eV
12 printf("\nThe wavelength of emitted photon = %5.1f
    nm", lamda/1e-09);
13 printf("\nThe frequency of emitted photon = %4.2e Hz
    ", f);
14 printf("\nEnergy of emitted photon = %4.1f eV", E);
15
16 // Part (b)
17 mc_square = 938.8e+06; // Energy of recoil of
    hydrogen atom, eV
18 K = 0.5*(E^2/mc_square); // Recoil kinetic
    energy of H atom, eV
19 E_difference = K/E; // Energy difference
20 printf("\nRecoil kinetic energy of H atom = %4.2e eV
    ", K);
21 printf("\nThe fraction of energy difference = %3.1e"
    , E_difference);
22
23 // Result
24 // The wavelength of emitted photon = 121.5 nm
25 // The frequency of emitted photon = 2.47e+15 Hz
26 // Energy of emitted photon = 10.2 eV
27 // Recoil kinetic energy of H atom = 5.55e-08 eV
28 // The fraction of energy difference = 5.4e-09

```

---

#### Scilab code Exa 4.8 series for Hydrngen

```

1 // Scilab code Ex4.8: Pg 136 (2005)
2 clc; clear;
3
4 // Part (a)
5 n_i = 3; // Initial level of electron
6 n_f = 2; // Final level of electron
7 R = 1.097e+07; // Rydberg constant, per metre

```

```

8 c = 3e+08;          // Velocity of light , m/s
9 h = 6.626e-34;     // Plank's constant , Js
10 lamda_max = (n_i^2*n_f^2)/((n_i^2-n_f^2)*R);      //
    Maximum wavelength of emitted photon , m
11 E_photon = (h*c)/(lamda_max*1.6e-19);           //
    Energy of emitted photon , eV
12 printf("\nThe maximum wavelength of emitted photon =
    %5.1f nm", lamda_max/1e-09);
13 printf("\nEnergy of emitted photon = %4.2f eV",
    E_photon);
14
15 // Part (b)
16 n_i = %inf;        // Initial level of electron
17 lamda_min = 1/(R*(1/n_f^2-1/n_i^2));
18 printf("\nThe wavelength corresponding to the series
    limit = %5.1f nm which is in the ultraviolet
    region", lamda_min/1e-09);
19
20 // Result
21 // The maximum wavelength of emitted photon = 656.3
    nm
22 // Energy of emitted photon =1.89 eV
23 //// The wavelength corresponding to the series
    limit = 364.6 nm which is in the ultraviolet
    region

```

---

#### Scilab code Exa 4.9 Hydrogen in its first excited state

```

1 // Scilab code Ex4.9: Pg 137 (2005)
2 clc; clear;
3 k_B = 8.62e-05;    // Boltzmann constant , eV/K
4 delta_E = 10.2;   // Average thermal energy , eV
5 // Since (3/2)*k_B*T = average thermal energy per
    atom = 10.2eV, solving for T
6 T = 10.2/(3/2*k_B); // Temperature at which H-

```

```

    atoms jump to first excited state, K
7  printf("\nThe temperature at which H-atoms jump to
    first excited state = %5d K", T);
8  N_ratio = 0.10;      // Number ratio of population of
    first excited state relative to the ground state
9  // As N_ratio = exp(-delta_E/(k_B*T)), solving for T
10 T = -delta_E/(k_B*log(N_ratio));    // Temperature
    at which H-atoms jump to first excited state, K
11 printf("\nThe temperature of excitation from
    Boltzmann distribution = %5d K", T);
12
13 // Result
14 // The temperature at which H-atoms jump to first
    excited state = 78886 K
15 // The temperature of excitation from Boltzmann
    distribution = 51389 K

```

---

# Chapter 5

## Matter waves

Scilab code Exa 5.1 Wave properties of a baseball

```
1 // Scilab code Ex5.1: Pg 154 (2005)
2 clc; clear;
3 h = 6.63e-34; // Plank's constant, Js
4 m = 140e-03; // Mass of baseball, kg
5 v = 27; // Velocity of baseball, m/s
6 p = m*v; // Momentum of baseball, kgm/s
7 lamda = h/p; // de Broglie wavelength associated
  with baseball, m
8 printf("\\nde-Broglie wavelength associated with
  baseball = %3.1e m", lamda);
9
10 // Result
11 // de-Broglie wavelength associated with baseball =
  1.8e-34 m
```

---

Scilab code Exa 5.2 de Broglie wavelength of an electron

```
1 // Scilab code Ex5.2: Pg 154 (2005)
```



```

2  clc; clear;
3
4  // Part (b)
5  h = 6.63e-34;    // Plank's constant, Js
6  m_e = 9.11e-31; // Mass of electron, kg
7  q = 1.6e-19;    // Charge on electron, C
8  V = 50;        // Electric potential applied, V
9  lamda = h/(sqrt(2*m_e*q*V)); // de Broglie
    wavelength of an electron, m
10 printf("\nde Broglie wavelength of an electron = %3
    .1f angstrom", lamda/1e-10);
11
12 // Result
13 // de Broglie wavelength of an electron = 1.7
    angstrom

```

---

### Scilab code Exa 5.3 Diffraction of neutrons at the crystal lattice

```

1  // Scilab code Ex5.3: Pg 158 (2005)
2  clc; clear;
3  h = 6.63e-34; // Plank's constant, J-s
4  lamda = 1e-10; // de Broglie wavelength of
    neutron, m
5  p = h/lamda; // Momentum associated with neutron
    , kg-m/s
6  m_n = 1.66e-27; // Mass of neutron, kg
7  e = 1.6e-19; // Energy equivalent of 1 eV, J/eV
8  K = p^2/(2*m_n); // Kinetic energy of neutron, eV
9  printf("\nThe momentum of neutrons = %4.2e kg-m/s",
    p)
10 printf("\nThe kinetic energy of neutrons = %4.2 fe -20
    J = %6.4f eV", K*1e+20, K/e);
11
12 // Result
13 // The momentum of neutrons = 6.63e-24 kg-m/s

```

```
14 // The kinetic energy of neutrons = 1.32e-20 J =  
    0.0828 eV
```

---

**Scilab code Exa 5.8** Uncertainty principle for macroscopic objects

```
1 // Scilab code Ex5.8: Pg 177 (2005)  
2 clc; clear;  
3 h_cross = 1.05e-34; // Reduced Plank's constant,  
    J-s  
4 delta_x = 15; // Uncertainty in position, m  
5 v_x = 2; // Velocity of ball, m/s  
6 m = 100e-03; // Mass of ball, kg  
7 delta_p_x = h_cross/(2*delta_x); // Uncertainty  
    in momentum, kg-m/s  
8 delta_v_x = delta_p_x/m; // Minimum spread in  
    velocity, m/s  
9 U_r = delta_v_x/v_x; // Relative uncertainty in  
    velocity of ball  
10 printf("\nThe minimum spread in velocity of ball =  
    %3.1e m/s", delta_v_x);  
11 printf("\nThe relative uncertainty in velocity of  
    ball = %4.2e", U_r);  
12  
13 // Result  
14 // The minimum spread in velocity of ball = 3.5e-35  
    m/s  
15 // The relative uncertainty in velocity of ball =  
    1.75e-35
```

---

**Scilab code Exa 5.9** Kinetic energy of electron confined within the nucleus

```
1 // Scilab code Ex5.9: Pg 178 (2005)  
2 clc; clear;
```

```

3 delta_x = 1.0e-14/2; //
  Uncertainty in position of electron , m
4 q = 1.6e-19; // Charge on
  electron , C
5 h_cross = 1.05e-34; // Reduced Plank '
  s constant , J-s
6 c = 3e+08; // Velocity of
  light , m/s
7 delta_p_x = (h_cross*c)/(2*delta_x*q);
  // Uncertainty in momentum,
  eV/c
8 E_r = 0.551e+06; // Rest
  mass energy if electron , eV
9 E = sqrt((delta_p_x)^2 + (E_r)^2);
10 K = E - E_r; // Kinetic
  energy of electron within nucleus , eV
11 printf("\nKinetic energy of electron within nucleus
  = %4.1f MeV", K/1e+06);
12
13 // Result
14 // Kinetic energy of electron within nucleus = 19.1
  MeV

```

---

### Scilab code Exa 5.10 Width of spectral lines

```

1 // Scilab code Ex5.10: Pg 178 (2005)
2 clc; clear;
3
4 // Part (a)
5 h_cross = 1.05e-34; // Reduced Plank 's constant , J
  -s
6 h = 6.63e-34; // Plank 's constant , J-s
7 delta_t = 1.0e-08; // Average time to measure the
  excited state , s
8 delta_E = h_cross/(2*delta_t); // Uncertainty in

```

```

    energy of the excited state , J
9 // Since  $\Delta E = h \cdot \Delta f$  , solving for  $\Delta f$ 
10  $\Delta f = \Delta E / h$ ; // Line width of emitted
    light , Hz
11 printf(" \nLine width of emitted light = %2.0e Hz" ,
     $\Delta f$ );
12
13 // Part (b)
14  $c = 3e+08$ ; // Velocity of light , m/s
15  $\lambda = 500e-09$ ; // Wavelength of spectral line
    , m
16  $f_o = c / \lambda$ ; // Center frequency of spectral
    line , Hz
17  $f_b = \Delta f / f_o$ ; // Fractional broadening of
    spectral line
18 printf(" \nFractional broadening of spectral line =
    %3.1e" ,  $f_b$ );
19
20 // Result
21 // Line width of emitted light =  $8.0e+06$  Hz
22 // Fractional broadening of spectral line =  $1.3e-08$ 

```

---

# Chapter 6

## Quantum mechanics in one dimension

Scilab code Exa 6.2 Probability from wave function

```
1 // Scilab code Ex6.2: Pg 193 (2005)
2 clc; clear;
3 x0 = 1; // For simplicity assume x0 = 1
4 C = 1/sqrt(x0); // Normalization constant
5 P = 2*C^2*integrate('exp(-2*x/x0)', 'x', 0, x0);
6 printf("\nThe probability that the particle will be
    found in the interval  $-x_0 \leq x \leq x_0$  is %6.4f or
    %4.1f percent", P, P*100);
7
8 // Result
9 // The probability that the particle will be found
    in the interval  $-x_0 \leq x \leq x_0$  is 0.8647
```

---

Scilab code Exa 6.4 Dispersion of matter waves

```
1 // Scilab code Ex6.4: Pg 197 (2005)
```

```

2 clc; clear;
3 delta_x0 = 1e-010;    // Initial width of the
   localized space, m
4 delta_xt = 10*delta_x0;    // Final width at which
   the wave packet is dispersed, m
5 h_cross = 1.055e-034;    // Reduced Planck's
   constant, Js
6 m = 9.11e-031;    // Mass of the electron, kg
7 // From Dispersion relation,  $\Delta x_t^2 - \Delta x_0^2 = \sqrt{h_{cross} * t / (2 * m * \Delta x_0^2)}$ , solving for t
8 t = 2*m*sqrt(delta_xt^2 - delta_x0^2)*delta_x0/
   h_cross;    // Time which elapses before
   delocalization
9 printf("\nThe time which elapses before the
   localization of electron destroys = %3.1e s", t);
10 m = 1e-03;    // Mass of marble, kg
11 delta_x0 = 1e-004;    // Initial width of the
   localized space, m
12 delta_xt = 10*delta_x0;    // Final width at which
   the wave packet is dispersed, m
13 t = 2*m*sqrt(delta_xt^2 - delta_x0^2)*delta_x0/
   h_cross;    // Time which elapses before
   delocalization
14 printf("\nThe time which elapses before the
   localization of marble destroys = %3.1e s", t);
15 printf("\nFor all the practical purposes, the marble
   will remain localized for ever");
16 // Result
17 //

```

---

### Scilab code Exa 6.5 Energy Quantization for Macroscopic Object

```

1 // Scilab code Ex6.5: Pg 202 (2005)
2 clc; clear;
3 h = 6.626e-034;    // Planck's constant, Js

```

```

4 m = 1e-06; // Mass of the object , kg
5 n = 1; // Quantum number for minimum energy level
6 L = 1e-02; // Distance between two rigid walls ,
    m
7 E1 = n^2*h^2/(8*m*L^2); // Minimum energy of the
    object , J
8 v1 = sqrt(2*E1/m); // Minimum speed of the object
    , m/s
9 v = 3.00e-02; // Given speed of the objct , m/s
10 E = 1/2*m*v^2; // Energy of the object for given
    speed , J
11 n = sqrt(8*m*L^2*E)/h; // Quantum number
    corresponding to the given speed
12 printf("\nThe minimum speed of the object = %4.2e m/
    s", v1);
13 printf("\nThe quantum number corresponding to the
    speed of %4.2e m/s is n = %4.2e", v1, n);
14
15 // Result
16 // The minimum speed of the object = 3.31e-26 m/s
17 // The quantum number corresponding to the speed of
    3.31e-26 m/s is n = 9.06e+23

```

---

### Scilab code Exa 6.6 Model of an Atom

```

1 // Scilab code Ex6.6: Pg 203 (2005)
2 clc; clear;
3 c = 1; // Assume speed of light to be unity , m/
    s
4 h_cross = 197.3; // Reduced Planck's constant , eV
    .nm/c^2
5 m_e = 511e+03; // Mass of an electron , eV/c
    ^2
6 L = 0.200; // Length of the box , nm
7 E1 = %pi^2*(h_cross/c)^2/(2*m_e*L^2); // Ground

```

```

state energy of atomic electron , eV
8 E2 = 2^2*E1; // Excited state energy of the
atomic electron , eV
9 delta_E = E2- E1; // Energy that must be applied
to the electron to raise it from ground to the
first excited state , eV
10 h = 2*pi*h_cross; // Planck's constant , Js
11 lambda = h*c/delta_E; // Wavelength of the photon
to cause the electron transition , nm
12 printf("\nThe energy that must be applied to the
electron to raise it from ground to the first
excited state = %4.1f eV", delta_E);
13 printf("\nThe wavelength of the photon to cause this
electron transition = %4.1f nm", lambda);
14 printf("\nThis wavelength is in the far ultraviolet
region.");
15
16 // Result
17 // The energy that must be applied to the electron
to raise it from ground to the first excited
state = 28.2 eV
18 // The wavelength of the photon to cause this
electron transition = 44.0 nm
19 // This wavelength is in the far ultraviolet region.

```

---

#### Scilab code Exa 6.7 Probabilities for a particle in a Box

```

1 // Scilab code Ex6.7: Pg 205 (2005)
2 clc; clear;
3 L = 1; // For simplicity assume length of finite
square well to be unity , m
4 P = 2/L*integrate('sin(%pi*x/L)^2', 'x', L/4, 3*L/4)
; // Probability that the particle will be found
in the middle half of the well
5 printf("\nThe probability that the particle will be

```



```

        found in the middle half of the well = %5.3f", P)
    ;
6
7 // Result
8 // The probability that the particle will be found
    in the middle half of the well = 0.818

```

---

**Scilab code Exa 6.8** Ground state energy of an electron confined to a potential well

```

1
2 // Scilab code Ex6.8: Pg 211 (2005)
3 clc; clear;
4 c = 1; // Assume speed of light to be unity, m/
    s
5 L = 0.200; // Width of the potential well, nm
6 h_cross = 197.3; // Reduced Planck's constant, eV
    .nm/c^2
7 m = 511e+03; // Mass of an electron, eV/c^2
8 U = 100; // Height of potential well, eV
9 delta = h_cross/sqrt(2*m*U); // Decay length of
    electron, nm
10 L = L + 2*delta; // Effective length of the
    infinite potential well, nm
11 E = %pi^2*(h_cross/c)^2/(2*m*L^2); // Ground state
    energy of the electron with effective length, eV
12 U = U - E; // New potential energy, eV
13 delta = h_cross/sqrt(2*m*U); // New decay length
    of electron, nm
14 printf("\\nThe ground state energy of an electron
    confined to the potential well = %4.2f eV", E);
15 printf("\\nThe new decay length of the electron = %6
    .4f nm", delta);
16
17 // Result
18 // The ground state energy of an electron confined

```

```
to the potential well = 6.58 eV
19 // The new decay length of the electron = 0.0202 nm
```

---

**Scilab code Exa 6.12** The quantum oscillator in nonclassical region

```
1 // Scilab code Ex6.12: Pg 214 (2005)
2 clc; clear;
3 P = 2/sqrt(%pi)*integrate('exp(-z^2)', 'z', 1, 100);
   // Probability that the quantum oscillator in
   its ground state will be found in the
   nonclassical region
4 printf("\nThe probability that the quantum
   oscillator in its ground state will be found in
   the nonclassical region = %5.3f", P);
5
6 // Result
7 // The probability that the quantum oscillator in
   its ground state will be found in the
   nonclassical region = 0.157
```

---

**Scilab code Exa 6.13** Quantization of vibrational energy

```
1 // Scilab code Ex6.13: Pg 211 (2005)
2 clc; clear;
3 h_cross = 6.582e-016; // Reduced Planck's constant
   , eV-s
4 // For spring-mass system
5 K = 0.100; // Force constant of the spring-mass
   system, N/m
6 m = 0.0100; // Mass attached to the spring, kg
7 omega = sqrt(K/m); // Angular frequency of
   oscillations, rad/s
```

```

8 delta_E = h_cross*omega;    // Energy spacing
   between quantum levels , eV
9 printf("\nThe energy spacing between quantum levels
   for spring-mass system = %4.2e eV\nwhich is far
   below present limits of detection", delta_E);
10 // For vibrating hydrogen molecule
11 K = 510.5; // Force constant of the hydrogen
   molecule system , N/m
12 mu = 8.37e-028; // Reduced mass of the hydrogen
   molecule , kg
13 omega = sqrt(K/mu); // Angular frequency of
   oscillations , rad/s
14 delta_E = h_cross*omega;    // Energy spacing
   between quantum levels , eV
15 printf("\nThe energy spacing between quantum levels
   for hydrogen molecule = %5.3f eV\nwhich can be
   measured easily", delta_E);
16
17 // Result
18 // The energy spacing between quantum levels for
   spring-mass system = 2.08e-15 eV
19 // which is far below present limits of detection
20 // The energy spacing between quantum levels for
   hydrogen molecule = 0.514 eV
21 // which can be measured easily

```

---

#### Scilab code Exa 6.14 Standard Deviations from Averages

```

1 // Scilab code Ex6.14: Pg 219 (2005)
2 clc; clear;
3 x = [2.5, 3.7, 1.4, 7.9, 6.2, 5.4, 8.0, 6.4, 4.1,
   5.4, 7.0, 3.3, 4.2, 8.8, 6.2, 7.1, 5.4, 5.3]; //
   Data entries
4 sum_x = 0; // Initialize the accumulator
5 sum_x_sq = 0; // Initialize the second accumulator

```

```

6 N = 18;          // Total number of data points
7 for i = 1:1:N
8     sum_x = sum_x + x(i);    // Sum of data
9     sum_x_sq = sum_x_sq + x(i)^2; // Sum of square
        of data
10 end
11 x_av = sum_x/N;          // Average of data
12 x_sq_av = sum_x_sq/N;    // Mean square value
13 sigma = sqrt(x_sq_av-x_av^2); // Standard
        deviation from averages
14 printf("\nThe standard deviation from averages = %4
        .2f", sigma);
15
16
17 // Result
18 // The standard deviation from averages = 1.93

```

---

#### Scilab code Exa 6.15 Location of a particle in the box

```

1 // Scilab code Ex6.15: Pg 219 (2005)
2 clc; clear;
3 L = 1; // For simplicity assume length of the box
        to be unity, unit
4 x_av = 2*L/%pi^2*integrate('theta*sin(theta)^2', '
        theta', 0, %pi); // Average value of x
5 x_sq_av = L^2/%pi^3*(integrate('theta^2', 'theta',
        0, %pi)-integrate('theta^2*cos(2*theta)', 'theta',
        , 0, %pi)); // Average value of x square
6 delta_x = sqrt(x_sq_av - x_av^2); // Uncertainty
        in the position for this particle, unit
7 printf("\nThe average position of the particle in
        the box = L/%1d", x_av*4);
8 printf("\nThe uncertainty in the position for the
        particle = %5.3fL", delta_x);
9

```

```
10 // Result
11 // The average position of the particle in the box =
    L/2
12 // The uncertainty in the position for the particle
    = 0.181L
```

---

# Chapter 7

## Tunnelling phenomena

Scilab code Exa 7.1 Transmission coefficient for an oxide layer

```
1 // Scilab code Ex7.1: Pg 235 (2005)
2 clc; clear;
3 c = 3e+08; // Velocity of light, m/s
4 m_e = 511e+03/(c^2); // Mass of electron, eV
5 U = 3.00; // Ground state energy neglecting E, eV
6 h_cross = (1.973e+03)/c; // Reduced planck's
   constant, eV
7 alpha = sqrt(2*m_e*U)/h_cross;
8 L = 50; // Thickness of the layer, angstrom
9 T = 1/(1+1/4*10^2/(7*3)*sinh(alpha*L)^2);
10 printf("\n\nThe transmission coefficient for the layer
   thickness of");
11 printf("\n\n%2d angstrom = %5.3e", L, T);
12 L = 10; // // Thickness of the layer, angstrom
13 T = 1/(1+1/4*10^2/(7*3)*sinh(alpha*L)^2);
14 printf("\n\n%2d angstrom = %5.3e", L, T);
15
16 // Result
17 // The transmission coefficient for the layer
   thickness of
18 // 50 angstrom = 9.628e-39
```

19 // 10 angstrom = 6.573e-08

---

### Scilab code Exa 7.2 Tunnelling current through an oxide layer

```
1 // Scilab code Ex7.2: Pg 236 (2005)
2 clc; clear;
3 e = 1.60e-19; // Electric charge, C
4 i = 1.00e-03; // Electron current, A
5 N = i/e; // Electrons per second
6 T = 0.657e-07; // Fraction of electrons
   transmitted
7 T_e = N*T; // Number of electrons transmitted per
   second
8 T_i = T_e*e; // Transmitted current, A
9 printf("\nThe transmitted current through the oxide
   layer = %4.1f pA", T_i*1e+12);
10
11 // Result
12 // The transmitted current through the oxide layer =
   65.7 pA
```

---

### Scilab code Exa 7.5 Tunnelling in a parallel plate capacitor

```
1 // Scilab code Ex7.5: Pg 241 (2005)
2 clc; clear;
3 epsilon_c = 5.5e+10; // Characteristic field
   strength, V/m
4 epsilon = 1.0e+09; // Electric field, V/m
5 f = 1.0e+30; // Collision frequency, s(-1)cm(-2)
6 lamda = f*exp(-epsilon_c/epsilon); // Electron
   emission rate, electrons/sec
7 e = 1.60e-19; // Electric charge, C
8 I = lamda*e; // Tunelling current, A
```

```

9 printf("\nTunelling current in parallel plate
   capacitor = %4.2f pA", I/1e-12);
10 printf("\n");
11
12 // Result
13 // Tunelling current in parallel plate capacitor =
   0.21 pA

```

---

### Scilab code Exa 7.6 Estimating halflives of Thorium and Polonium

```

1 // Scilab code Ex7.6: Pg 244 (2005)
2 clc; clear;
3 Z_T = 88; // Atomic number of daughter nucleus
4 E_T = 4.05e+06; // Energy of ejected alphas, eV
5 R = 9.00e-15; // Nuclear radius, m
6 r_o = 7.25e-15; // Bohr radius, m
7 E_o = 0.0993e+06; // Energy analogous to the
   Rydberg in Atomic Physics
8 T_T = exp(-4*%pi*Z_T*sqrt(E_o/E_T) + 8*sqrt((Z_T*R)/
   r_o)); // Transmission factor in case of
   Thorium
9 f = 1e+21; // Frequency of collisions, Hz
10 lamda_T = f*T_T; // Decay rate in case of Thorium
   , s(-1)
11 t_T = 0.693/lamda_T; // Half-life time of
   Thorium, s
12 Z_P = 82; // Atomic number of daughter nucleus
13 E_P = 8.95e+06; // Energy of ejected alphas, eV
14 R = 9.00e-15; // Nuclear radius, m
15 r_o = 7.25e-15; // Bohr radius, m
16 E_o = 0.0993e+06; // Energy unit, eV
17 T_P = exp(-4*%pi*Z_P*sqrt(E_o/E_P) + 8*sqrt((Z_P*R)/
   r_o)); // Transmission factor in case of
   Polonium
18 f = 1e+21; // Frequency of collisions, Hz

```



```

19 lamda_P = f*T_P;          // Decay rate in case of
    Thorium, s(-1)
20 t_P = 0.693/lamda_P;     // Half-life time of
    Polonium, s
21
22 printf("\nHalf-life time of Thorium = %3.1e s = %3.1
    e yrs", t_T, t_T/(365*24*60*60));
23 printf("\nHalf-life time of Polonium = %3.1e s", t_P
    );
24
25 // Result
26 // Half-life time of Thorium = 5.3e+17 s = 1.7e+10
    yrs
27 // Half-life time of Polonium = 8.4e-10 s

```

---

## Chapter 8

# Quantum Mechanics in Three Dimensions

Scilab code Exa 8.4 Orbital quantum number for a stone

```
1 // Scilab code Ex8.4: Pg 270 (2005)
2 clc; clear;
3 R = 1.00; // Radius of circle , m
4 T = 1.00; // Time period of revolution , s
5 v = (2*%pi*R)/T; // Speed of stone in its orbit ,
   m/s
6 m = 1.00; // Mass of stone , kg
7 L = m*v*R; // Angular momentum of stone , kg-m^2/s
8 h_cross = 1.055e-34; // Reduced Planck 's constant
   , kg-m^2/s
9 l = L/h_cross; // Orbital quantum number
10 printf("\\nThe orbital quantum number for stone = %4.2
   fe+34" , l*1e-34);
11
12 // Result
13 // Orbital quantum number for stone = 5.96e+34
```

---

### Scilab code Exa 8.6 Space quantisation for an atomic electron

```
1 // Scilab code Ex8.6: Pg 272 (2005)
2 clc; clear;
3 // For simplicity let h_cross = 1
4 h_cross = 1; // Reduced planck's constant
5 l = 3; // Given orbital quantum number
6 L = sqrt(l*(l+1)*h_cross); // Magnitude of total
    angular momentum, in h_cross units
7 m_l = [-3, -2, -1, 0, 1, 2, 3];
8 L_z = m_l*h_cross; // Allowed values of L_z
9 cos_theta = L_z/L;
10 theta = acosd(L_z/L); // Orientations of L_z,
    degrees
11 for i = 1:1:7
12     if theta(i) > 90 then
13         theta(i) = theta(i)-180;
14     end
15 end
16 printf("\nThe magnitude of total angular momentum =
    2*sqrt(%d)*h_cross\n", L^2/4);
17 printf("\nThe allowed values of L_z in units of
    h_cross are :");
18 disp(L_z);
19 printf("\nThe orientations of L_z in degrees are:");
20 disp(theta);
21
22 // Result
23 // The magnitude of total angular momentum = 2*sqrt
    (2)*h_cross
24
25 // The allowed values of L_z in units of h_cross are
    :
26 //   - 3.   - 2.   - 1.    0.    1.    2.    3.
27
28 // The orientations of L_z in degrees are:
29 //   - 30.   - 54.73561  - 73.221345   90.
    73.221345   54.73561   30.
```

---

Scilab code Exa 8.7 Energy of Hydrogen atom at first excited state

```
1 // Scilab code Ex8.7: Pg 281 (2005)
2 clc; clear;
3 k = 9e+09; // Coulomb constant, N/Sq.m/C
4 e = 1.6e-019; // Electronic charge, C
5 a_0 = 0.529e-010; // Bohr's radius, m
6 n = 2; // Principal quantum number
7 l = [0, 1]; // Orbital quantum number
8 m_l = [-1, 0, 1]; // Orbital magnetic quantum
   number
9 Z = 1; // Atomic number of hydrogen
10 E2 = -k*e^2/(2*a_0)*Z^2/n^2; // Energy of first
   excited level of hydrogen,
11 printf("\nThe energy of first excited level of
   hydrogen = %3.1f eV", E2/e);
12
13 // Result
14 // The energy of first excited level of hydrogen =
   -3.4 eV
```

---

Scilab code Exa 8.8 Probabilities for the Electron in Hydrogen

```
1 // Scilab code Ex8.8: Pg 284 (2005)
2 clc; clear;
3 P = 1/2*integrate('z^2*exp(-z)', 'z', 2, 100); //
   Take some large value of upper limit
4 printf("\nP(electron in the ground state of hydrogen
   will be found outside the first Bohr radius) =
   %4.1f percent", P*100);
5
```

```
6 // Result
7 // P(electron in the ground state of hydrogen will
  be found outside the first Bohr radius) = 67.7
  percent
```

---

# Chapter 9

## Atomic Structure

Scilab code Exa 9.1 Magnetic energy of electron in Hydrogen

```
1 // Scilab code Ex9.1: Pg 300 (2005)
2 clc; clear;
3 // Since  $\mu_B = (e \cdot h_{\text{cross}}) / (2 \cdot m_e)$ 
4  $\mu_B = 9.27e-24$ ; // Bohr magneton, J/T
5  $B = 1.00$ ; // Magnetic flux, T
6 // Since  $1 \text{ eV} = 1.6e-19 \text{ J}$ 
7  $eV = 1.6e-19$ ; // Energy, J
8  $h_{\text{cross}} = 6.58e-16$ ; // Reduced Plank's constant,
   eV-s
9  $\omega_L = (\mu_B \cdot B) / (eV \cdot h_{\text{cross}})$ ; // Larmor
   frequency, rad/s
10 printf(" \nLarmour frequency at n = 2 is %4.2 fe+10
   rad/s",  $\omega_L \cdot 1e-10$ );
11
12 // Result
13 // Larmour frequency at n = 2 is  $8.81e+10 \text{ rad/s}$ 
```

---

Scilab code Exa 9.2 Angles between z axis and the spin angular momentum vector

```

1 // Scilab code Ex9.2: Pg 307 (2005)
2 clc; clear;
3 h_cross = 6.58e-16; // Reduced Plank's constant,
   eV-s
4 S = h_cross*sqrt(3)/2; // Spin angular momentum,
   eV-s
5 S_z = h_cross/2; // Z-component of spin angular
   momentum, eV-s
6 theta_up = acosd(S_z/S);
7 theta_down = acosd(-S_z/S);
8 printf("\nFor up spin state, theta = %4.2f degrees",
   theta_up);
9 printf("\nFor down spin state, theta = %5.1f degrees
   ", theta_down);
10
11 // Result
12 // For up spin state, theta = 54.74 degrees
13 // For down spin state, theta = 125.3 degrees

```

---

### Scilab code Exa 9.3 Zeeman Spectrum of Hydrogen Including Spin

```

1 // Scilab code Ex9.3: Pg 311 (2005)
2 clc; clear;
3 e = 1.6e-019; // Energy equivalent of 1 eV, J/eV
4 B = 1.00; // Magnitude of magnetic field, tesla
5 n = 2; // Initial state of the hydrogen atom
6 mu_B = 9.27e-024; // Bohr's magneton, J/T
7 E_Z = mu_B*B/e; // Zeeman energy, eV
8 E2 = -13.6/n^2; // Energy of first excited state
   , eV
9 m_l = [-2, -1, 0, 1, 2]; // Orbital magnetic
   quantum number for l = 2
10 printf("\nThe energies of the electron (in eV) in n
   = 2 state are:\n");
11 for i = 1:1:5

```

```

12     if m_l(i) < 0 then
13         sig = '-';
14     else
15         sig = '+';
16     end
17     printf(" (%4.2f %s %4.2e) ", E2, sig, abs(E_Z*
        m_l(i)));
18 end
19
20 // Result
21 // The energies of the electron (in eV) in n = 2
    state are:
22 // (-3.40 - 1.16e-04) (-3.40 - 5.79e-05) (-3.40 +
    0.00e+00) (-3.40 + 5.79e-05) (-3.40 + 1.16e-04

```

---

#### Scilab code Exa 9.4 Spin orbit energy of Sodium doublet

```

1 // Scilab code Ex9.4: Pg 311 (2005)
2 clc; clear;
3 hc = 1240; // Product of plank's constant &
    velocity of light, eV
4 lamda_1 = 588.995; // Wavelength of first doublet
    of Na lines, nm
5 lamda_2 = 589.592; // Wavelength of second
    doublet of Na lines, nm
6 delta_E = hc*(lamda_2 - lamda_1)/(lamda_1*lamda_2);
    // Spin orbit energy, eV
7 printf("\nSpin orbit energy from doublet spacing =
    %4.2fe-03 eV", delta_E*1e+03);
8
9 // Result
10 // Spin orbit energy from doublet spacing = 2.13e-03
    eV

```

---



### Scilab code Exa 9.5 Ground state of Helium atom

```
1 // Scilab code Ex9.5: Pg 316 (2005)
2 clc; clear;
3 n = 1; // Principal quantum number
4 Z = 2; // Atomic number of Helium
5 E_a = (-13.6*Z^2)/n^2; // Energy of the
   electron in state 'a', eV
6 E_b = (-13.6*Z^2)/n^2; // Energy of the
   electron in state 'b', eV
7 E = E_a + E_b; // Total electronic energy of
   Helium, eV
8 printf("\nTotal electronic energy of Helium = %5.1f
   eV", E);
9
10 // Result
11 // Total electronic energy of Helium = -108.8 eV
```

---

### Scilab code Exa 9.6 Effective atomic number for 3s electron in Na

```
1 // Scilab code Ex9.6: Pg 317 (2005)
2 clc; clear;
3 E_i = 5.14; // Ionisation energy of Na, eV
4 n = 3; // Principal quantum number
5 Z_eff = sqrt((n^2*E_i)/13.6); // Effective atmic
   number
6 printf("\nEffective atomic number for 3s electron in
   Na = %4.2f", Z_eff);
7
8 // Result
9 // Effective atomic number for 3s electron in Na =
   1.84
```



# Chapter 10

## Statistical Physics

Scilab code Exa 10.1 Population of excited states with respect to ground states in

```
1 // Scilab code Ex10.1: Pg 340 (2005)
2 clc; clear;
3 // Part (a)
4 E1 = -13.6; // Energy of ground state, eV
5 E2 = -3.40; // Energy of first excited state, eV
6 E3 = -1.51; // Energy of second excited state, eV
7 g1 = 2; // Degeneracy for ground state
8 g2 = 8; // Degeneracy for first excited state
9 g3 = 18; // Degeneracy for second excited state
10 kB = 8.617e-05; // Boltzmann constant, eV/K
11 Ta = 300; // Temperature, K
12 // As  $n_2/n_1 = (g_2 * A * e^{(-E_2/(k_B * T))}) / (g_1 * A * e^{(-E_1/(k_B * T))})$ , on simplifying we get
13 N21 = (g2/g1)*exp((E1 - E2)/(kB*Ta)); // The
// population of first excited state w.r.t ground
// state
14 printf("\nThe population of first excited state w.r.
// t. ground state at %3d K = %1d", Ta, N21);
15
16 // Part (b)
17 Tb = 20000; // Temperature, K
```

```

18 n21 = (g2/g1)*exp((E1 - E2)/(kB*Tb)); // The
    population of first excited state w.r.t ground
    state
19 n31 = (g3/g1)*exp((E1 - E3)/(kB*Tb)); // The
    population of second excited state w.r.t ground
    state
20 printf("\nThe population of first excited state w.r.
    t. ground state at %4d K = %6.4f", Tb, n21);
21 printf("\nThe population of second excited state w.r
    .t ground state at %4d K = %6.4f", Tb, n31);
22
23 // Part (c)
24 E_strength = (g3/g2)*exp((E2 - E3)/(kB*Tb)); //
    Emission strength
25 printf("\nEmission strength of spectral lines = %3.2
    f", E_strength);
26
27 // Result
28 // The population of first excited state w.r.t.
    ground state at 300 K = 0
29 // The population of first excited state w.r.t.
    ground state at 20000 K = 0.0108
30 // The population of second excited state w.r.t
    ground state at 20000 K = 0.0081
31 // Emission strength of spectral lines = 0.75

```

---

### Scilab code Exa 10.2 Validity of Maxwell Boltzmann Statistics

```

1 // Scilab code Ex10.2: Pg 345 (2005)
2 clc; clear;
3 // Part (a)
4 N = 6.02e+23; // Number of molecules at STP
5 m = 3.34e-27; // Mass of H-molecule , kg
6 h_cross = 1.055e-34; // Reduced Plank's constant ,
    J-s

```

```

7 V = 22.4e-03;      // Volume occupied by molecules at
  STP, m^3
8 T = 273;          // Absolute temperature, K
9 k_B = 13.8e-24;   // Boltzmann constant, J/K
10 x_H = N/V*h_cross^3/(8*(m*k_B*T)^(3/2)); //
  Particle concentration at STP
11 printf("\nx_H = %4.2e", x_H);
12 if (x_H < 1)
13 printf("\nThe criterion for the validity of
  Maxwell Boltzmann Statistics is satisfied in
  hydrogen.");
14
15 // Part (b)
16 d_Ag = 10.5;      // Density of silver, g/m^3
17 M_Ag = 107.9;     // Molar weight of silver, g
18 NV_Ag = (d_Ag/M_Ag)*(6.02e+023)*1e+06; // Density
  of free electrons in silver, electrons/m^3
19 me = 9.109e-031; // Mass of an electron, kg
20 T = 300;         // Room temperature, K
21 x_Ag = ((NV_Ag)*h_cross^3)/(8*(me*k_B*T)^(3/2));
  // Particle concentration at STP
22 printf("\nx_Ag = %4.2f", x_Ag);
23 if (x_Ag > 1)
24 printf("\nThe criterion for the validity of
  Maxwell Boltzmann Statistics does not hold for
  electrons in silver");
25
26 // Result
27 // x_H = 8.84e-08
28 // The criterion for the validity of
  Maxwell Boltzmann Statistics is satisfied in
  hydrogen.
29 // x_Ag = 37.13
30 // The criterion for the validity of
  Maxwell Boltzmann Statistics does not hold for
  electrons in silver

```

---

### Scilab code Exa 10.3 Photons in a box

```
1 // Scilab code Ex10.3: Pg 352 (2005)
2 clc; clear;
3 // Part (b)
4 I = integrate('z^2/(exp(z)-1)', 'z', 0, 100); //
    Integral value
5 k_B = 8.62e-05; // Boltzmann constant, eV/K
6 T = 3000; // Temperature, K
7 h = 4.136e-15; // Plank's constant, eV
8 c = 3e+10; // Velocity of light, cm/s
9 N_V = 8*pi*((k_B*T)/(h*c))^3*I; // Number of
    photons/cc
10 printf("\nThe density of photons inside the cavity =
    %4.2fe+11 photons/cc", N_V*1e-11);
11
12 // Result
13 // The density of photons inside the cavity = 5.47e
    +11 photons/cc
```

---

### Scilab code Exa 10.4 Specific Heat of Diamond

```
1 // Scilab code Ex10.4: Pg 356 (2005)
2 clc; clear;
3
4 // Part (a)
5 k_B = 8.62e-05; // Boltzmann constant, eV/K
6 T_E = 1300; // Temperature, K
7 h_cross = 6.58e-16; // Reduced plank's constant,
    eV-s
8 omega = (k_B*T_E)/h_cross; // Frequency of
    vibration of carbon atom in diamond, Hz
```

```

9 spacing = (h_cross*omega); // Spacing between
    adjacent oscillator energy level, eV
10 printf("\nFrequency of vibration of carbon atom in
    diamond = %4.2e Hz", omega);
11 printf("\nSpacing between adjacent oscillator energy
    level = %5.3f eV", spacing);
12
13 // Part (b)
14 T_R = 300; // Room temperature, K
15 p = exp((h_cross*omega)/(k_B*T_R)); // For
    simplification
16 E_R = (h_cross*omega)/(p-1); // Average energy of
    oscillator at room temperature, eV
17 T = 1500; // Temperature, K
18 q = exp((h_cross*omega)/(k_B*T)); // For
    simplification
19 E_bar = (h_cross*omega)/(q-1); // Average energy
    at 1500 K, eV
20 printf("\nAverage energy of oscillator at room
    temperature = %7.5f eV", E_R);
21 printf("\nAverage oscillator energy at %4d K = %7.5f
    eV", T, E_bar);
22
23
24 // Result
25 // Frequency of vibration of carbon atom in diamond
    = 1.70e+14 Hz
26 // Spacing between adjacent oscillator energy level
    = 0.112 eV
27 // Average energy of oscillator at room temperature
    = 0.00149 eV
28 // Average oscillator energy at 1500 K = 0.0813 eV

```

---

Scilab code Exa 10.5 Fermi Energy of Gold

```

1 // Scilab code Ex10.5: Pg 360 (2005)
2 clc; clear;
3
4 // Part (a)
5 h = 6.625e-34; // Plank's constant, J-s
6 m_e = 9.11e-31; // Mass of electron, kg
7 density = 19.32/(1e-02)^3; // Density of gold, g/m
  ^3
8 weight = 197; // Molar weight, g/mol
9 N_V = (density/weight)*6.02e+23; // Number of
  electrons per mole
10 E_F = (h^2/(2*m_e*1.6e-19))*((3*(N_V))/(8*pi))
  ^((2/3)); // Fermi energy of Gold at 0 K
11 printf("\nFermi energy of Gold at 0 K = %4.2f eV",
  E_F);
12
13 // Part (b)
14 v_F = sqrt((2*E_F*1.6e-19)/m_e); // Fermi speed of
  Gold at 0 K
15 printf("\nFermi speed of Gold at 0 K = %4.2fe+06 m/s
  ", v_F*1e-06);
16
17 // Part (c)
18 k_B = 8.62e-05; // Boltzmann constant, eV/K
19 T_F = (E_F)/(k_B); // Fermi temperature for Gold
  at 0 K, K
20 printf("\nFermi temperature for Gold at 0 K = %5d K"
  , T_F);
21
22 // Result
23 // Fermi energy of Gold at 0 K = 5.53 eV
24 // Fermi speed of Gold at 0 K = 1.39fe+06 m/s
25 // Fermi temperature for Gold at 0 K = 64201 K

```

---



# Chapter 11

## Molecular Structure

Scilab code Exa 11.1 Rotation of CO molecule

```
1 // Scilab code Ex11.1: Pg 380 (2005)
2 clc; clear;
3 // Part (a)
4 f = 1.15e+11; // Frequency of transitions, Hz
5 omega = 2*(%pi)*f; // Angular frequency of
   absorbed radiations, Hz
6 h_cross = 1.055e-34; // Reduced planks constant, J
   -s
7 // Since  $E = (h\_cross)^2/I\_CM = h\_cross*omega$ ,
   solving for I_CM
8 I_CM = h_cross/omega; // Moment of inertia of
   molecule about its center of mass, kg-m^2
9 printf("\nThe moment of inertia of molecule about
   its center of mass = %4.2e kg-m^2", I_CM);
10
11 // Part (b)
12 m_O = 16; // Mass of oxygen atom, a.m.u
13 m_C = 12; // Mass of carbon atom, a.m.u
14 mu = ( m_O * m_C * 0.166e-26)/(m_O + m_C); //
   Reduced mass, kg
15 // Since  $I\_CM = mew*R\_o^2$ , solving for R_o
```

```

16 R_0 = sqrt(I_CM/mu);    // Bond length of carbon
    monoxide molecule , m
17 printf("\nThe bond length of carbon monoxide
    molecule = %5.3f nm", R_0/1e-09);
18
19 // Result
20 // The moment of inertia of molecule about its
    center of mass = 1.46e-046 kg-m^2
21 // The bond length of carbon monoxide molecule =
    0.113 nm

```

---

#### Scilab code Exa 11.2 Variation of CO molecule

```

1 // Scilab code Ex11.2: Pg 383 (2005)
2 clc; clear;
3
4 // Part (a)
5 f = 6.42e+13;    // Frequency of absorption , Hz
6 omega = 2*(%pi)*f;    // Angular frequency of
    absorbed radiations , Hz
7 mu = 1.14e-26;    // Reduced mass of CO molecule , kg
8 K = mu*(omega^2);    // Effective force constant of
    CO molecule , N/m
9 printf("\nThe effective force constant of CO
    molecule = %4.2e N/m", K);
10
11 // Part (b)
12 h_cross = 1.055e-34;    // Reduced Planck's constant ,
    J-s
13 A = sqrt(h_cross/(mu*omega));    // Amplitude of
    vibrations , m
14 printf("\nThe amplitude of vibrations = %7.5f nm", A
    /1e-09);
15
16 // Result

```

```
17 // The effective force constant of CO molecule =  
    1.85e+003 N/m  
18 // The amplitude of vibrations = 0.00479 nm
```

---

# Chapter 12

## The Solid State

Scilab code Exa 12.1 Classical free electron model

```
1 // Scilab code Ex12.1: Pg 418 (2005)
2 clc; clear;
3 // Part (a)
4 k_B = 1.38e-23; // Boltzmann constat, J/K
5 m_e = 9.11e-31; // Mass of electron, kg
6 T = 300; // Temperature, K
7 N_A = 6.023e+023; // Avogadro's number
8 v_rms = sqrt((3*k_B*T)/m_e); // Root mean
    square velocity of electrons, m/s
9 I = 10; // Electric current, A
10 A = 4e-06; // Area of cross-section of copper
    wire, m^2
11 J = I/A; // Current density, A-m^(-2)
12 d = 8.96; // Density of copper at room
    temperature, g/cc
13 M = 63.5; // Atomic mass of Cu, g
14 n = d*N_A/M*1e+06; // Number of electrons per
    metre cube
15 e = 1.6e-19; // Charge on electron, C
16 v_d = J/(n*e); // Drift velocity, m/s
17 v_d_rms = v_d/v_rms; // Ratio of drift speed to
```

```

    rms speed
18 printf("\nThe ratio of drift speed to rms speed is =
    %3.1e", v_d_rms);
19
20 // Part (b)
21 L = 2.6e-10;
22 tau = L/v_rms;    // Average time between two
    collisions , s
23 printf("\nAverage time between two collisions = %2.2
    e s", tau);
24
25 // Part (c)
26 sigma = (n*e^2*L)/sqrt(3*k_B*T*m_e);    //
    Conductivity of copper , per ohm-m
27 printf("\nConductivity of copper at room temperature
    = %3.1e per ohm-m", sigma);
28
29
30 // Result
31 // The ratio of drift speed to rms speed is = 1.6e
    -009
32 // Average time between two collisions = 2.23e-015 s
33 // Conductivity of copper at room temperature = 5.3e
    +006 per ohm-m

```

---

### Scilab code Exa 12.2 Conduction in diamond

```

1 // Scilab code Ex12.2: Pg 429 (2005)
2 clc; clear;
3 V = 7;    // Energy gap, V
4 L = 5e-08;    // Mean free path , m
5 E = V/L;    // Electric field , V/m
6 printf("\nThe electric field strength required to
    produce conduction in diamond = %3.1fe+08 V/m", E
    *1e-08);

```

```

7 printf("\n");
8
9 // Result
10 // The electric field strength required to produce
    conduction in diamond = 1.4e+08 V/m

```

---

### Scilab code Exa 12.3 Forward and reverse currents in diode

```

1 // Scilab code Ex12.3: Pg 436 (2005)
2 clc; clear;
3 e_V = 1; // Energy applied to diode, eV
4 k_B_T = 0.025; // Product of Boltzmann constant
    and temperature, eV
5 // For simplicity let (q*V)/(k_B*T) = x
6 x = (e_V/(k_B_T));
7 I_f_r = (exp(x)-1)/(exp(-x)-1); // Ratio of
    forward current to reverse current in diode
8 printf("\nThe ratio of forward current to reverse
    current in diode = %3.1fe+17", I_f_r*1e-17);
9
10 // Result
11 // The ratio of forward current to reverse current
    in diode = -2.4e+17

```

---

# Chapter 13

## Nuclear Structure

### Scilab code Exa 13.1 The Atomic Mass Unit

```
1 // Scilab code Ex13.1: Pg 466 (2005)
2 clc; clear;
3 M = 0.012; // Atomic mass of carbon , kg
4 N_A = 6.02e+023; // Avogadro's number
5 m = M/N_A; // Mass of one Carbon-12 atom, kg
6 // As m = 12*u, twelve mass units, solving for u
7 u = m/12; // The atomic mass unit, kg
8 printf("\\nThe atomic mass unit = %4.2e kg", u);
9
10 // Result
11 // The atomic mass unit = 1.66e-27 kg
```

---

### Scilab code Exa 13.2 The Volume and Density of Nucleus

```
1 // Scilab code Ex13.2: Pg 468 (2005)
2 clc; clear;
3 r0 = 1.2e-015; // Nuclear mean radius, m
4 m = 1.67e-027; // Mass of the nucleon, kg
```

```

5 rho_0 = 3*m/(4*pi*r0^3); // Density of the
    nucleus, kg per metre cube
6 printf("\nThe mass of the nucleus = Am approx.");
7 printf("\nThe volume of the nucleus = 4/3*pi*r0^3*A"
    );
8 printf("\nThe density of the nucleus = %3.1e kg per
    metre cube", rho_0);
9
10 // Result
11 // The mass of the nucleus = Am approx.
12 // The volume of the nucleus = 4/3*pi*r0^3*A
13 // The density of the nucleus = 2.3e+17 kg per metre
    cube

```

---

### Scilab code Exa 13.3 Binding energy of the Deuteron

```

1 // Scilab code Ex13.3: Pg 473 (2005)
2 clc; clear;
3 M2 = 2.014102; // Atomic mass of deuteron, u
4 M_H = 1.007825; // Atomic mass of hydrogen, u
5 m_n = 1.008665; // Mass of a neutron, u
6 E_b = (M_H + m_n - M2)*931.494; // Binding
    energy of the deuteron, MeV/u
7 printf("\nThe binding energy of the Deuteron = %5.3 f
    MeV", E_b);
8
9 // Result
10 // The binding energy of the Deuteron = 2.224 MeV

```

---

### Scilab code Exa 13.4 Left out sample during radioactive decay

```

1 // Scilab code Ex13.4: Pg 482 (2005)
2 clc; clear;

```



```

3 T = 5730;    // Half life of the carbon-14 isotope ,
   years
4 N0 = 1000;  // Initial number of carbon-14 isotope
5 t = 22920;  // Time of decay, years
6 n = t/T;    // Total number of half lives
7 N = (1/2)^n*N0; // Sample remains after 22920 years
8 printf("\nNumber of C-14 isotopes remained after %d
   years = %d", t, N);
9
10 // Result
11 // Number of C-14 isotopes remained after 22920
   years = 62

```

---

### Scilab code Exa 13.5 The Activity of Radium

```

1 // Scilab code Ex13.5: Pg 483 (2005)
2 clc; clear;
3 T_half = 1.6e+03*3.16e+07;    // Half life of
   radioactive nucleus Ra-226, s
4 lambda = 0.693/T_half; // Decay constant of Ra-226,
   per second
5 N0 = 3.0e+016; // Number of radioactive nuclei at t
   = 0
6 R0 = lambda*N0; // Activity of sample at t = 0,
   decays/s
7 t = 2.0e+003*3.16e+07; // Time during which the
   radioactive disintegration takes place, s
8 R = R0*exp(-1*lambda*t); // Decay rate after 2.0e
   +003 years, decay/s
9 printf("\nThe decay constant of Ra-226 = %3.1e per
   second", lambda);
10 printf("\nThe activity of sample at t = 0 = %4.1f
   micro-Ci", R0/(3.7e+010*1e-006));
11 printf("\nThe activity of sample after %3.1e years =
   %3.1e decays/s", t, R);

```

```

12
13 // Result
14 // The decay constant of Ra-226 = 1.4e-11 per second
15 // The activity of sample at t = 0 = 11.1 micro-Ci
16 // The activity of sample after 6.3e+10 years = 1.7e
    +05 decays/s

```

---

### Scilab code Exa 13.6 The Activity of Carbon

```

1 // Scilab code Ex13.6: Pg 483 (2005)
2 clc; clear;
3 M = 11.0; // Atomic mass of C-11 isotope , g
4 NA = 6.02e+023; // Avogadro's number
5 m = 3.50e-06; // Given mass of Carbon-11, g
6
7 // Part (a)
8 N = m/M*NA; // Number of C-11 atoms in 3.50
    micro-g of sample
9 printf("\\nThe number of C-11 atoms in %4.2f micro-g
    of sample = %4.2e nuclei", m/1e-06, N);
10
11 // Part (b)
12 T_half = 20.4*60; // Half life of radioactive
    nucleus C-11, s
13 lambda = 0.693/T_half; // Decay constant of C-11,
    per second
14 R0 = lambda*N; // Activity of sample at t = 0,
    decays/s
15 t = 8.00*60*60; // Time during which the
    radioactive disintegration takes place, s
16 R = R0*exp(-1*lambda*t); // Decay rate after 2.0e
    +003 years , decay/s
17
18 printf("\\nThe activity of C-11 sample at t = 0 is %4
    .2e decays/s", R0);

```

```

19 printf("\nThe activity of sample after %4.2f hours =
      %4.2e decays/s", t/3600, R);
20
21 // Result
22 // The number of C-11 atoms in 3.50 micro-g of
      sample = 1.92e+17 nuclei
23 // The activity of C-11 sample at t = 0 is 1.08e+14
      decays/s
24 // The activity of sample after 8.00 hours = 8.99e
      +06 decays/s

```

---

#### Scilab code Exa 13.7 The Radiative Isotope of Iodine

```

1 // Scilab code Ex13.7: Pg 484 (2005)
2 clc; clear;
3 R0 = 5; // Activity of I-131 isotope at the time of
      shipment, mCi
4 R = 4.2; // Activity of I-131 isotope at the time
      of receipt by the medical laboratory, mCi
5 T_half = 8.04; // Half life of radioactive
      nucleus I-131, days
6 lambda = 0.693/T_half; // Decay constant of C-11,
      per second
7 // As  $\log(R/R_0) = -\lambda t$ , solving for t
8 t = -1/lambda*log(R/R0); // Time that has elapsed
      between two measurements, days
9 printf("\nThe time that has elapsed between two
      measurements = %4.2f days", t);
10
11 // Result
12 // The time that has elapsed between two
      measurements = 2.02 days

```

---

### Scilab code Exa 13.8 Energy Liberated during Decay of Radium

```
1 // Scilab code Ex13.8: Pg 486 (2005)
2 clc; clear;
3 M_X = 226.025406; // Atomic mass of Ra-226, u
4 M_Y = 222.017574; // Atomic mass of Rn-222, u
5 M_alpha = 4.002603; // Mass of alpha particle, u
6 Q = (M_X - M_Y - M_alpha)*931.494; // Q-value for
   Radium Decay, MeV/u
7 printf("\nThe Q-value for Radium Decay = %4.2f MeV",
   Q);
8
9 // Result
10 // The Q-value for Radium Decay = 4.87 MeV
```

---

### Scilab code Exa 13.9 Probability of Alpha Decay

```
1 // Scilab code Ex13.9: Pg 487 (2005)
2 clc; clear;
3 Z = 86; // Atomic number of radon
4 A = 222; // Mass number of radon
5 k = 9e+09; // Coulomb constant, N-metre square per
   C-square
6 e = 1.6e-019; // Charge on an electron, C
7 r0 = 7.25e-015; // Bohr radius for alpha particle, m
8 E0 = k*e^2/(2*r0*1e+06*e); // Rydberg energy, MeV
9 R = 1.2e-015*A^(1/3); // Radius of radon nucleus,
   fm
10 E = 5; // Disintegration energy during alpha decay,
   MeV
11 T_E = exp(-4*pi*Z*sqrt(E0/E)+8*sqrt(Z*R/r0)); //
   Decay probability for alpha disintegration
12 printf("\nThe decay probability for alpha
   disintegration at %d MeV energy = %4.2e", E, T_E)
   ;
```

```
13
14 // Result
15 // The decay probability for alpha disintegration at
    5 MeV energy = 1.29e-34
```

---

### Scilab code Exa 13.11 Radioactive Dating

```
1 // Scilab code Ex13.11: Pg 490 (2005)
2 clc; clear;
3 T_half = 5370*3.6e+07; // Half life of C-14, s
4 lambda = 0.693/T_half; // // Decay constant for C
    -14 disintegration , per sec
5 N_C12 = 6.02e+023/12*25; // Number of C-12 nuclei
    in 25.0 g of carbon
6 N0_C14 = 1.3e-012*N_C12; // Number of C-14 nuclei
    in 25.0 g of carbon before decay
7 R0 = N0_C14*3.83e-012*60; // Initial activity of the
    sample , decays/min
8 R = 250; // Present activity of the sample
9 // As  $R = R0 \cdot \exp(-\lambda \cdot t)$ , solving for t
10 t = -1/lambda*log(R/R0); // Time during which the
    tree dies , s
11 printf("\\nThe lifetime of the tree = %3.1e yr", t
    /(365*24*60*60));
12
13 // Result
14 // The lifetime of the tree = 3.6e+03 yr
```

---

# Chapter 14

## Nuclear Physics Applications

Scilab code Exa 14.1 Energy released in Fission

```
1 // Scilab code Ex14.1: Pg 513 (2005)
2 clc; clear;
3 // Part (a)
4 u = 931.5; // Atomic mass unit, MeV
5 M_Li = 7.016003; // Mass of Lithium, kg
6 M_H = 1.007825; // Mass of Hydrogen, kg
7 M_He = 4.002603; // Mass of Helium, kg
8 Q = (M_Li + M_H - 2*M_He)*u; // Q-value of the
   reaction, MeV
9 // Part (b)
10 K_incident = 0.6; // Kinetic energy of the
   incident protons, MeV
11 K_products = Q + K_incident; // Kinetic energy of
   the products
12 printf("\\nThe Q value of the reaction = %4.1f MeV",
   Q);
13 printf("\\nThe kinetic energy of the products (two
   alpha particles) = %4.1f MeV", K_products);
14
15 // Result
16 // The Q value of the reaction = 17.3 MeV
```

```
17 // The kinetic energy of the products (two alpha
    particles) = 17.9 MeV
```

---

#### Scilab code Exa 14.2 Neutron capture by Al

```
1 // Scilab code Ex14.2: Pg 509 (2005)
2 clc; clear;
3 roh = 2.7e+06; // Density of Al, g/cm^3
4 A = 27; // Mass number of Al
5 n = (6.02e+23*roh)/A; // Number of nuclei/m^3
6 sigma = 2.0e-31; // Effective area of nucleus
    normal to motion, m^2
7 R_0 = 5.0e+12; // Rate of incident particles
    per unit area, neutrons/cm^2-s
8 x = 0.30e-03; // Thickness of foil, m
9 R = (R_0*sigma*n*x) // Number of neutrons captured
    by foil, neutrons/cm^2-s
10 printf("\nThe number of neutrons captured by foil =
    %3.1fe+07 neutrons/Sq.cm-s", R*1e-07);
11
12 // Result
13 // The number of neutrons captured by foil = 1.8e+07
    neutrons/Sq.cm-s
```

---

#### Scilab code Exa 14.4 Energy released in the Fission of U235

```
1 // Scilab code Ex14.4: Pg 513 (2005)
2 clc; clear;
3 m = 1; // Mass of Uranium taken, kg
4 Q = 208; // Disintegration energy per event, MeV
5 A = 235; // Mass number of Uranium
6 N = (6.02e+23*m)/A; // Number of nuclei
7 E = N*Q; // Disintegration energy, MeV
```

```

8 printf("\n\nThe total energy released if %1d kg of
   Uranium undergoes fission = %4.2fe+26 MeV", m, E
   *1e-23);
9
10 // Result
11 // The total energy released if 1 kg of Uranium
   undergoes fission = 5.33e+26 MeV

```

---

#### Scilab code Exa 14.5 A Rough Mechanism for Fission Process

```

1 // Scilab code Ex14.5: Pg 513 (2005)
2 clc; clear;
3 A_Ba = 141; // Mass number of Barium
4 A_Kr = 92; // Mass number of Barium
5 r_0 = 1.2e-15; // Separation constant, m
6 r_Ba = r_0*A_Ba^(1/3); // Nuclear radius of Barium
   , m
7 r_Kr = r_0*A_Kr^(1/3); // Nuclear radius of
   Krypton, m
8 r = r_Ba + r_Kr; // Separation between two atoms,
   m
9 Z_1 = 56; // Atomic number of Barium
10 Z_2 = 36; // Atomic number of Barium
11 k = 1.440e-09; // Coulomb constant, eV-m
12 U = k*Z_1*Z_2/r // Coulomb Potential
   energy of two charges, MeV
13 printf("\n\nThe Coulomb potential energy for two
   charges = %3d MeV" , U/1e+06);
14 printf("\n\nThis shows that the fission mechanism is
   plausible");
15
16 // Result
17 // The Coulomb potential energy for two charges =
   248 MeV
18 // This shows that the fission mechanism is

```



plausible

---

### Scilab code Exa 14.6 The Fusion of Two Deutrons

```
1 // Scilab code Ex14.6: Pg 519 (2005)
2 clc; clear;
3 // Part (a)
4 e = 1.6e-19; // Charge on electron , C
5 k = 8.99e-09; // Coulomb constant , N-m^2/C^2
6 r = 1.0e-14; // Distance between two duetrons , m
7 // We have  $U = (k*q1*q2)/r$ , for duetrons  $q1 = q2 = e$ 
  , therefore we get
8 U = (k*e^2)/r; // Potential energy of duetrons , J
9 E_C = 1.1e-014; // The coulomb energy per deuteron
  , J
10 k_B = 1.38e-023; // Boltzmann constant , J/mol/K
11 T = 2/3*E_C/k_B; // Effective temperature
  required for deuteron to overcome the potential
  barrier , K
12 printf("\\nThe potential energy of two duetrons
  separated by the distance of %1.0de-14 m = %4.2 f
  MeV", r*1e+14, (U*1e+12)/e);
13 printf("\\nThe effective temperature required for
  deuteron to overcome the potential barrier = %3.1e
  K", T);
14
15 // Result
16 // The potential energy of two duetrons separated by
  the distance of 1e-14 m = 0.14 MeV
17 // The effective temperature required for deuteron to
  overcome the potential barrier = 5.3e+008 K
18 // Result
19 // The potential energy of two duetrons separated by
  the distance of 1e-14 m = 0.14 MeV
```

---

### Scilab code Exa 14.7 Half value thickness

```
1 // Scilab code Ex14.7: Pg 530 (2005)
2 clc; clear;
3 mew = 55e-02; // Linear absorption coefficient ,
    per m
4 // In equation  $I(x) = I_0 \exp(-mew*x)$ , replacing  $I(x)$ 
    by  $I_0/2$  & solving for  $x$ , we get
5 x = log(2)/mew; // Half value thickness , m
6 printf("\\nThe half value thickness for lead = %4.2fe
    -02 cm", x);
7
8 // Result
9 // The half value thickness for lead = %1.26e-02 cm
```

---

# Chapter 15

## Elementary Particle

Scilab code Exa 15.2 Checking Baryon Numbers

```
1 // Scilab code Ex15.2: Pg 560 (2005)
2 clc; clear;
3 // Data for Reaction 1
4 R1 = cell(6,2); // Declare a 6X2 cell
5 R1(1,1).entries = 'p';
6 R1(2,1).entries = 'n';
7 R1(3,1).entries = 'p';
8 R1(4,1).entries = 'p';
9 R1(5,1).entries = 'n';
10 R1(6,1).entries = 'p_bar';
11 R1(1,2).entries = 1;
12 R1(2,2).entries = 1;
13 R1(3,2).entries = 1;
14 R1(4,2).entries = 1;
15 R1(5,2).entries = 1;
16 R1(6,2).entries = -1;
17 // Data for reaction 2
18 R2 = cell(5,2); // Declare a 5X2 cell
19 R2(1,1).entries = 'p';
20 R2(2,1).entries = 'n';
21 R2(3,1).entries = 'p';
```

```

22 R2(4,1).entries = 'p';
23 R2(5,1).entries = 'p_bar';
24 R2(1,2).entries = 1;
25 R2(2,2).entries = 1;
26 R2(3,2).entries = 1;
27 R2(4,2).entries = 1;
28 R2(5,2).entries = -1;
29 // Check baryon number conservation for first
    reaction
30 if (R1(1,2).entries+R1(2,2).entries) == (R1(3,2).
    entries+R1(4,2).entries+R1(5,2).entries+R1(6,2).
    entries) then
31     printf("\nThe reaction %s + %s --> %s + %s + %s
        + %s can occur (B is conserved)", R1(1,1).
        entries, R1(2,1).entries, R1(3,1).entries, R1
        (4,1).entries, R1(5,1).entries, R1(6,1).
        entries);
32 else
33     printf("\nThe reaction %s + %s --> %s + %s + %s
        + %s cannot occur (B is not conserved)", R1
        (1,1).entries, R1(2,1).entries, R1(3,1).
        entries, R1(4,1).entries, R1(5,1).entries, R1
        (6,1).entries);
34 end
35 // Check baryon number conservation for second
    reaction
36 if R2(1,2).entries+R2(2,2).entries == R2(3,2).
    entries+R2(4,2).entries+R2(5,2).entries then
37     printf("\nThe reaction %s + %s --> %s + %s + %s
        can occur (B is conserved)", R2(1,1).entries,
        R2(2,1).entries, R2(3,1).entries, R2(4,1).
        entries, R2(5,1).entries);
38 else
39     printf("\nThe reaction %s + %s --> %s + %s + %s
        cannot occur (B is not conserved)", R2(1,1).
        entries, R2(2,1).entries, R2(3,1).entries, R2
        (4,1).entries, R2(5,1).entries);
40 end

```

```

41
42 // Result
43 // The reaction  $p + n \rightarrow p + p + n + p_{\text{bar}}$  can
    occur (B is conserved)
44 // The reaction  $p + n \rightarrow p + p + p_{\text{bar}}$  cannot occur
    (B is not conserved)

```

---

### Scilab code Exa 15.3 Checking Lepton Numbers

```

1 // Scilab code Ex15.3: Pg 561 (2005)
2 clc; clear;
3 // Data for Reaction 1
4 R1 = cell(4,3); // Declare a 4X3 cell
5 R1(1,1).entries = 'mu';
6 R1(2,1).entries = 'e-';
7 R1(3,1).entries = 'nue_bar';
8 R1(4,1).entries = 'nu_mu';
9 R1(1,2).entries = 1; // Muon number for mu
10 R1(2,2).entries = 0; // Muon number for e-
11 R1(3,2).entries = 0; // Muon number for nue_bar
12 R1(4,2).entries = 1; // Muon number for nu_mu
13 R1(1,3).entries = 0; // Lepton number for mu
14 R1(2,3).entries = 1; // Lepton number for e-
15 R1(3,3).entries = -1; // Lepton number for
    nue_bar
16 R1(4,3).entries = 0; // Lepton number for nu_mu
17 // Data for Reaction 2
18 R2 = cell(4,3); // Declare a 4X3 cell
19 R2(1,1).entries = 'Pi+';
20 R2(2,1).entries = 'mu+';
21 R2(3,1).entries = 'nu_mu';
22 R2(4,1).entries = 'nu_e';
23 R2(1,2).entries = 0; // Muon number for Pi+
24 R2(2,2).entries = -1; // Muon number for mu+
25 R2(3,2).entries = 1; // Muon number for nu_mu

```

```

26 R2(4,2).entries = 0;      // Muon number for nu_e
27 R2(1,3).entries = 0;      // Lepton number for Pi+
28 R2(2,3).entries = 0;      // Lepton number for mu+
29 R2(3,3).entries = 0;      // Lepton number for nu_mu
30 R2(4,3).entries = 1;      // Lepton number for nu_e
31 // Check lepton number conservation for first
    reaction
32 if (R1(1,2).entries== R1(2,2).entries+R1(3,2).
    entries+R1(4,2).entries) & (R1(1,3).entries == R1
    (2,3).entries+R1(3,3).entries+R1(4,3).entries)
    then
33     printf("\nThe reaction %s --> %s + %s + %s can
        occur (Both L_mu and L_e are conserved)", R1
        (1,1).entries, R1(2,1).entries, R1(3,1).
        entries, R1(4,1).entries);
34 else
35     printf("\nThe reaction %s + %s --> %s + %s + %s
        + %s cannot occur (L_mu and L_e are not
        conserved)", R1(1,1).entries, R1(2,1).entries
        , R1(3,1).entries, R1(4,1).entries);
36 end
37 // Check lepton number conservation for second
    reaction
38 if (R2(1,2).entries== R2(2,2).entries+R2(3,2).
    entries+R2(4,2).entries) & (R2(1,3).entries == R2
    (2,3).entries+R2(3,3).entries+R2(4,3).entries)
    then
39     printf("\nThe reaction %s --> %s + %s + %s can
        occur (Both L_mu and L_e are conserved)", R2
        (1,1).entries, R2(2,1).entries, R2(3,1).
        entries, R2(4,1).entries);
40 else
41     printf("\nThe reaction %s --> %s + %s + %s
        cannot occur (L_mu is conserved but L_e is
        not conserved)", R2(1,1).entries, R2(2,1).
        entries, R2(3,1).entries, R2(4,1).entries);
42 end
43

```

```

44 // Result
45 // The reaction  $\mu^- \rightarrow e^- + \bar{\nu}_e + \nu_\mu$  can
    occur (Both  $L_\mu$  and  $L_e$  are conserved)
46 // The reaction  $\pi^+ \rightarrow \mu^+ + \nu_\mu + \nu_e$  cannot
    occur ( $L_\mu$  is conserved but  $L_e$  is not
    conserved)

```

---

#### Scilab code Exa 15.4 Conservation of strangeness

```

1 // Scilab code Ex15.4: Pg 563 (2005)
2 clc; clear;
3 // Data for Reaction 1
4 R1 = cell(4,2); // Declare a 4X2 cell
5 R1(1,1).entries = 'Pi0';
6 R1(2,1).entries = 'n';
7 R1(3,1).entries = 'K+';
8 R1(4,1).entries = 'sigma-';
9 R1(1,2).entries = 0; // Strangeness number for
    Pi0
10 R1(2,2).entries = 0; // Strangeness number for n
11 R1(3,2).entries = 1; // Strangeness number for K+
12 R1(4,2).entries = -1; // Strangeness number for
    sigma-
13 // Data for Reaction 2
14 R2 = cell(4,2); // Declare a 4X2 cell
15 R2(1,1).entries = 'Pi-';
16 R2(2,1).entries = 'p';
17 R2(3,1).entries = 'pi-';
18 R2(4,1).entries = 'sigma+';
19 R2(1,2).entries = 0; // Strangeness number for Pi
    -
20 R2(2,2).entries = -1; // Strangeness number for p
21 R2(3,2).entries = 1; // Strangeness number for pi
    -
22 R2(4,2).entries = 0; // Strangeness number for

```

```

sigma+
23 // Check strangeness number conservation for first
    reaction
24 if R1(1,2).entries + R1(2,2).entries == R1(3,2).
    entries+R1(4,2).entries then
25     printf("\nThe reaction %s + %s --> %s + %s can
        occur (Strangness is conserved)", R1(1,1).
        entries, R1(2,1).entries, R1(3,1).entries, R1
        (4,1).entries);
26 else
27     printf("\nThe reaction %s + %s --> %s + %s
        cannot occur (Strangness is not conserved)",
        R1(1,1).entries, R1(2,1).entries, R1(3,1).
        entries, R1(4,1).entries);
28 end
29 // Check strangeness number conservation for second
    reaction
30 if R2(1,2).entries + R2(2,2).entries == R2(3,2).
    entries+R2(4,2).entries then
31     printf("\nThe reaction %s + %s --> %s + %s can
        occur (Strangness is conserved)", R2(1,1).
        entries, R2(2,1).entries, R2(3,1).entries, R2
        (4,1).entries);
32 else
33     printf("\nThe reaction %s + %s --> %s + %s
        cannot occur (Strangness is not conserved)",
        R2(1,1).entries, R2(2,1).entries, R2(3,1).
        entries, R2(4,1).entries);
34 end
35
36 // Result
37 // The reaction  $\text{Pi}^0 + n \rightarrow \text{K}^+ + \text{sigma}^-$  can occur (
    Strangness is conserved)
38 // The reaction  $\text{Pi}^- + p \rightarrow \text{Pi}^- + \text{sigma}^+$  cannot
    occur (Strangness is not conserved)

```

---



### Scilab code Exa 15.5 Making virtual particle real

```
1 // Scilab code Ex15.5: Pg 570 (2005)
2 clc; clear;
3 m_pi = 135; // Mass of pion, MeV/c^2
4 m_p = 938.3; // Mass of proton, MeV/c^2
5 // For simplification, let velocity of light be
  unity
6 c = 1; // Velocity of light, m/s
7 // Simplifying  $K_{th} = (m_3 + m_4 + m_5 + \dots)^2 * c^2 - (m_1 + m_2)^2 * c^2$ , we get
8 K_th = 2*m_pi*c^(2) + ((m_pi*c)^2/(2*m_p)); //
  Required kinetic energy of proton, MeV
9 printf("\nRequired kinetic energy of proton = %3d
  MeV", ceil(K_th));
10
11 // Result
12 // Required kinetic energy of proton = 280 MeV
```

---

# Chapter 16

## Cosmology

Scilab code Exa 16.1 Hubbles law

```
1 // Scilab code Ex16.1: Pg 15 (2005)
2 clc; clear;
3 c = 3e+05; // Velocity of light , km/s
4 v = c/4; // Recessional velocity , km/s
5 H_0 = 20e-06; // Hubble's constant , km/s/
   lightyear
6 // From Hubble's law,  $v = H_0 \cdot R_{\max}$ , solving for
   R_max
7 R_max = v/H_0; // Maximum distance at which Hubble's
   law applies without relativistic correction ,
   lightyears
8 printf("\\nThe maximum distance at which Hubbles law
   applies without relativistic correction = %1.0e
   ly", R_max);
9 printf("\\n");
10
11 // Result
12 // The maximum distance at which Hubbles law applies
   without relativistic correction = 4e+09 ly
```

---

### Scilab code Exa 16.2 Critical density of universe

```
1 // Scilab code Ex16.2: Pg 22 (2005)
2 clc; clear;
3 H = 23e-03/(9.46e15); // Hubble's constant, km/s/
  ly
4 G = 6.67e-11; // Gravitational constant, N-m^2/
  kg^2
5 // Since  $H^2 = (8*\%pi*G*p_c)/3$ , solving for p_c
6 p_c = (3*H^2)/(8*%pi*G); // Critical mass density
  of universe, kg/m^3
7 printf("\\nCritical mass density of universe = %4.2e
  kg per metre cube", p_c);
8
9
10 // Result
11 // Critical mass density of universe = %1.06e-27 kg/
  m^3
```

---